1	Main text: 3161
2	Abstract: 149
3	Number of tables: 5
4	Number of figures: 4
5	Number of references: 48
6	Supplementary Tables: 1
7	
8	GENOME-WIDE SPERM DNA METHYLATION CHANGES AFTER THREE
9	MONTHS OF EXERCISE TRAINING IN HUMANS
10	
11	Joshua Denham ¹ , Brendan J. O'Brien ² , Jack T. Harvey ² , Fadi J. Charchar ^{1*}
12	
13	¹ Faculty of Science and Technology, Federation University Australia, Mt Helen,
14	Victoria, Australia.
15	² Faculty of Health, Federation University Australia, Mt Helen, Victoria, Australia.
16	
17	*Corresponding author. Prof Fadi Charchar, Y Building, University Drive, Federation
18	University Australia, Mt Helen, Victoria, Australia 3350. Ph: +613 5327 6098, Fax:
19	+613 5327 9602, Email: f.charchar@federation.edu.au
20	
21	Keywords: Epigenetics, sperm, transgenerational inheritance, disease risk
22	
23	
24	
25	

Abstract Aim: DNA methylation programs gene expression and is involved in numerous biological processes. Accumulating evidence supports transgenerational inheritance of DNA methylation changes in mammals via germ cells. Our aim was to determine the effect of exercise on sperm DNA methylation. Materials and Methods: Twenty-four men were recruited and assigned to an exercise intervention or control group. Clinical parameters were measured and sperm samples were donated by subjects before and after the three month time-period. Mature sperm global and genome-wide DNA methylation was assessed using an ELISA assay and the 450K BeadChip (Illumina). Results: Global and genome-wide sperm DNA methylation was altered after three months of exercise training. DNA methylation changes occurred in genes related to numerous diseases such as Schizophrenia and Parkinson's disease. **Conclusions:** Our study provides the first evidence showing exercise training reprograms the sperm methylome. Whether these DNA methylation changes are inherited to future generations warrants attention.

Executive summary

52 *Aim*

51

58

59

60

61

62

63

64

65

66

68

69

70

71

72

73

- To determine the impact of exercise training on human sperm DNA
 methylation.
- 55 Materials and Methods
- Twenty-four healthy young men were recruited and allocated to an exercise
 training or control group.
 - Health and fitness parameters were measured and sperm samples were collected before and after three months.
 - Global and genome-wide DNA methylation was quantified using an ELISA assay and the Infinium HumanMethylation450 BeadChip (Illumina), respectively.
 - Significant alterations in CpG methylation and corresponding gene pathways
 modulated by exercise training were analysed using Partek (Genomic Suite).
 The disease association was performed using the Database for Annotation,
 Visualization and Integrated Discovery (DAVID).
- 67 Discussion
 - Exercise training was associated with global demethylation of human sperm.
 - DNA methylation changes occurred across the genome after exercise training in genes enriched for numerous developmental processes and pathways associated with the adaptation to exercise training.
 - Genes related to disease were methylated after exercise, indicating transcriptional silencing.

74

Conclusions

 We provide the first evidence demonstrating exercise training regulates the human sperm methylome at CpG sites in genes associated with disease.

79

80

78

76

77

Introduction

81 DNA methylation is crucial for long-term gene expression regulation [1, 2]. DNA 82 methylation is, however, reprogrammable through environmental conditions [3, 4]. Aberrant DNA methylation signatures are associated with age-related diseases such 83 84 as cancer [5-7], atherosclerosis [8, 9], type 2 diabetes mellitus [10, 11], Alzheimer's disease [12] and major psychosis disorders [13, 14]. 85 Mounting evidence indicates that environmentally induced DNA methylation changes 86 are inherited by subsequent generations and affects their phenotypes [15-22]. A 87 likely mechanism for inheritance of DNA methylation profile is via the germ cells. 88 89 Interestingly, aging is associated with sperm DNA methylation changes and may 90 influence disease risk of future generations [23, 24]. The differentially methylated regions in sperm obtained from old (12–14 month-old) versus young (3-month-old) 91 92 mice were observed in the brain of the subsequent offspring, causing altered gene expression and increased behaviors associated with psychiatric disorders 93 (schizophrenia and autism spectrum disorder) [24]. Although aging has been 94 95 investigated in context with sperm DNA methylation, the influence of other favorable environmental stimuli are yet to be determined. 96 97 We recently found that exercise training significantly improves cardiorespiratory fitness and lipid profile in conjunction with genome-wide leukocyte methylation 98 99 changes in young men [25]. Others have shown that exercise training modulates skeletal myocyte [26-28] and adipocyte [29] DNA methylation as well as improving 100

health and fitness phenotypes. Thus, somatic cells are vulnerable to epigenetic changes caused by exercise training, but whether exercise training regulates DNA methylation is germ cells (oocyte or sperm) is currently unknown.

The aim of this study was to determine the impact of exercise training on human sperm DNA methylation. Considering the mounting evidence supporting exercise training as an environmental modulator of somatic cell DNA methylation [25-29], we hypothesized that exercise training would significantly modify sperm DNA methylation in genes related to human diseases.

Materials and methods

Subjects

Twenty-four healthy young men were recruited for this study. Subjects were non-smoking, not taking any medications and free from any age-related chronic diseases, according to self-administered health questionnaires. Subjects were eligible if they had not engaged in any structured high-intensity aerobic exercise training during the past six months and were recruited predominantly from the University by flyers and by word-of-mouth.

The subjects' characteristics are outlined in Table 1. All subjects gave written informed consent to participate and this study was approved by the Federation University Australia Human Research Ethics Committee.

Procedures

On the initial day of testing, subjects were asked to complete health and physical activity questionnaires in order to establish their eligibility to participate in the study. Subjects were also familiarized with the testing procedures to follow and were given a collection tube for their sperm donation.

The second testing session involved a sperm donation, blood donation, physical measures and a test of maximal oxygen uptake (VO_{2max}) on a treadmill. All testing sessions were performed in the morning (7–10 AM). Height, weight and body mass index were recorded using scales and a stadiometer. Blood pressure was measured and augmentation index was assessed by non-invasive applanation tonometry using the SphygmoCor instrument (AtCor, Australia). Subjects completed a maximal treadmill exercise test to assess $\dot{V}O_{2max}$. After a three minute warm-up, subjects began running at 10 km⁻¹ and the treadmill speed was progressively increased by 1 km⁻¹ every second minute until the volitional exhaustion or the supervising Exercise Physiologist deemed it appropriate to stop the test. Participants were fitted with a two-way breathing valve (Hans Rudolph) and breath-by-breath analysis was performed by the online metabolic system (Moxus, Modular, USA). VO_{2max} was determined as the highest O₂ value averaged over 60 s. Maximal treadmill speed was deemed the highest speed successfully completed and was adjusted for time so that participants were encouraged to exercise to their limit. The same testing procedures were completed 3–5 days after the cases' final exercise session or three months after their initial donation for controls. Sperm and blood processing Subjects gave resting preprandial sperm and blood donations before and after the three month exercise intervention. Following an overnight fast, subjects extracted their sperm by masturbating at home and donated their sample to the researchers at the University within 30 min of extraction. All sperm donations and analyses were performed according the World Health Organization's recommendations. Sperm cell number and vitality was assessed by the same researcher using a sheep-counter.

Mature, motile sperm were isolated from lymphocytes, epithelial cells, immature or

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

abnormal sperm and other cellular debris using the PureSperm 40/80 reagents (Nidacon). Briefly, 2 ml of PureSperm 40 was carefully layered on top of PureSperm 80, followed by gently aliquoting 1.5 ml of ejaculate onto the density gradient. The density gradient was spun at 300 G for 20 min. The mature sperm were pelleted and supernatant was removed. The pellet was washed twice by resuspending it in 5 ml of PBS before being spun at 500 G for 10 min. DNA was extracted from mature sperm suspended in 200 µl of PBS using the Purelink Genomic DNA Mini kit (LifeTechnologies) following the manufacturer's recommendations with some small adjustments. To deform the tightly packaged DNA in mature sperm, 0.03 mmol of DL-Dithiothreitol solution (Sigma-Aldrich) was added to the sample before the addition of proteinase K solution and the protein digestion step (incubation at 55°C) was extended from the recommended 10 min to 90 min. All extractions were performed immediately after sperm donation to prevent any temporal de novo influences on DNA methylation. DNA yield was assessed using the Nanodrop 2000 Spectrophotometer (Thermo Scientific) before being stored at -20°C. A preprandial blood sample (24 ml) was drawn from the antecubital vein into serum separating tubes with the subject seated, using standard phlebotomy procedures. Blood was immediately stored on ice before further processing. Blood lipid assays were performed as described previously [25]. Global DNA methylation analysis Average global DNA methylation of sperm from all subjects was quantified using the 5-mC ELISA Kit (Zymo Research, USA). All experiments were performed according to the manufacturer's procedures. All paired samples before and after the intervention were run on the same plate, as well as 100% methylated and 0% methylated controls, all run in duplicate with a standard curve. Absorbance was

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

measured at 450 nm using the Multiskan FC Microplate Photometer (Thermo Scientific, Australia) and average DNA methylation was calculated using the formula generated from the standard curve. The average intra-assay coefficient of variation for duplicate samples was 4.85%. Genome-wide DNA methylation analysis Genome-wide sperm DNA methylation was quantified in 12 cases before and after the exercise intervention using the Infinium HumanMethylation 450K BeadChip (Illumina) as previously described [25]. Experiments were conducted following the manufacturer's guidelines, with the help of the Busselton Population Medical Research Institute. Briefly, raw β -values underwent Subset-quantile Within Array Normalization (SWAN) and were transformed to M-values using the logit function $(\log^2(\beta/(1-\beta)))$ before analyzing differentially methylated CpG sites after exercise training. M-values were converted to β -values for representation of differentially methylated CpG sites in tables and graphs because they correspond to the percentage of CpG methylation and are a good biologically relevant representation. Genome-wide DNA methylation data was analyzed using Partek (Genomic Suite, version 6.6). Exercise training Subjects were randomly allocated to either an exercise training (cases) or control group. Cases completed sprint interval training twice weekly for three months. Three months of exercise training was chosen because it covered one human spermatogenesis cycle. The controls did not engage in any form of training but were asked not to deviate from their routine physical activity habits. All subjects were asked to adhere to their normal diet. The sprint interval training schedule is outlined

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

in Table 2. Sprint interval training is an effective type of exercise training that rapidly improves cardiorespiratory fitness and endurance performance [30-33]. The exercise training protocol of this study was based on research shown to improve cardiorespiratory fitness and endurance performance [25, 34], but with some adjustments, to avoid overuse injuries that could occur with an extended training period (three months). Subjects completed 30 s sprints at maximal intensity with a passive four minute rest period between efforts. Training began with three sprints and was progressed by one effort every fortnight until week 7, when sprints were kept at six efforts for the remaining six weeks of the intervention. Before each session, subjects completed a short warm-up entailing a light 5 min jog, dynamic stretches and some 20 m sprints at 70, 80, 90 and 100% of maximal effort. Training was performed on the University's athletics oval and subjects were supervised and given constant encouragement during sprints by an Exercise Physiologist. Statistical analyses Kolmogorov-Smirnov and Shapiro-Wilks were performed to assess normality. Twotailed paired t-tests were used to assess within-subject changes to physical, health and fitness variables. A repeated measures ANOVA was used to determine the interaction between global sperm DNA methylation before and after the intervention in relation to cases and controls using IBM SPSS for Windows (version 21). Differentially methylated CpG sites in cases were determined by repeated measures ANOVA using Partek (Genomic Suite, version 6.6). In order to control for false positives, a false discovery rate correction (FDR, $q \le 0.1$) was applied to genome-wide data by converting p-values to q-values. Hierarchical clustering, gene ontology and pathway analysis on genome-wide CpG methylation data were performed using Partek (Genomic Suite, version 6.6). Disease annotation analysis was conducted

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

using the Database for Annotation, Visualization and Integrated Discovery (DAVID, version 6.7). Statistical significance was set at p<0.05.

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

225

226

Results

Exercise training improved cardiorespiratory fitness To determine whether exercise training alters human sperm DNA methylation, 13 untrained male volunteers completed three months of exercise training using a modified training protocol (Table 2) previously shown to improve cardiorespiratory fitness and induce genome-wide DNA methylation changes in leukocytes [25]. Although $\dot{V}O_{2max}$ was unchanged (2.2%, p=0.35), relative to baseline, the cases, on average, had a significantly lower resting heart rate by 10 beats min⁻¹ and completed an extra one min at a significantly higher maximal treadmill speed (p=0.01 and p=0.002, respectively), while controls showed no significant changes in these variables (Table 1). To assess whether exercise training influences average sperm DNA methylation, we assessed global sperm DNA methylation in cases and controls using ELISA-based chemistry. After adjusting for ELISA plate, relative to the unchanged global sperm DNA methylation observed in controls (n=11, 4.09%), sperm DNA was significantly de-methylated after three months of exercise training (n=13, -6.63%, interaction p=0.006) (Figure 1A and B). Based on the global DNA methylation results we only quantified genome-wide sperm DNA methylation in 12 cases before and after three months of exercise training using the 450K BeadChip (Illumina). After applying a FDR ($q \le 0.1$), we found DNA methylation changes occurred throughout the genome in relation to the nearest gene and CpG islands (Figure 1C and D). Of these, 7509 CpG sites relating to 4602

genes had significant DNA methylation changes ranging from 0.3 to 6.2% after exercise training (Figure 1E and 2). More CpG sites were demethylated rather than methylated after exercise training (4275 vs 3233, respectively). The top 25 methylated and demethylated CpG sites after exercise training are outlined in Table 3. To gain insight into the biological impact of the CpG methylation changes following exercise, we performed gene ontology and pathway analysis. Genes with altered CpG methylation were those enriched for crucial biological processes including developmental process, anatomical structure, embryonic morphogenesis and organ development (Figure 3 and Supplementary Table 1). Pathway analysis revealed genes with CpG site DNA methylation changes were enriched for pathways such as MAPK signaling, PI3K-Akt signaling, pathways in cancer and ErbB signaling (Table 4). In the search for the most biologically relevant CpG methylation changes, we focused on CpG sites with a ≥1.5% increase (n=703) or decrease (n=1247) in methylation after exercise training and entered them into DAVID for disease annotation analysis. While genes with demethylated CpG sites were those relating to normal variation, those with increased methylation were enriched for numerous diseases, with the most overrepresented diseases being Schizophrenia, Parkinson's Disease, autism, cervical cancer and leukemia (Figure 4). The global erasure of sperm CpG methylation that occurs after fertilization is well established [35-37]. The sperm CpG methylation at imprinted genes, however, are resistant to the global erasure after fertilization [38, 39] and as such, these genes are key candidates for possible transgenerational inheritance through sperm. We found 16 paternally imprinted genes were differentially methylated after the three month

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

exercise training intervention (Table 5), in genes associated with pathologies including EKG abnormalities (*KCNQ10T1* and *KCNQ1*), autism and alcohol dependence (*GABRG3*), Tourette syndrome, obsessive compulsive disorder and dystonia (*SGCE*), and Alzheimer's disease, obesity, type 2 diabetes mellitus, blood pressure and atherosclerosis (*IGF2* and *INS-IGF2*).

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

275

276

277

278

279

Discussion

Unlike the genome, the epigenome is malleable to changing environments and these changes are somewhat heritable [3, 4]. Germ cells (sperm and oocytes) are likely vectors that transfer environmentally-affected DNA methylation profiles to future generations. We are the first to show the exercise training-induced DNA methylation changes in human sperm. We found exercise training significantly manipulated genome-wide DNA methylation and most strikingly, increased methylation in disease-associated genes indicating transcriptional silencing. Moreover, methylation changes occurred in paternally imprinted genes that are exempt from the DNA methylation erasure after fertilization [38, 39]. Mounting evidence has demonstrated the capacity of exercise training to influence the human DNA methylome in numerous tissues, including skeletal myocytes [26-28], adipocytes [29] and leukocytes [25]. Exercise training or increased physical activity tends to be associated with a reduced level of somatic cell global DNA methylation. Leukocyte DNA methylation was anti-correlated to levels of physical activity in 509 older adults (>70 y) [40] and we previously reported more demethylated leukocyte CpG sites in response to a four week exercise training intervention in young men [25]. Furthermore, global and gene-specific demethylation occurred after acute exercise in human skeletal myocytes [41]. In the present study,

we found global sperm DNA methylation was reduced after three months of exercise training, suggesting that exercise may systemically demethylate both somatic and germ cells. Interestingly, older adults have more heavily methylated global sperm DNA compared to their younger counterparts [23, 42]. Also defective human sperm exhibit nearly two-fold higher levels of global DNA methylation [43]. To that end, the global demethylation observed in our subjects after exercise training could be favorable, possibly counteracting the increased DNA methylation associated with ageing. This, in turn, could positively influence fertility though future studies are necessary. Interestingly, exercise training was associated with sperm methylation changes to CpG sites in genes related to pathways and biological processes similar to those observed in skeletal muscle [26] and leukocytes [25] from previous studies. For example, pathways regulated by exercise training-induced leukocyte DNA methylation alterations [25] that were also observed in sperm include MAPK signaling, PI3K-Akt signaling and pathways in cancer. Similarly, overlapping pathways between skeletal muscle [26] and sperm pathways modulated by exercise training include MAPK signaling, insulin signaling and ErbB signaling. Therefore, it would seem that some exercise-responsive genes (CpG sites) are regulated by DNA methylation in human somatic cells and also in sperm. The genes with CpG methylation changes in the present study were enriched for numerous molecular and cellular processes including anatomical structure development and morphogenesis, single-organism development and embryonic morphogenesis. Genes with exercise-induced CpG methylation changes were also associated with many debilitating diseases such as Schizophrenia, Parkinson's disease, cervical cancer, leukemia and autism. The genes associated with diseases

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

were methylated after exercise training, indicating that these genes may have been transcriptionally silenced. Age-associated sperm CpG methylation patterns were observed in genes implicated in Schizophrenia in humans [23] and mice [24]. Some of the adverse DNA methylation patterns found in sperm of aged mice was transmitted to the brain of their offspring, leading to transcriptional dysregulation of genes involved in schizophrenia and autism, facilitating the disease-related traits [24]. Furthermore, offspring (F1 to F3) of gestating female rats treated with endocrine disruptors (bisphenol-A, bis[2-ethlhexyl]phthalate and dibutyl phthalate and vinclozolin) have altered sperm DNA methylation and an increased risk of testis disease, obesity and infertility [15, 44, 45]. These environmentally-induced DNA methylation aberrations led to transcriptional changes in common pathways across numerous somatic cells, including those regulated in the present study (MAPK signaling, insulin signaling and pathways in cancer) [46]. Whether the altered sperm DNA methylation observed in our subjects will be transmitted to their offspring and prevent the development of disease is speculative and would require large human studies involving the analysis of multiple generations. Germ cells undergo global demethylation following fertilization [35, 39]. Consequently, the likelihood of such CpG methylation changes surviving this demethylation event is uncertain. We did, however, reveal a number of paternally imprinted genes (e.g. IGF2, INS-IGF2, SGCE and GABRG3) had differentially methylated CpGs after exercise training. These genes are exempt from the global demethylation after fertilization [38, 39] and as such, are key candidates for exerciseinduced transgenerational inheritance of DNA methylation gene programming. Our disease analysis using DAVID revealed the differentially methylated paternally imprinted genes after exercise training were related to neurological disorders

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

(autism, Alzheimer's disease, Tourette's syndrome and obsessive compulsive disorder) and cardio-metabolic diseases (obesity, type 2 diabetes mellitus, high blood pressure and atherosclerosis). Whilst no evidence exists, it is tempting to speculate that exercise training may remodel the sperm methylome and that this epigenetic modification may reduce the disease risk of offspring. DNA methyltransferase enzymes (DNMT1, DNMT3A, DNMT3B and DNMT3L) as well as ten-eleven translocation (TET1, TET2 and TET3) proteins could be modulated by exercise training leading to altered DNA methylation. Our study was not designed to target the exact physiological or molecular mechanism by which exercise training influences DNA methylation in sperm, rather it was designed to investigate whether exercise training influences sperm DNA methylation. Therefore, further research is required to identify the molecular mechanisms evoking the changes in the sperm methylome after exercise training. Our study has several limitations. Firstly, our study had a modest sample size and there was a statistically different global sperm DNA methylation between cases and controls at baseline. While we acknowledge this is not desirable, our statistical analysis accounted for the observed difference. We do, however, recommend that our results should be verified in a larger population. The relative exercise workload was not standardized. We did, however, ask that all subjects perform each sprint maximally without pacing to ensure they are working at maximal intensity. Certain diets are associated with DNA methylation levels in somatic cells [47, 48]. Despite asking our subjects not to deviate from their routine dietary habits, the lack of control for diet in our study should also be acknowledged. Finally, we did not control for genotype-DNA methylation interactions or quantify DNA methylation in context with other epigenetic modifications or gene expression. Nonetheless, our novel finding

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

that exercise alters human sperm DNA methylation should encourage future investigations into the influence of exercise training on germ cell epigenetic modifications and gene regulation.

378

379

380

381

382

383

375

376

377

Conclusions

We report for the first time that exercise training significantly modifies the human sperm methylome. Whilst exercise training was associated with an increased global sperm demethylation overall, methylation in key genes associated with human disease was increased indicating transcriptional silencing of these genes.

384

385

References

- 1. Jones PA, Takai D. The role of DNA methylation in mammalian epigenetics.
- 387 *Science* 293(5532), 1068-1070 (2001).
- 388 2. Wu H, Tao J, Sun YE. Regulation and function of mammalian DNA
- methylation patterns: A genomic perspective. Brief. Funct. Genomics 11(3),
- 390 240-250 (2012).
- 391 3. Angers B, Castonguay E, Massicotte R. Environmentally induced phenotypes
- and DNA methylation: How to deal with unpredictable conditions until the next
- 393 generation and after. *Mol. Ecol* 19(7), 1283-1295 (2010).
- 394 4. Flores KB, Wolschin F, Amdam GV. The role of methylation of DNA in
- environmental adaptation. *Integr. Comp. Biol* 53(2), 359-372 (2013).
- 5. Cheung HH, Lee TL, Davis AJ, Taft DH, Rennert OM, Chan WY. Genome-
- wide DNA methylation profiling reveals novel epigenetically regulated genes
- and non-coding rnas in human testicular cancer. Br. J. Cancer 102(2), 419-
- 399 427 (2010).

- 400 6. Kanai Y. Genome-wide DNA methylation profiles in precancerous conditions 401 and cancers. *Cancer Sci* 101(1), 36-45 (2010).
- Weber M, Davies JJ, Wittig D *et al.* Chromosome-wide and promoter-specific
 analyses identify sites of differential DNA methylation in normal and
 transformed human cells. *Nat. Genet* 37(8), 853-862 (2005).
- 405 8. Zaina S, Heyn H, Carmona FJ *et al.* DNA methylation map of human atherosclerosis. *Circ. Cardiovasc. Genet* 7(5), 692-700 (2014).
- 9. Sharma P, Garg G, Kumar A *et al.* Genome wide DNA methylation profiling for epigenetic alteration in coronary artery disease patients. *Gene* 541(1), 31-409 40 (2014).
- Toperoff G, Aran D, Kark JD *et al.* Genome-wide survey reveals predisposing
 diabetes type 2-related DNA methylation variations in human peripheral
 blood. *Hum. Mol. Genet* 21(2), 371-383 (2012).
- Volkmar M, Dedeurwaerder S, Cunha DA *et al.* DNA methylation profiling
 identifies epigenetic dysregulation in pancreatic islets from type 2 diabetic
 patients. *EMBO J* 31(6), 1405-1426 (2012).
- 416 12. Bakulski KM, Dolinoy DC, Sartor MA *et al.* Genome-wide DNA methylation 417 differences between late-onset alzheimer's disease and cognitively normal 418 controls in human frontal cortex. *J. Alzheimers Dis* 29(3), 571-588 (2012).
- Wockner LF, Noble EP, Lawford BR *et al.* Genome-wide DNA methylation
 analysis of human brain tissue from schizophrenia patients. *Transl. Psychiatry* 421
 4, e339 (2014).
- 14. Nishioka M, Bundo M, Koike S *et al.* Comprehensive DNA methylation analysis of peripheral blood cells derived from patients with first-episode schizophrenia. *J. Hum. Genet* 58(2), 91-97 (2013).

- 425 15. Anway MD, Cupp AS, Uzumcu M, Skinner MK. Epigenetic transgenerational
- actions of endocrine disruptors and male fertility. Science 308(5727), 1466-
- 427 1469 (2005).
- 428 16. Manikkam M, Tracey R, Guerrero-Bosagna C, Skinner MK. Plastics derived
- endocrine disruptors (bpa, dehp and dbp) induce epigenetic transgenerational
- inheritance of obesity, reproductive disease and sperm epimutations. *PLoS*
- 431 One 8(1), e55387 (2013).
- 432 17. Manikkam M, Guerrero-Bosagna C, Tracey R, Haque MM, Skinner MK.
- Transgenerational actions of environmental compounds on reproductive
- disease and identification of epigenetic biomarkers of ancestral exposures.
- 435 PLoS One 7(2), e31901 (2012).
- 436 18. Guerrero-Bosagna C, Savenkova M, Hague MM, Nilsson E, Skinner MK.
- Environmentally induced epigenetic transgenerational inheritance of altered
- sertoli cell transcriptome and epigenome: Molecular etiology of male infertility.
- 439 PLoS One 8(3), e59922 (2013).
- 440 19. Michaud EJ, Van Vugt MJ, Bultman SJ, Sweet HO, Davisson MT, Woychik
- RP. Differential expression of a new dominant agouti allele (aiapy) is
- correlated with methylation state and is influenced by parental lineage. *Genes*
- 443 Dev 8(12), 1463-1472 (1994).
- 20. Carone BR, Fauquier L, Habib N et al. Paternally induced transgenerational
- environmental reprogramming of metabolic gene expression in mammals. *Cell*
- 446 143(7), 1084-1096 (2010).
- 21. Cooney CA, Dave AA, Wolff GL. Maternal methyl supplements in mice affect
- epigenetic variation and DNA methylation of offspring. *J. Nutr* 132(8 Suppl),
- 449 2393S-2400S (2002).

- 450 22. Ng SF, Lin RC, Laybutt DR, Barres R, Owens JA, Morris MJ. Chronic high-fat
- diet in fathers programs beta-cell dysfunction in female rat offspring. *Nature*
- 452 467(7318), 963-966 (2010).
- 453 23. Jenkins TG, Aston KI, Pflueger C, Cairns BR, Carrell DT. Age-associated
- sperm DNA methylation alterations: Possible implications in offspring disease
- 455 susceptibility. *PLoS Genet* 10(7), e1004458 (2014).
- 456 24. Milekic MH, Xin Y, O'donnell A et al. Age-related sperm DNA methylation
- changes are transmitted to offspring and associated with abnormal behavior
- and dysregulated gene expression. *Mol. Psychiatry*, (2014).
- 25. Denham J, O'brien BJ, Marques FZ, Charchar FJ. Changes in the leukocyte
- methylome and its effect on cardiovascular related genes after exercise. J.
- 461 Appl. Physiol (1985), JAP 00878 02014 (2014).
- 462 26. Nitert MD, Dayeh T, Volkov P et al. Impact of an exercise intervention on DNA
- methylation in skeletal muscle from first-degree relatives of patients with type
- 464 2 diabetes. *Diabetes* 61(12), 3322-3332 (2012).
- 27. Rowlands DS, Page RA, Sukala WR et al. Multi-omic integrated networks
- connect DNA methylation and mirna with skeletal muscle plasticity to chronic
- exercise in type 2 diabetic obesity. *Physiol. Genomics* 46(20), 747-765
- 468 (2014).
- Lindholm ME, Marabita F, Gomez-Cabrero D et al. An integrative analysis
- reveals coordinated reprogramming of the epigenome and the transcriptome
- in human skeletal muscle after training. *Epigenetics* 9(12), 1557-1569 (2014).
- 29. Ronn T, Volkov P, Davegardh C et al. A six months exercise intervention
- influences the genome-wide DNA methylation pattern in human adipose
- 474 tissue. *PLoS Genet* 9(6), e1003572 (2013).

- 475 30. Khaled MB, A M, S TK, S PC, R D. Effect of traditional aerobic exercises
- versus sprint interval training on pulmonary function tests in young sedentary
- 477 males: A randomised controlled trial. J. Clin. Diagn. Res 7(9), 1890-1893
- 478 (2013).
- 479 31. Macpherson RE, Hazell TJ, Olver TD, Paterson DH, Lemon PW. Run sprint
- interval training improves aerobic performance but not maximal cardiac
- output. Med. Sci. Sports. Exerc 43(1), 115-122 (2011).
- 482 32. Cicioni-Kolsky D, Lorenzen C, Williams MD, Kemp JG. Endurance and sprint
- benefits of high-intensity and supramaximal interval training. Eur. J. Sport. Sci
- 484 13(3), 304-311 (2013).
- 485 33. Sloth M, Sloth D, Overgaard K, Dalgas U. Effects of sprint interval training on
- vo2max and aerobic exercise performance: A systematic review and meta-
- analysis. Scand. J. Med. Sci. Sports 23(6), e341-352 (2013).
- 488 34. Denham J, Feros SA, O'brien BJ. Four weeks of sprint interval training
- improves 5 km run performance. J. Strength. Cond. Res, (2015).
- 490 35. Mayer W, Niveleau A, Walter J, Fundele R, Haaf T. Demethylation of the
- zygotic paternal genome. *Nature* 403(6769), 501-502 (2000).
- 492 36. Kafri T, Gao X, Razin A. Mechanistic aspects of genome-wide demethylation
- in the preimplantation mouse embryo. *Proc. Natl. Acad. Sci USA* 90(22),
- 494 10558-10562 (1993).
- 495 37. Oswald J, Engemann S, Lane N et al. Active demethylation of the paternal
- genome in the mouse zygote. *Curr. Biol* 10(8), 475-478 (2000).
- 497 38. Morgan HD, Santos F, Green K, Dean W, Reik W. Epigenetic reprogramming
- in mammals. *Hum. Mol. Genet* 14 Spec No 1, R47-58 (2005).

- 39. Reik W, Dean W, Walter J. Epigenetic reprogramming in mammalian development. *Science* 293(5532), 1089-1093 (2001).
- 501 40. Luttropp K, Nordfors L, Ekstrom TJ, Lind L. Physical activity is associated with decreased global DNA methylation in swedish older individuals. *Scand. J.*
- 503 Clin. Lab. Invest 73(2), 184-185 (2013).
- 504 41. Barres R, Yan J, Egan B *et al.* Acute exercise remodels promoter methylation 505 in human skeletal muscle. *Cell Metab* 15(3), 405-411 (2012).
- Jenkins TG, Aston KI, Cairns BR, Carrell DT. Paternal aging and associated intraindividual alterations of global sperm 5-methylcytosine and 5-hydroxymethylcytosine levels. *Fertil. Steril* 100(4), 945-951 (2013).
- 509 43. Barzideh J, Scott RJ, Aitken RJ. Analysis of the global methylation status of 510 human spermatozoa and its association with the tendency of these cells to 511 enter apoptosis. *Andrologia* 45(6), 424-429 (2013).
- 512 44. Guerrero-Bosagna C, Settles M, Lucker B, Skinner MK. Epigenetic 513 transgenerational actions of vinclozolin on promoter regions of the sperm 514 epigenome. *PLoS One* 5(9), (2010).
- 515 45. Nilsson EE, Skinner MK. Environmentally induced epigenetic 516 transgenerational inheritance of disease susceptibility. *Transl. Res* 165(1), 12-517 17 (2015).
- 518 46. Skinner MK, Manikkam M, Haque MM, Zhang B, Savenkova MI. Epigenetic 519 transgenerational inheritance of somatic transcriptomes and epigenetic 520 control regions. *Genome Biol* 13(10), R91 (2012).
- Zhang FF, Morabia A, Carroll J *et al.* Dietary patterns are associated with
 levels of global genomic DNA methylation in a cancer-free population. *J. Nutr* 141(6), 1165-1171 (2011).

48. Anderson OS, Sant KE, Dolinoy DC. Nutrition and epigenetics: An interplay of dietary methyl donors, one-carbon metabolism and DNA methylation. *J. Nutr Biochem* 23(8), 853-859 (2012).

Acknowledgements

We would like to thank the subjects for successfully completing such as physically demanding study. JD is supported by an Australian Post-Graduate Award (APA) scholarship. The work was supported by the National Health and Medical Research Council of Australia and LEW CARTY Charitable Foundation.

Competing interests

The authors have no competing interests to disclose.

Figure Legends

Figure 1. Sperm DNA methylation changes after exercise training. Average percent (±SE) of global sperm DNA methylation is given for 13 cases (A) and 11 controls (B) after the three month intervention. Relative to the controls who showed no global methylation change, cases exhibited a global demethylation (-6.63%) after three months of exercise training (mean±SE: 39.1±3.85 to 43.2±3.6 and 52.95±3.56 to 46.3±3.4, respectively, interaction *p*=0.006). Data are from repeated measures ANOVA adjusting for ELISA plate number. Genome-wide DNA methylation changes at CpG sites in relation to the nearest gene (C) and CpG islands (D) are shown. Data are from Chi² tests. E) The number of CpG sites with increased methylation (black bars) and decreased methylation (grey bars) are given on the y-axis with the

CpG sites (*q*≤0.1). 549 Figure 2. Hierarchical clustering of CpG sites after three months of exercise 550 551 training. The CpG methylation status is indicated by blue (low methylation) and yellow (high methylation) from 12 subjects before (grey bar) and after (purple bar) 552 three months of exercise training. Each branch of the array tree on the left of the 553 purple and grey bars are subjects. The array tree at the top of the hierarchical cluster 554 555 indicates clusters of CpG sites modified by exercise training. Data are from differentially methylated CpG sites ($q \le 0.1$). 556 Figure 3. Gene ontology for genes with differentially methylated CpG sites 557 558 after exercise training. Genes with altered CpG site methylation after exercise are 559 grouped for function on the y-axis with corresponding enrichment scores on the xaxis. Data are from differentially methylated CpG sites (*q*≤0.1). 560 Figure 4. Disease annotation analysis for genes with differentially methylated 561 562 **CpG** sites after exercise training. Data are from genes with a ≥1.5% increase or decrease in methylation after exercise training ($q \le 0.1$). The y-axis represents the 563 564 number of genes for each annotated disease (x-axis). 565 566 567 568 569 570

magnitude of change illustrated on the x-axis. Data are from differentially methylated

Table 1. Participant characteristics before and after three months of exercise training.

Variable	Ca	ases	Controls			
	(n	=13)	(n:	=11)		
	Before	After	Before	After		
	Exercise	Exercise				
Age (y)	24.4	1±5.19	22.4	5±4.74		
Ht (cm)	181.	1±6.54	177.9	5±4.07		
Wt (kg)	83.58±10.7	84.01±11.97	82.22±11.83	82.93±10.64		
BMI (Wt/Ht²)	25.53±3.34	25.89±3.61	25.94±3.5	25.9±3.08		
HR (beats min ⁻¹)	78.54±15.64	68.23±7.15*	68.18±11.58	66.45±8.19		
SBP (mm Hg)	130.46±11.82	129.85±10.64	123.45±9.2	123.09±10.24		
DBP (mm Hg)	77.23±9.87	76.38±7.85	75.82±8.51	72.27±7.14		
CSBP (mm Hg)	110.92±9.86	111.08±8.43	106.91±8.38	105.72±9.32		
CDBP (mm Hg)	76.08±17.35	77.69±7.77	76.36±8.64	73.64±7.38		
Aix (%)	-4.15±9.73	-4.15±8.73	-2.36±12.8	-1.18±11.71		
Cholesterol	4.82±0.88	4.77±1.04	4.01±1.0	3.97±0.77		
(mmol/l)^						
Triglyceride	1.01±0.48	1.07±0.46	0.9±0.38	0.82±0.18		
(mmol/l)^						
HDLC (mmol/l)^	1.43±0.25	1.43±0.3	1.37±0.2	1.29±0.14		
LDLC (mmol/l)^	2.92±0.79	2.87±0.88	2.22±0.75	2.32±0.74		
Glucose (mmol/l)^	4.78±0.38	5.22±0.52	4.9±0.46	4.92±0.31		
Insulin (mmol/l)^	6.94±2.72	10.72±6.44	7.15±3.65	12.78±16.42		
ŸO₂ _{max} (ml¹kg⁺min⁻¹)	47.43±5.64	48.47±6.0	45.85±7.34	44.92±5.49		
Maximal treadmill	14.4±1.12	14.9±1.11**	14.07±1.52	13.84±1.8		
speed (km·h ⁻¹)						
Sperm volume (ml)	4.7±1.4	5.17±0.87	3.95±1.7	3.97±1.79 ²⁴		

10 ⁹
10 ⁸ ±
10 ⁸
±14.14

Legend: Ht, height; Wt, weight; BMI, body mass index; HR, resting heart rate; SBP, systolic blood pressure; DBP, diastolic blood pressure; CSBP, central (aortic) blood pressure; CDBP, central (aortic) diastolic blood pressure; augmentation index; HDLC, high-density lipo-protein cholesterol; LDLC, low-density lipoprotein cholesterol; $\dot{V}O_{2max}$, maximal oxygen uptake; ^, n=8; #, number; %, percentage. Data are from two-tailed paired t-test and expressed as mean±SD. Sperm analysis was conducted on semen samples and not from highly motile sperm isolated using the PureSperm 40/80 reagents (Nidacon).

Table 2. Description of the three month exercise program.

Week	Session #	Training load*	Training sprint time (min)	Total session time (min)
1	1	3 sprints	1.5	9.5
	2	3 sprints	1.5	9.5
2	3	3 sprints	1.5	9.5
	4	3 sprints	1.5	9.5
3	5	4 sprints	2	14
	6	4 sprints	2	14
4	7	4 sprints	2	14
	8	4 sprints	2	14
5	9	5 sprints	2.5	18.5
	10	5 sprints	2.5	18.5
6	11	5 sprints	2.5	18.5
	12	5 sprints	2.5	18.5
7	13	6 sprints	3	23
	14	6 sprints	3	23
8	15	6 sprints	3	23
	16	6 sprints	3	23
9	17	6 sprints	3	23
	18	6 sprints	3	23
10	19	6 sprints	3	23
	20	6 sprints	3	23
11	21	6 sprints	3	23
	22	6 sprints	3	23
12	23	6 sprints	3	23
	24	6 sprints	3	23

Total Time 120 60 454

Legend: # number; * all sprints were completed at maximum intensity, separated by a four

minute passive recovery.

Location in relation to					DNA i	methylation (_		
CpG site	Closest	Gene	СрG	Chr	Before	After	Δ	<i>p</i> -value	<i>q</i> -value
	gene	region	island		(mean±SD)	exercise			(≤0.1)
						(mean±SD)			
cg02598618	-	-	Open	10	94.0±1.4	98.6±1.1	4.6	1.79×10 ⁻⁵	0.06
			sea						
cg02655397	-	-	N shelf	11	80.7±4.4	85.1±3.3	4.5	0.0005	0.08
cg10243075	CACNA1A	Body	N shore	19	23.2±3.6	27.6±5.0	4.4	0.0004	0.08
cg27222157	RBM14	TSS1500	N shore	11	22.2±6.8	26.5±8.5	4.4	0.0009	0.08
cg10413297	RFPL3S	TSS1500	Open	22	25.6±6.7	29.7±6.6	4.1	0.001	0.09
			sea						
cg16533336	WBSCR17	Body	Open	7	76.7±2.6	80.8±2.3	4.1	1.17×10 ⁻⁷	0.02
			sea						
cg09194930	MT1L	TSS200	Island	16	14.7±9.5	18.8±12.7	4.1	0.001	0.09

cg00685853	-	-	N shore	2	14.4±3.9	18.4±6.2	4.0	0.0001	0.07
cg27205941	GMDS	TSS200	Island	6	29.0±4.1	32.9±3.5	4.0	0.001	0.09
cg17745251	C10orf140	Body	Island	10	25.0±2.7	29.0±3.0	4.0	0.002	0.09
cg25485435	RAMP1	TSS1500	N shore	2	16.9±3.5	20.8±3.0	3.9	0.0002	0.07
cg11888420	-	-	Open	6	33.0±2.7	36.8±4.6	3.8	0.001	0.09
			sea						
cg08715827	-	-	Open	12	6.1±2.3	9.8±5.8	3.8	0.001	0.09
			sea						
cg15617548	GORASP2	5'UTR	Island	2	23.5±2.8	27.1±2.8	3.7	0.0009	0.08
		1stExon							
cg15488194	MBNL1	TSS1500	Open	3	23.9±11.9	27.5±12.8	3.7	0.002	0.097
		5'UTR	sea						
cg17271592	HIST2H2AA4	TSS1500	Island	1	13.8±3.1	17.4±2.6	3.6	0.0006	0.08
	HIST2H2AA3								
cg06785129	RFPL3	TSS1500	Open	22	18.6±3.2	22.2±4.2	3.6	0.002	0.09
			sea						

cg14047339	RUNX2	Body	Island	6	33.7±2.2	37.3±2.9	3.6	0.002	0.09
		5'UTR							
	SUPT3H	1stExon							
		5'UTR							
cg13916261	FNBP1	Body	N shelf	9	24.4±3.9	27.9±4.9	3.5	0.0003	0.08
cg09303484	-	-	Open	7	63.3±4.3	66.8±3.6	3.5	0.0008	0.08
			sea						
cg05528975	-	-	Open	2	25.9±2.5	29.4±3.1	3.5	0.0005	0.08
			sea						
cg17627829	HCG18	Body	N shore	6	9.5±3.4	13.0±4.1	3.5	0.002	0.09
cg05629323	TFDP3	3'UTR	Open	Χ	8.8±2.7	12.3±4.4	3.5	0.002	0.09
		1stExon	sea						
cg20681184	PIK3CD	5'UTR	S shore	1	14.5±2.8	18.0±4.5	3.5	0.002	0.09
cg15069995	HORMAD1	5'UTR	Open	1	5.9±2.1	9.3±3.9	3.4	0.0009	0.08
			sea						
cg13062455	SEL1L	Body	Open	14	88.2±2.3	85.4±3.3	-2.9	1.27×10 ⁻⁵	0.05

			sea						
cg24360197	-	-	Open	15	70.1±8.4	67.2±7.9	-2.9	0.001	0.09
			sea						
cg03359503	CLIC2	5'UTR	Open	X	87.7±4.1	84.7±.5	-3.0	0.0009	0.08
		1 st Exon	sea						
cg24121979	-	-	Open	12	94.9±3.2	91.9±2.0	-3.0	0.0003	0.08
			sea						
cg20909579	PDE6B	TSS1500	Island	4	89.6±2.9	86.6±5.4	-3.0	0.0008	0.08
		Body							
cg02507663	NCRNA00219	Body	S shore	5	54.9±8.4	51.9±8.4	-3.0	0.001	0.09
	SNORA13	TSS200							
cg27569203	CELSR1	Body	Island	22	90.2±2.4	87.3±1.6	-3.0	0.002	0.09
cg26235273	DKK3	3'UTR	Open	11	71.0±4.0	67.8±3.8	-3.1	0.0006	0.08
			sea						
cg17518550	C1orf150	TSS200	Open	1	82.3±4.0	79.2±6.4	-3.1	0.001	0.09
			sea						

cg16514287	SLC25A36	Body	Open	3	94.0±1.4	90.8±1.1	-3.1	1.7×10 ⁻⁶	0.03
			sea						
cg26057840†	-	-	N shore	20	89.4±2.7	86.2±2.6	-3.2	0.0005	0.08
cg24544876	-	-	S shore	5	78.3±3.3	75.1±2.5	-3.2	8000.0	0.08
cg14630692	URM1	Body	Open	9	86.4±1.7	83.2±2.4	-3.2	0.001	0.09
			sea						
cg03440485	-	-	Island	X	22.9±2.3	19.7±2.1	-3.2	0.001	0.09
cg15832577‡	C1orf53	Body	S shore	1	78.6±2.5	75.3±3.2	-3.3	0.0002	0.07
cg17494438	DRGX	TSS1500	N shelf	10	82.6±1.8	79.2±2.8	-3.3	0.001	0.09
cg16320838	DNAJB7	TSS1500	Open	22	92.5±14.5	89.0±19.1	-3.5	0.0005	0.08
	XPNPEP3	Body	sea						
cg11342670	KIAA0319	Body	Open	6	78.9±9.3	75.5±10.3	-3.5	0.001	0.09
			sea						
cg03987985	L1TD1	3'UTR	Open	1	75.6±4.6	72.0±5.6	-3.6	0.001	0.09
			sea						
cg08377570	LPAR6	TSS1500	Open	13	87.5±2.1	83.8±2.5	-3.7	0.0005	0.07

	RB1	Body	sea						596
cg17486639	-	-	N shelf	19	89.8±1.6	86.0±3.1	-3.8	0.0006	0.08 597
cg03834767	CDK14	3'UTR	Open	7	90.6±2.0	86.4±6.0	-4.1	0.002	0.09 598
			sea						599
cg16631698	-	-	Island	10	81.7±4.2	77.2±2.3	-4.5	0.002	0.09 600
cg08269188	CBX5	5'UTR	Open	12	83.7±2.7	79.0±4.5	-4.7	5.17×10 ⁻⁵	0.06 601
			sea						602
cg05946920	GGA1	Body	Island	22	39.7±6.7	33.6±3.4	-6.2	0.002	0.096603
									604

Data are expressed are beta values ± standard deviation.

Legend: Chr, chromosome number; Δ, difference; 3', three prime; 5', five prime; UTR, untranslated region; TSS1500, 1500 bases upstream of transcription start site; *DMR, differentially methylated region; †CDMR, cancer differentially methylated region; ‡RDMR, reprogrammed differentially methylated region.

Table 4. Molecular pathways related to genes with differentially methylated CpG sites after exercise training ($p \le 0.05$).

Pathway Name	Enrichment	<i>p</i> -value	#	Pathway ID
	Score		genes	
Circadian entrainment	8.0	0.0003	24	kegg pathway 51
Axon guidance	7.2	0.001	29	kegg pathway 176
Endocytosis	7.0	0.001	39	kegg pathway 211
Glutamatergic synapse	6.1	0.002	25	kegg pathway 182
Protein digestion and	5.9	0.003	22	kegg pathway 46
absorption				
ECM-receptor interaction	5.4	0.004	19	kegg pathway 41
GABAergic synapse	4.7	0.01	22	kegg pathway 133
Bile secretion	4.4	0.01	18	kegg pathway 74
MAPK signaling pathway	4.4	0.01	41	kegg pathway 119
Focal adhesion	4.3	0.01	34	kegg pathway 5
PI3K-Akt signaling pathway	4.3	0.01	54	kegg pathway 45
Huntington's disease	4.3	0.01	33	kegg pathway 53
Transcriptional misregulation	4.3	0.01	31	kegg pathway 94
in cancer				
African trypanosomiasis	4.1	0.02	10	kegg pathway 69
Colorectal cancer	4.1	0.02	13	kegg pathway 148
Viral myocarditis	3.9	0.02	15	kegg pathway 105
Taurine and hypotaurine	3.7	0.02	4	kegg pathway 195
metabolism				
Galactose metabolism	3.7	0.03	8	kegg pathway 15

Long-term depression	3.6	0.03	12	kegg pathway 250
Endocrine and other factor-	3.6	0.03	10	kegg pathway 258
regulated calcium				
reabsorption				
Retrograde endocannabinoid	3.6	0.03	21	kegg pathway 129
signaling				
Glycosaminoglycan	3.4	0.03	7	kegg pathway 248
biosynthesis - chondroitin				
sulfate / dermatan sulfate				
Nicotine addiction	3.4	0.03	10	kegg pathway 62
Allograft rejection	3.4	0.03	10	kegg pathway 177
Selenocompound	3.4	0.04	5	kegg pathway 135
metabolism				
Morphine addiction	3.2	0.04	20	kegg pathway 86
Amoebiasis	3.2	0.04	21	kegg pathway 56
Graft-versus-host disease	3.1	0.04	10	kegg pathway 209
Maturity onset diabetes of	3.1	0.04	7	kegg pathway 241
the young				
Pathways in cancer	3.1	0.05	49	kegg pathway 128
ErbB signaling pathway	3.1	0.05	15	kegg pathway 121
Type I diabetes mellitus	2.9	0.05	10	kegg pathway 242
Insulin secretion	2.9	0.05	17	kegg pathway 4

Table 5. Paternally imprinted gene CpG methylation changes caused by exercise training.

Paternally imprinted gene		ion to	n to		DNA methylation (%)		-		
	CpG site	Gene	СрG	Chr	Basal	After	Δ	<i>p</i> -value	<i>q</i> -value
		region	island		(mean±SD)	exercise			(≤0.1)
						(mean±SD)			
GRB10	cg12358041	5'UTR	Open sea	7	92.3±1.2	91.0±1.2	-1.3	0.0004	0.08
		Body							
	cg26050906	5'UTR	Open sea	7	91.8±1.1	90.9±1.1	-0.9	0.003	0.10
		TSS200							
SGCE	cg22090863*	Body	N shore	7	4.0±0.6	4.6±0.6	0.7	0.0007	0.08
PEG10		TSS1500							
SGCE	cg18139769	TSS1500	S shore	7	10.5±2.0	13.5±4.5	3.0	0.0006	0.08
PEG10		5'UTR							
DLGAP2	cg27316067	Body	S shore	8	64.3±11.2	62.2±11.3	-2.1	0.002	0.09
SFMBT2	cg15362304	Body	S shelf	10	93.4±1.1	92.5±1.1	-0.9	0.0007	0.08
INS-IGF2	cg00273464	Body	Open sea	11	89.8±1.7	87.7±1.2	-2.0	0.00002	0.06

IGF2		1stExon							
		5'UTR							
INS-IGF2	cg17665927	Body	Open sea	11	93.2±1.3	91.6±1.4	-1.5	0.0002	0.07
IGF2		1stExon							
		5'UTR							
INS-IGF2	cg05384664	Body	S shore	11	88.6±2.2	87.3±2.8	-1.4	0.001	0.09
IGF2		Body							
KCNQ10T1	cg20406758‡	Body	Open sea	11	95.5±1.0	94.1±1.0	-1.4	0.001	0.09
KCNQ1		Body							
SLC38A4	cg12152384	Body	Open sea	12	93.4±1.0	92.3±0.9	-1.1	0.002	0.09
WIF1	cg15862358	TSS1500	S shore	12	6.7±2.6	8.6±5.5	1.9	0.001	0.09
DLK1	cg17412258	TSS1500	CpG island	14	2.0±0.5	2.8±0.6	8.0	0.0005	0.08
	cg06504820	TSS1500	N shore	14	21.5±6.7	24.1±7.6	2.5	0.0003	0.07
RTL1	cg23546343	1stExon	N shore	14	93.0±0.9	91.8±1.0	-1.2	0.002	0.10
MIR431		TSS200							
MIR433		TSS1500							
GABRG3	cg08325929	Body	S shelf	15	86.2±2.5	84.9±2.7	-1.3	0.003	0.10

	ZIM2	cg16008476	Body	N shore	19	5.9±1.6	7.6±2.7	1.7	0.0006	0.08
	HM13	cg06000530	Body	CpG island	20	4.0±1.0	5.6±2.2	1.6	0.002	0.10
	PSIMCT-1		TSS200							
2	Data are expressed are beta values ± standard deviation.									
}	Legend: Chr, chromosome number; Δ, difference; 3', three prime; 5', five prime; UTR, untranslated region; TSS1500, 1500 bases upstream of									
Ļ	transcription start site; TSS20	00, 200 bases up	ostream of transcription	n start site; *DI	MR, di	fferentially me	thylated region	n; ‡RDI	MR, reprogra	ammed
5	differentially methylated regio	n.								
,										
?										
,										
,										
)										
-										
)										
}										
Ļ										

Supplementary Table 1. Gene ontology for genes with DNA methylation changes after exercise training ($q \le 0.1$).

Function	Туре	Enrichment	<i>p</i> -value	<i>q</i> -value	# genes	Go ID
		score				
Single-organism	BP	34.82	7.57×10 ⁻¹⁶	3.88×10 ⁻¹²	1428	44699
process						
Single-organism cellular	BP	26.32	3.70×10 ⁻¹²	9.50×10 ⁻⁹	1251	44763
process						
Biological process	BP	23.74	4.89×10 ⁻¹¹	8.22×10 ⁻⁸	1968	8150
Cellular process	BP	23.47	6.41×10 ⁻¹¹	8.22×10 ⁻⁸	1678	9987
Developmental process	BP	21.96	2.89×10 ⁻¹⁰	2.97×10 ⁻⁷	591	32502
Biological regulation	BP	19.53	3.31×10 ⁻⁹	2.83×10 ⁻⁶	1227	65007
Regulation of biological	BP	18.66	7.88×10 ⁻⁹	5.78×10 ⁻⁶	1168	50789
process						
Anatomical structure	BP	18.31	1.12×10 ⁻⁸	7.18×10 ⁻⁶	374	48856
development						
Anatomical structure	BP	17.79	1.87×10 ⁻⁸	1.07×10 ⁻⁵	213	9653
morphogenesis						

Cellular developmental	BP	17.46	2.62×10 ⁻⁸	1.29×10 ⁻⁵	312	48869
process						
Regulation of cellular	BP	17.38	2.84×10 ⁻⁸	1.29×10 ⁻⁵	1113	50794
process						
Single-organism	BP	17.32	3.01×10 ⁻⁸	1.29×10 ⁻⁵	405	44767
developmental process						
Ras guanyl-nucleotide	MF	17.22	3.32×10 ⁻⁸	1.31×10 ⁻⁵	36	5088
exchange factor activity						
Guanyl-nucleotide	MF	17.04	3.97×10 ⁻⁸	1.45×10 ⁻⁵	50	5085
exchange factor activity						
Gtpase regulator activity	MF	15.96	1.18×10 ⁻⁷	4.02×10 ⁻⁵	96	30695
Sequence-specific DNA	MF	15.66	1.58×10 ⁻⁷	4.68×10 ⁻⁵	182	3700
binding transcription						
factor activity						
Pattern specification	BP	15.65	1.60×10 ⁻⁷	4.68×10 ⁻⁵	84	7389
process						
Nucleic acid binding	MF	15.52	1.81×10 ⁻⁷	4.68×10 ⁻⁵	182	1071

transcription factor						
activity						
Small gtpase regulator	MF	15.52	1.82×10 ⁻⁷	4.68×10 ⁻⁵	71	5083
activity						
Ion binding	MF	15.49	1.88×10 ⁻⁷	4.68×10 ⁻⁵	867	43167
Nucleoside-	MF	15.47	1.91×10 ⁻⁷	4.68×10 ⁻⁵	97	60589
triphosphatase regulator						
activity						
Multicellular organismal	BP	14.95	3.22×10 ⁻⁷	7.51×10 ⁻⁵	435	32501
process						
Embryonic	BP	14.90	3.37×10 ⁻⁷	7.51×10 ⁻⁵	81	48598
morphogenesis						
Single-multicellular	BP	14.75	3.91×10 ⁻⁷	8.37×10 ⁻⁵	428	44707
organism process						
Molecular function	MF	14.47	5.18×10 ⁻⁷	0.0001	2015	3674
Regulation of cell	BP	14.16	7.09×10 ⁻⁷	0.0001	319	10646
communication						

Regulation of signaling	BP	14.14	7.22×10 ⁻⁷	0.0001	318	23051
Regulation of cell	BP	13.86	9.59×10 ⁻⁷	0.0002	124	60284
development						
Cell junction	CC	13.47	1.42×10 ⁻⁶	0.0003	136	30054
Sequence-specific DNA	MF	13.17	1.90×10 ⁻⁶	0.0003	127	43565
binding						
Binding	MF	12.88	2.55×10 ⁻⁶	0.0004	1597	5488
Cell differentiation	BP	12.46	3.88×10 ⁻⁶	0.0006	218	30154
Regulation of	BP	12.44	3.96×10 ⁻⁶	0.0006	80	50767
neurogenesis						
Regionalization	BP	12.19	5.08×10 ⁻⁶	0.0008	53	3002
Regulation of nervous	BP	12.16	5.24×10 ⁻⁶	0.0008	87	51960
system development						
Negative regulation of	BP	11.75	7.85×10 ⁻⁶	0.001	88	45596
cell differentiation						
Cellular response to	BP	11.54	9.69×10 ⁻⁶	0.001	653	51716
stimulus						

Extracellular matrix	CC	11.31	1.22×10 ⁻⁵	0.002	65	31012
Regulation of neuron	BP	11.27	1.28×10 ⁻⁵	0.002	67	45664
differentiation						
Motor activity	MF	11.21	1.35×10 ⁻⁵	0.002	34	3774
Extracellular matrix part	CC	11.21	1.35×10 ⁻⁵	0.002	44	44420
Organ development	BP	11.20	1.37×10 ⁻⁵	0.002	159	48513
Positive regulation of	BP	11.08	1.55×10 ⁻⁵	0.002	471	48522
cellular process						
Regulation of cell	BP	10.78	2.08×10 ⁻⁵	0.002	174	45595
differentiation						
Negative regulation of	BP	10.76	2.12×10 ⁻⁵	0.002	103	51093
developmental process						
Positive regulation of	BP	10.76	2.13×10 ⁻⁵	0.002	210	31328
cellular biosynthetic						
process						
Dendritic spine	CC	10.73	2.19×10 ⁻⁵	0.002	18	43197
Synaptic transmission	BP	10.72	2.22×10 ⁻⁵	0.002	81	7268

Neuron part	CC	10.65	2.36×10 ⁻⁵	0.002	105	97458
System development	BP	10.60	2.50×10 ⁻⁵	0.003	115	48731
Response to external	BP	10.57	2.57×10 ⁻⁵	0.003	168	9605
stimulus						
Synapse part	CC	10.35	3.21×10 ⁻⁵	0.003	71	44456
Cell communication	BP	10.34	3.22×10 ⁻⁵	0.003	148	7154
Signal transduction	BP	10.33	3.26×10 ⁻⁵	0.003	538	7165
Axon guidance	BP	10.31	3.32×10 ⁻⁵	0.003	68	7411
Negative regulation of	BP	10.15	3.91×10⁻⁵	0.003	18	45665
neuron differentiation						
Neuron spine	CC	10.15	3.91×10⁻⁵	0.003	18	44309
Cation binding	MF	10.14	3.94×10 ⁻⁵	0.003	581	43169
Metal ion binding	MF	10.07	4.23×10 ⁻⁵	0.004	572	46872
Positive regulation of	BP	10.06	4.26×10 ⁻⁵	0.004	211	9891
biosynthetic process						
Positive regulation of	BP	10.05	4.33×10 ⁻⁵	0.004	196	10557
macromolecule						

biosynthetic process						
Regulation of signal	BP	10.04	4.38×10 ⁻⁵	0.004	271	9966
transduction						
Phospholipid binding	MF	9.97	4.66×10 ⁻⁵	0.004	93	5543
Ion channel activity	MF	9.89	5.05×10 ⁻⁵	0.004	73	5216
Anterior/posterior	BP	9.83	5.40×10 ⁻⁵	0.004	36	9952
pattern specification						
Substrate-specific	MF	9.68	6.24×10 ⁻⁵	0.005	74	22838
channel activity						
Rho guanyl-nucleotide	MF	9.61	6.69×10 ⁻⁵	0.005	22	5089
exchange factor activity						
Negative regulation of	BP	9.56	7.04×10 ⁻⁵	0.005	110	9968
signal transduction						
Cell part	CC	9.55	7.10×10 ⁻⁵	0.005	1907	44464
Negative regulation of	BP	9.48	7.66×10 ⁻⁵	0.006	131	48585
response to stimulus						
Positive regulation of	BP	9.45	7.84×10 ⁻⁵	0.006	194	45935

nucleobase-containing compound metabolic process 8.13×10⁻⁵ ΒP 9.42 0.006 7 Regulation of axon 48841 extension involved in axon guidance 9.50×10⁻⁵ Negative regulation of BP 9.26 0.007 456 48519 biological process BP 9.26 9.55×10⁻⁵ 0.007 56 51056 Regulation of small gtpase mediated signal transduction 9.69×10⁻⁵ 72 ΒP 9.24 0.007 1900542 Regulation of purine nucleotide metabolic process Cell development ΒP 9.12 0.0001 0.007 79 48468 Negative regulation of 9.04 0.0001 800.0 114 10648 BP cell communication

Gated channel activity	MF	9.03	0.0001	0.008	60	22836
lon gated channel	MF	9.03	0.0001	0.008	60	22839
activity						
DNA binding	MF	9.00	0.0001	0.008	351	3677
Response to stimulus	BP	8.99	0.0001	0.008	827	50896
Positive regulation of	BP	8.94	0.0001	0.008	196	51173
nitrogen compound						
metabolic process						
Regulation of nucleotide	BP	8.91	0.0001	0.008	72	6140
metabolic process						
Channel activity	MF	8.88	0.0001	0.008	76	15267
Passive transmembrane	MF	8.88	0.0001	0.008	76	22803
transporter activity						
Organ morphogenesis	BP	8.80	0.0002	0.009	76	9887
Extracellular matrix	BP	8.78	0.0002	0.009	22	22617
disassembly						
Negative regulation of	BP	8.74	0.0002	0.009	113	23057

signaling						
Negative regulation of	BP	8.67	0.0002	0.01	421	48523
cellular process						
Neuron projection	CC	8.64	0.0002	0.01	80	43005
Cell projection part	CC	8.61	0.0002	0.01	111	44463
Positive regulation of	BP	8.61	0.0002	0.01	292	31325
cellular metabolic						
process						
Positive regulation of	BP	8.52	0.0002	0.01	173	51254
RNA metabolic process						
Regulation of cellular	BP	8.51	0.0002	0.01	202	51128
component organization						
Intracellular part	CC	8.51	0.0002	0.01	1607	44424
Enzyme regulator	MF	8.51	0.0002	0.01	161	30234
activity						
Positive regulation of	BP	8.48	0.0002	0.01	306	9893
metabolic process						

Regulation of Ras	BP	8.45	0.0002	0.01	35	46578
protein signal						
transduction						
Regulation of	BP	8.45	0.0002	0.01	275	51239
multicellular organismal						
process						
Cation channel activity	MF	8.40	0.0002	0.01	54	5261
Negative regulation of	BP	8.39	0.0002	0.01	22	90090
canonical Wnt receptor						
signaling pathway						
Collagen metabolic	BP	8.39	0.0002	0.01	22	32963
process						
Plasma membrane part	CC	8.38	0.0002	0.01	284	44459
Collagen	CC	8.37	0.0002	0.01	24	5581
Locomotion	BP	8.34	0.0002	0.01	159	40011
Regulation of	BP	8.27	0.0003	0.01	178	2000026
multicellular organismal						

development						
Regulation of Wnt	BP	8.25	0.0003	0.01	40	30111
receptor signaling						
pathway						
Collagen catabolic	BP	8.25	0.0003	0.01	20	30574
process						
Multicellular organismal	BP	8.20	0.0003	0.01	24	44236
metabolic process						
Regulation of	BP	8.14	0.0003	0.01	214	32879
localization						
Connective tissue	BP	8.08	0.0003	0.01	26	61448
development						
Positive regulation of	BP	8.07	0.0003	0.01	164	45893
transcription, DNA-						
dependent						
Positive regulation of	BP	8.06	0.0003	0.01	34	43547
gtpase activity						

Positive regulation of	BP	7.97	0.0003	0.02	284	10604
macromolecule						
metabolic process						
Neuron fate commitment	BP	7.96	0.0004	0.02	11	48663
Ionotropic glutamate	CC	7.96	0.0004	0.02	11	8328
receptor complex						
Synaptic membrane	CC	7.95	0.0004	0.02	45	97060
Multicellular organismal	BP	7.93	0.0004	0.02	21	44243
catabolic process						
Ion transmembrane	MF	7.90	0.0004	0.02	127	15075
transporter activity						
Cellular component	BP	7.86	0.0004	0.02	491	16043
organization						
Establishment of	BP	7.84	0.0004	0.02	427	51234
localization						
Embryonic placenta	BP	7.84	0.0004	0.02	10	1892
development						

Localization	BP	7.84	0.0004	0.02	71	51179
Tissue development	BP	7.83	0.0004	0.02	96	9888
Regulation of canonical	BP	7.68	0.0005	0.02	30	60828
Wnt receptor signaling						
pathway						
Energy reserve	BP	7.67	0.0005	0.02	32	6112
metabolic process						
Regulation of cellular	BP	7.67	0.0005	0.02	68	44087
component biogenesis						
Membrane-bounded	CC	7.66	0.0005	0.02	104	31988
vesicle						
Intracellular signal	BP	7.65	0.0005	0.02	206	35556
transduction						
Regulation of Rho	BP	7.59	0.0005	0.02	23	35023
protein signal						
transduction						
Axis specification	BP	7.52	0.0005	0.02	18	9798

Ion channel complex	CC	7.50	0.0006	0.02	41	34702
Microtubule motor	MF	7.48	0.0006	0.02	20	3777
activity						
Signaling	BP	7.46	0.0006	0.02	118	23052
Single organism	BP	7.46	0.0006	0.02	118	44700
signaling						
Negative regulation of	BP	7.44	0.0006	0.02	27	30178
Wnt receptor signaling						
pathway						
Embryonic organ	BP	7.44	0.0006	0.02	27	48562
morphogenesis						
Cell-cell signaling	BP	7.44	0.0006	0.02	115	7267
Positive regulation of	BP	7.39	0.0006	0.02	174	10628
gene expression						
Multicellular organismal	BP	7.32	0.0007	0.02	22	44259
macromolecule						
metabolic process						

Establishment or	BP	7.32	0.0007	0.02	22	7163
maintenance of cell						
polarity						
Regulation of	BP	7.32	0.0007	0.02	26	48638
developmental growth						
Lipid binding	MF	7.30	0.0007	0.02	123	8289
Regulation of cell	BP	7.29	0.0007	0.02	183	42127
proliferation						
Substrate-specific	MF	7.22	0.0007	0.03	134	22891
transmembrane						
transporter activity						
Regulation of	BP	7.21	0.0007	0.03	223	50793
developmental process						
Taxis	BP	7.16	0.0008	0.03	92	42330
Chemotaxis	BP	7.16	0.0008	0.03	92	6935
Ligand-gated ion	MF	7.15	0.0008	0.03	30	15276
channel activity						

Ligand-gated channel	MF	7.15	0.0008	0.03	30	22834
activity						
Cell fate determination	BP	7.15	0.0008	0.03	13	1709
Vesicle	CC	7.13	0.0008	0.03	116	31982
Negative regulation of	BP	7.02	0.0009	0.03	27	10721
cell development						
Wnt receptor signaling	BP	7.02	0.0009	0.03	44	16055
pathway						
Cellular component	BP	7.02	0.0009	0.03	491	71840
organization or						
biogenesis						
Positive regulation of	BP	7.00	0.0009	0.03	513	48518
biological process						
Membrane	CC	6.94	0.0010	0.03	801	16020
Regulation of metabolic	BP	6.92	0.0010	0.03	672	19222
process						
Positive regulation of	BP	6.90	0.0010	0.03	118	45944

transcription from RNA						
polymerase II promoter						
Negative regulation of	BP	6.90	0.0010	0.03	13	50771
axonogenesis						
Regulation of respiratory	BP	6.89	0.0010	0.03	8	43576
gaseous exchange						
Transport	BP	6.87	0.0010	0.03	416	6810
Transport vesicle	CC	6.86	0.0011	0.03	17	30658
membrane						
Energy derivation by	BP	6.85	0.0011	0.03	37	15980
oxidation of organic						
compounds						
Positive regulation of	BP	6.84	0.0011	0.03	33	10720
cell development						
Regulation of cell	BP	6.82	0.0011	0.03	51	30155
adhesion						
Actin filament-based	BP	6.82	0.0011	0.03	54	30029

process						
Cell surface receptor	BP	6.81	0.0011	0.03	312	7166
signaling pathway						
Cell-cell junction	CC	6.79	0.0011	0.03	52	5911
Activation of Ras gtpase	BP	6.79	0.0011	0.03	10	32856
activity						
Chromatin binding	MF	6.73	0.0012	0.04	62	3682
Tissue morphogenesis	BP	6.72	0.0012	0.04	63	48729
Regulation of	BP	6.70	0.0012	0.04	487	19219
nucleobase-containing						
compound metabolic						
process						
Negative regulation of	BP	6.69	0.0012	0.04	22	50768
neurogenesis						
Microtubule associated	CC	6.64	0.0013	0.04	29	5875
complex						
Transcription, DNA-	BP	6.59	0.0014	0.04	308	6351

dependent						
Early endosome	CC	6.57	0.0014	0.04	21	31901
membrane						
Cation channel complex	CC	6.52	0.0015	0.04	29	34703
Positive regulation of	BP	6.49	0.0015	0.04	26	50769
neurogenesis						
Coated vesicle	CC	6.49	0.0015	0.04	25	30662
membrane						
Columnar/cuboidal	BP	6.42	0.0016	0.05	7	2066
epithelial cell						
development						
Regulation of steroid	BP	6.42	0.0016	0.05	13	50810
biosynthetic process						
ER to Golgi transport	CC	6.42	0.0016	0.05	11	12507
vesicle membrane						
Dynein complex	CC	6.40	0.0017	0.05	12	30286
Positive regulation of	BP	6.40	0.0017	0.05	88	51130

cellular component						
organization						
Site of polarized growth	CC	6.39	0.0017	0.05	22	30427
Skeletal system	BP	6.37	0.0017	0.05	23	48705
morphogenesis						
Negative regulation of	BP	6.37	0.0017	0.05	23	10741
intracellular protein						
kinase cascade						
Common myeloid	BP	6.33	0.0018	0.05	3	35726
progenitor cell						
proliferation						
Regulation of	BP	6.33	0.0018	0.05	3	14807
somitogenesis						
Detection of	BP	6.33	0.0018	0.05	3	34287
monosaccharide						
stimulus						
Detection of glucose	BP	6.33	0.0018	0.05	3	51594

Detection of	BP	6.33	0.0018	0.05	3	9730
carbohydrate stimulus						
Detection of hexose	BP	6.33	0.0018	0.05	3	9732
stimulus						
Insulin-like growth	MF	6.33	0.0018	0.05	3	5010
factor-activated receptor						
activity						
Laminin-10 complex	CC	6.33	0.0018	0.05	3	43259
Cell projection	CC	6.26	0.0019	0.05	139	42995
Multicellular organismal	BP	6.25	0.0019	0.05	81	7275
development						
Positive regulation of	BP	6.23	0.0020	0.05	23	32320
Ras gtpase activity						
Endoplasmic reticulum	CC	6.22	0.0020	0.05	142	44432
part						
Regulation of GTP	BP	6.21	0.0020	0.05	44	33124
catabolic process						

Regulation of nitrogen	BP	6.20	0.0020	0.05	495	51171
compound metabolic						
process						
Neuron recognition	BP	6.19	0.0020	0.05	10	8038
Negative regulation of	BP	6.17	0.0021	0.05	88	8285
cell proliferation						
Regulation of cellular	BP	6.16	0.0021	0.05	612	31323
metabolic process						
Negative regulation of	BP	6.14	0.0022	0.05	5	48843
axon extension involved						
in axon guidance						
Negative regulation of	BP	6.14	0.0022	0.05	5	46426
JAK-STAT cascade						
N-methyl-D-aspartate	CC	6.14	0.0022	0.05	5	17146
selective glutamate						
receptor complex						
Regulation of	BP	6.10	0.0022	0.05	22	50770

axonogenesis						
Neuron differentiation	BP	6.09	0.0023	0.06	38	30182
Cytoskeleton	BP	6.05	0.0024	0.06	93	7010
organization						
Regulation of growth	BP	6.05	0.0024	0.06	85	40008
Regulation of	BP	6.01	0.0025	0.06	48	9118
nucleoside metabolic						
process						
Establishment of protein	BP	5.98	0.0025	0.06	9	90150
localization to						
membrane						
Integral to endoplasmic	CC	5.97	0.0025	0.06	21	30176
reticulum membrane						
Growth cone	CC	5.97	0.0025	0.06	21	30426
Cartilage development	BP	5.97	0.0025	0.06	6	60351
involved in						
endochondral bone						

morphogenesis						
Enzyme linked receptor	BP	5.96	0.0026	0.06	112	7167
protein signaling						
pathway						
Intramembranous	BP	5.93	0.0027	0.06	4	1957
ossification						
Direct ossification	BP	5.93	0.0027	0.06	4	36072
Regulation of transport	BP	5.91	0.0027	0.06	157	51049
Embryonic skeletal	BP	5.90	0.0027	0.06	18	48704
system morphogenesis						
Regulation of catabolic	BP	5.90	0.0027	0.06	88	9894
process						
Regulation of gtpase	BP	5.90	0.0027	0.06	43	43087
activity						
Transmembrane	BP	5.90	0.0028	0.06	87	7169
receptor protein tyrosine						
kinase signaling						

pathway						
Extracellular vesicular	CC	5.85	0.0029	0.07	16	70062
exosome						
Antigen processing and	BP	5.83	0.0029	0.07	21	19886
presentation of						
exogenous peptide						
antigen via MHC class II						
Tight junction	CC	5.83	0.0029	0.07	23	5923
Occluding junction	CC	5.83	0.0029	0.07	23	70160
Serine-type	MF	5.83	0.0029	0.07	22	4867
endopeptidase inhibitor						
activity						
Cellular component	BP	5.83	0.0029	0.07	138	6928
movement						
Membrane-enclosed	CC	5.78	0.0031	0.07	101	31974
lumen						
Vesicle membrane	CC	5.77	0.0031	0.07	61	12506

Regulatory region	MF	5.77	0.0031	0.07	62	1067
nucleic acid binding						
Regulatory region DNA	MF	5.77	0.0031	0.07	62	975
binding						
Regulation of primary	BP	5.76	0.0031	0.07	601	80090
metabolic process						
Regulation of purine	BP	5.72	0.0033	0.07	47	33121
nucleotide catabolic						
process						
Cartilage development	BP	5.72	0.0033	0.07	19	51216
Response to glucagon	BP	5.71	0.0033	0.07	12	33762
stimulus						
Regulation of	BP	5.70	0.0034	0.07	98	22603
anatomical structure						
morphogenesis						
Cytoplasmic vesicle	CC	5.65	0.0035	0.07	59	30659
membrane						

Regulation of biological	BP	5.65	0.0035	0.07	314	65008
quality						
Regulation of nucleotide	BP	5.64	0.0035	0.07	47	30811
catabolic process						
Metal ion	MF	5.64	0.0036	0.08	65	46873
transmembrane						
transporter activity						
Ionotropic glutamate	MF	5.62	0.0036	0.08	7	4970
receptor activity						
Cell motility	BP	5.60	0.0037	0.08	102	48870
Regulation of stress-	BP	5.60	0.0037	0.08	31	32872
activated MAPK						
cascade						
Single-organism	BP	5.59	0.0037	0.08	323	44765
transport						
Regulation of cellular	BP	5.58	0.0038	0.08	111	60341
localization						

Antigen processing and	BP	5.56	0.0038	0.08	21	2495
presentation of peptide						
antigen via MHC class II						
Response to	BP	5.56	0.0039	0.08	156	9719
endogenous stimulus						
Actin cytoskeleton	BP	5.55	0.0039	0.08	44	30036
organization						
Anchoring collagen	CC	5.55	0.0039	0.08	5	30934
Axonal growth cone	CC	5.55	0.0039	0.08	5	44295
Mannose metabolic	BP	5.55	0.0039	0.08	5	6013
process						
Semaphorin-plexin	BP	5.52	0.0040	0.08	6	71526
signaling pathway						
Biological adhesion	BP	5.52	0.0040	0.08	121	22610
Cell adhesion	BP	5.52	0.0040	0.08	121	7155
Extracellular organelle	CC	5.51	0.0040	0.08	16	43230
Extracellular membrane-	CC	5.51	0.0040	0.08	16	65010

bounded organelle						
Regulation of stress-	BP	5.50	0.0041	0.08	31	70302
activated protein kinase						
signaling cascade						
Positive regulation of	BP	5.50	0.0041	0.08	12	32024
insulin secretion						
Macromolecule	BP	5.47	0.0042	0.08	57	33036
localization						
Intrinsic to endoplasmic	CC	5.46	0.0043	0.08	23	31227
reticulum membrane						
Homeostatic process	BP	5.45	0.0043	0.08	152	42592
Transmembrane	MF	5.44	0.0043	0.08	141	22857
transporter activity						
Endocytosis	BP	5.44	0.0044	0.08	46	6897
Negative regulation of	BP	5.43	0.0044	0.08	20	7162
cell adhesion						
Integral to lumenal side	CC	5.41	0.0045	0.09	9	71556

of endoplasmic						
reticulum membrane						
Embryonic skeletal	BP	5.41	0.0045	0.09	10	48706
system development						
Protein binding	MF	5.40	0.0045	0.09	965	5515
Small gtpase mediated	BP	5.36	0.0047	0.09	71	7264
signal transduction						
Regulation of steroid	BP	5.35	0.0047	0.09	16	19218
metabolic process						
Secretion by cell	BP	5.33	0.0048	0.09	63	32940
Liver development	BP	5.32	0.0049	0.09	17	1889
Antigen processing and	BP	5.30	0.0050	0.09	21	2504
presentation of peptide						
or polysaccharide						
antigen via MHC class II						
Epithelial tube	BP	5.29	0.0050	0.09	19	60562
morphogenesis						

Regulation of response	BP	5.28	0.0051	0.09	348	48583
to stimulus						
Gtpase activator activity	MF	5.27	0.0051	0.09	47	5096
Organelle	CC	5.27	0.0051	0.09	1126	43226
Extracellular-glutamate-	MF	5.27	0.0051	0.09	7	5234
gated ion channel						
activity						
Osteoblast development	BP	5.27	0.0051	0.09	7	2076
Adherens junction	CC	5.25	0.0053	0.10	37	5912
Cardiac septum	BP	5.22	0.0054	0.10	11	60411
morphogenesis						
Cation transmembrane	MF	5.22	0.0054	0.10	94	8324
transporter activity						
Secretion	BP	5.22	0.0054	0.10	75	46903
Regulation of insulin	BP	5.21	0.0054	0.10	29	50796
secretion						
Regulation of peptide	BP	5.21	0.0055	0.10	31	90276

hormone secretion						
Regulation of hormone	BP	5.19	0.0056	0.10	36	46883
secretion						
Dorsal spinal cord	BP	5.19	0.0056	0.10	4	21516
development						
Peripheral nervous	BP	5.19	0.0056	0.10	4	32287
system myelin						
maintenance						
Positive regulation of	BP	5.19	0.0056	0.10	4	32930
superoxide anion						
generation						
Lipid particle	BP	5.19	0.0056	0.10	4	34389
organization						
Morphogenesis of an	BP	5.17	0.0057	0.10	50	2009
epithelium						
Positive regulation of	BP	5.17	0.0057	0.10	17	46887
hormone secretion						

Establishment of protein	BP	5.17	0.0057	0.10	8	90002
localization to plasma						
membrane						
Rab guanyl-nucleotide	MF	5.17	0.0057	0.10	8	17112
exchange factor activity						
Negative regulation of	BP	5.17	0.0057	0.10	10	45668
osteoblast differentiation						
Response to chemical	BP	5.16	0.0057	0.10	366	42221
stimulus						
Ras gtpase activator	MF	5.16	0.0057	0.10	25	5099
activity						
Response to nitrogen	BP	5.15	0.0058	0.10	96	1901698
compound						
Axonogenesis	BP	5.13	0.0059	0.10	24	7409
Regulation of JNK	BP	5.13	0.0059	0.10	27	46328
cascade						
Substrate-specific	MF	5.12	0.0059	0.10	146	22892

transporter activity						
Positive regulation of	BP	5.12	0.0060	0.10	81	45597
cell differentiation						
Cytoplasmic membrane-	CC	5.11	0.0060	0.10	89	16023
bounded vesicle						
RNA biosynthetic	BP	5.11	0.0061	0.10	323	32774
process						
Regulation of secretion	BP	5.10	0.0061	0.10	74	51046
Regulation of cell-	BP	5.10	0.0061	0.10	23	10810
substrate adhesion						
Neuropeptide receptor	MF	5.10	0.0061	0.10	12	8188
activity						
Establishment of cell	BP	5.10	0.0061	0.10	12	30010
polarity						
Cellular component	BP	5.09	0.0061	0.10	63	32989
morphogenesis						
Cell fate commitment	BP	5.09	0.0062	0.10	26	45165

Positive regulation of	BP	5.09	0.0062	0.10	40	51047
secretion						
Golgi apparatus part	CC	5.07	0.0063	0.10	103	44431
Response to organic	BP	5.05	0.0064	0.10	92	10243
nitrogen						
Regulation of	BP	5.05	0.0064	0.10	5	10889
sequestering of						
triglyceride						
Platelet-derived growth	MF	5.05	0.0064	0.10	5	48407
factor binding						
MHC class II receptor	MF	5.05	0.0064	0.10	5	32395
activity						
Positive regulation of	BP	5.04	0.0065	0.10	16	45666
neuron differentiation						
Small conductance	MF	5.04	0.0065	0.10	3	16286
calcium-activated						
potassium channel						

activity						
Cardiac endothelial cell	BP	5.04	0.0065	0.10	3	3348
differentiation						
Endocardial cell	BP	5.04	0.0065	0.10	3	60956
differentiation						
N-acetylgalactosaminyl-	MF	5.04	0.0065	0.10	3	50510
proteoglycan 3-beta-						
glucuronosyltransferase						
activity						
Positive regulation of	BP	5.04	0.0065	0.10	3	7221
transcription of Notch						
receptor target						
Glossopharyngeal nerve	BP	5.04	0.0065	0.10	3	21615
morphogenesis						
Extracellular matrix-cell	BP	5.04	0.0065	0.10	3	35426
signaling						
Immunoglobulin	BP	5.04	0.0065	0.10	3	2381

production involved in immunoglobulin mediated immune response Nerve growth factor ΒP 5.04 0.0065 0.10 32455 3 processing Enzyme binding MF 5.04 0.0065 0.10 162 19899 Cell recognition ΒP 5.03 0.0066 0.10 17 8037 Regulation of peptide ΒP 5.02 0.0066 0.10 31 2791 secretion Regulation of peptide BP 5.02 0.0066 0.10 31 90087 transport Cytoplasmic vesicle part CC 4.99 0.0068 0.10 73 44433 Skeletal system ΒP 4.97 0.0069 0.11 30 1501 development Postsynaptic membrane CC 4.97 0.0070 0.11 35 45211 Postsynaptic density CC 4.96 0.0070 0.11 22 14069

Bone morphogenesis	BP	4.95	0.0071	0.11	7	60349
Synaptic transmission,	BP	4.95	0.0071	0.11	7	35249
glutamatergic						
Cortical actin	CC	4.95	0.0071	0.11	7	30864
cytoskeleton						
Cellular response to	BP	4.95	0.0071	0.11	10	71377
glucagon stimulus						
Negative regulation of	BP	4.94	0.0071	0.11	36	45926
growth						
Regulation of	BP	4.92	0.0073	0.11	185	6357
transcription from RNA						
polymerase II promoter						
Embryonic digit	BP	4.91	0.0074	0.11	12	42733
morphogenesis						
Regulation of extent of	BP	4.91	0.0074	0.11	12	61387
cell growth						
Potassium channel	CC	4.90	0.0075	0.11	16	34705

complex						
Voltage-gated	CC	4.90	0.0075	0.11	16	8076
potassium channel						
complex						
Regulation of JAK-STAT	BP	4.90	0.0075	0.11	16	46425
cascade						
Regulation of	BP	4.88	0.0076	0.11	186	51174
phosphorus metabolic						
process						
Gland development	BP	4.84	0.0079	0.12	22	48732
Anion binding	MF	4.84	0.0079	0.12	380	43168
Regulation of	BP	4.83	0.0080	0.12	34	42391
membrane potential						
Regulation of phosphate	BP	4.81	0.0081	0.12	184	19220
metabolic process						
Epithelial cell	BP	4.80	0.0082	0.12	24	2064
development						

Embryonic heart tube	BP	4.79	0.0083	0.12	14	3143
morphogenesis						
Inner ear	BP	4.79	0.0083	0.12	14	42472
morphogenesis						
Regulation of vesicle-	BP	4.79	0.0084	0.12	40	60627
mediated transport						
Clathrin-coated vesicle	CC	4.77	0.0085	0.12	19	30665
membrane						
Cell migration	BP	4.77	0.0085	0.12	91	16477
Central nervous system	BP	4.77	0.0085	0.12	15	21953
neuron differentiation						
Negative regulation of	BP	4.75	0.0086	0.12	6	30517
axon extension						
Protein kinase C activity	MF	4.75	0.0086	0.12	6	4697
Regulation of cellular	BP	4.75	0.0086	0.12	75	31329
catabolic process						
Response to growth	BP	4.75	0.0086	0.12	83	70848

factor stimulus						
Organic substance	BP	4.75	0.0087	0.12	500	1901576
biosynthetic process						
L-amino acid	MF	4.73	0.0088	0.12	10	15179
transmembrane						
transporter activity						
Embryo implantation	BP	4.73	0.0088	0.12	10	7566
Endocytic vesicle	CC	4.73	0.0088	0.12	22	30666
membrane						
Glycogen metabolic	BP	4.73	0.0088	0.12	12	5977
process						
Transcription-coupled	BP	4.73	0.0088	0.12	12	6283
nucleotide-excision						
repair						
Neuron projection	BP	4.72	0.0089	0.12	32	48812
morphogenesis						
Epidermis development	BP	4.69	0.0092	0.13	28	8544

Regulation of actin	BP	4.68	0.0093	0.13	37	32970
filament-based process						
Cellular response to	BP	4.68	0.0093	0.13	81	71363
growth factor stimulus						
Response to nicotine	BP	4.67	0.0094	0.13	9	35094
Midbrain development	BP	4.67	0.0094	0.13	9	30901
Cell body	CC	4.67	0.0094	0.13	44	44297
Positive regulation of	BP	4.67	0.0094	0.13	13	90277
peptide hormone						
secretion						
G-protein coupled	BP	4.67	0.0094	0.13	84	7186
receptor signaling						
pathway						
Cellular nitrogen	BP	4.66	0.0095	0.13	366	44271
compound biosynthetic						
process						
Pancreas development	BP	4.65	0.0095	0.13	7	31016

Ventricular septum	BP	4.65	0.0095	0.13	7	60412
morphogenesis						
Protein localization	BP	4.65	0.0096	0.13	51	8104
Regulation of actin	BP	4.64	0.0096	0.13	35	32956
cytoskeleton						
organization						
Regulation of cyclic	BP	4.64	0.0096	0.13	25	30799
nucleotide metabolic						
process						
Binding, bridging	MF	4.64	0.0097	0.13	27	60090
Neuronal cell body	CC	4.64	0.0097	0.13	40	43025
Canonical Wnt receptor	BP	4.62	0.0099	0.13	17	60070
signaling pathway						
Atrial septum	BP	4.61	0.0099	0.13	5	60413
morphogenesis						
Outer membrane-	CC	4.61	0.0099	0.13	5	30288
bounded periplasmic						

space						
Periplasmic space	CC	4.61	0.0099	0.13	5	42597
Detection of bacterium	BP	4.61	0.0099	0.13	5	16045
Endocytic recycling	BP	4.61	0.0099	0.13	5	32456
Microtubule	CC	4.61	0.01	0.13	58	5874
Positive regulation of	BP	4.61	0.01	0.13	79	51050
transport						
Hepatocyte	BP	4.59	0.01	0.13	4	70365
differentiation						
Glomerular visceral	BP	4.59	0.01	0.13	4	72015
epithelial cell						
development						
Hydrogen peroxide	BP	4.59	0.01	0.13	4	50665
biosynthetic process						
RNA-induced silencing	CC	4.59	0.01	0.13	4	16442
complex						
Rnai effector complex	CC	4.59	0.01	0.13	4	31332

Regulation of dendritic	BP	4.59	0.01	0.13	4	2604
cell antigen processing						
and presentation						
Left/right axis	BP	4.59	0.01	0.13	4	70986
specification						
CARD domain binding	MF	4.59	0.01	0.13	4	50700
Neuron migration	BP	4.59	0.01	0.13	21	1764
Cellular localization	BP	4.59	0.01	0.13	49	51641
Regulation of cell	BP	4.59	0.01	0.13	34	10769
morphogenesis involved						
in differentiation						
Regulation of cellular	BP	4.58	0.01	0.13	73	51270
component movement						
Glucan metabolic	BP	4.55	0.01	0.13	12	44042
process						
Cellular glucan	BP	4.55	0.01	0.13	12	6073
metabolic process						

Brain development	BP	4.55	0.01	0.13	32	7420
Transcription regulatory	MF	4.55	0.01	0.13	58	44212
region DNA binding						
Cell proliferation	BP	4.55	0.01	0.13	88	8283
Regulation of	BP	4.53	0.01	0.13	10	32651
interleukin-1 beta						
production						
Biosynthetic process	BP	4.52	0.01	0.13	507	9058
Cytoplasmic vesicle	CC	4.51	0.01	0.14	97	31410
Positive regulation of	BP	4.51	0.01	0.14	13	2793
peptide secretion						
Positive regulation of	BP	4.49	0.01	0.14	14	42108
cytokine biosynthetic						
process						
Appendage	BP	4.48	0.01	0.14	21	35107
morphogenesis						
Limb morphogenesis	BP	4.48	0.01	0.14	21	35108

Embryonic forelimb	BP	4.45	0.01	0.14	9	35115
morphogenesis						
Tight junction assembly	BP	4.45	0.01	0.14	9	70830
Positive regulation of	BP	4.45	0.01	0.14	25	45785
cell adhesion						
Cellular response to	BP	4.44	0.01	0.14	11	71229
acid						
Golgi membrane	CC	4.43	0.01	0.14	80	139
Asymmetric protein	BP	4.42	0.01	0.14	6	8105
localization						
Positive regulation of	BP	4.42	0.01	0.14	106	51094
developmental process						
Neuron development	BP	4.41	0.01	0.15	22	48666
Potassium channel	MF	4.41	0.01	0.15	22	5267
activity						
Regulation of camp	BP	4.41	0.01	0.15	22	30814
metabolic process						

Cytoplasmic part	CC	4.40	0.01	0.15	888	44444
Intrinsic to plasma	CC	4.40	0.01	0.15	156	31226
membrane						
Coated vesicle	CC	4.40	0.01	0.15	26	30135
Clathrin-coated vesicle	CC	4.40	0.01	0.15	24	30136
Negative regulation of	BP	4.40	0.01	0.15	8	48640
developmental growth						
Insulin-like growth factor	MF	4.40	0.01	0.15	8	5520
binding						
Regulation of	BP	4.39	0.01	0.15	73	40012
locomotion						
Intracellular organelle	CC	4.39	0.01	0.15	1114	43229
Cellular response to	BP	4.38	0.01	0.15	7	35924
vascular endothelial						
growth factor stimulus						
Positive regulation of	BP	4.38	0.01	0.15	7	10718
epithelial to						

mesenchymal transition								
DNA-dependent	BP	4.37	0.01	0.15	38	6352		
transcription, initiation								
Calcium channel activity	MF	4.37	0.01	0.15	21	5262		
Lung development	BP	4.36	0.01	0.15	17	30324		
Regulation of	BP	4.34	0.01	0.15	14	14013		
gliogenesis								
Regulation of axon	BP	4.34	0.01	0.15	10	30516		
extension								
Positive regulation of	BP	4.34	0.01	0.15	10	31281		
cyclase activity								
Aromatic compound	BP	4.34	0.01	0.15	355	19438		
biosynthetic process								
Organic cyclic	BP	4.32	0.01	0.15	370	1901362		
compound biosynthetic								
process								
Organelle lumen	CC	4.32	0.01	0.15	90	43233		

Cellular_component	CC	4.30	0.01	0.15	2166	5575
Aminoglycan metabolic	BP	4.30	0.01	0.15	29	6022
process						
Digestive tract	BP	4.27	0.01	0.16	11	48565
development						
Cortical cytoskeleton	CC	4.27	0.01	0.16	11	30863
Inorganic cation	MF	4.26	0.01	0.16	74	22890
transmembrane						
transporter activity						
Anterior/posterior axis	BP	4.25	0.01	0.16	9	9948
specification						
Cytoskeleton	CC	4.23	0.01	0.16	106	5856
Cellular membrane	BP	4.23	0.01	0.16	64	16044
organization						
Plasma membrane	CC	4.23	0.01	0.16	488	5886
Dopaminergic neuron	BP	4.23	0.01	0.16	5	71542
differentiation						

Regulation of respiratory	BP	4.23	0.01	0.16	5	2087
gaseous exchange by						
neurological system						
process						
Positive regulation of	BP	4.22	0.01	0.16	16	30177
Wnt receptor signaling						
pathway						
Phosphoglucomutase	MF	4.22	0.01	0.16	3	4614
activity						
Microvillus assembly	BP	4.22	0.01	0.16	3	30033
Microvillus organization	BP	4.22	0.01	0.16	3	32528
Noradrenergic neuron	BP	4.22	0.01	0.16	3	3357
differentiation						
Positive regulation of	BP	4.22	0.01	0.16	3	2000096
Wnt receptor signaling						
pathway, planar cell						
polarity pathway						

Response to oleic acid	BP	4.22	0.01	0.16	3	34201
Chemorepellent activity	MF	4.22	0.01	0.16	3	45499
Positive regulation of	BP	4.22	0.01	0.16	3	90031
steroid hormone						
biosynthetic process						
Membrane-bounded	CC	4.19	0.02	0.17	1000	43227
organelle						
Generation of precursor	BP	4.17	0.02	0.17	63	6091
metabolites and energy						
Non-canonical Wnt	BP	4.17	0.02	0.17	8	35567
receptor signaling						
pathway						
Microtubule-based	BP	4.16	0.02	0.17	28	7018
movement						
Heterocycle biosynthetic	BP	4.16	0.02	0.17	354	18130
process						
Nucleobase-containing	BP	4.15	0.02	0.17	345	34654

compound biosynthetic						
process						
Glycogen biosynthetic	BP	4.13	0.02	0.17	6	5978
process						
Glucan biosynthetic	BP	4.13	0.02	0.17	6	9250
process						
Cell differentiation in	BP	4.13	0.02	0.17	6	21533
hindbrain						
MHC class II protein	CC	4.13	0.02	0.17	6	42613
complex						
Regulation of timing of	BP	4.11	0.02	0.17	4	48505
cell differentiation						
Glomerular epithelial cell	BP	4.11	0.02	0.17	4	72310
development						
Negative regulation of	BP	4.11	0.02	0.17	4	35414
catenin import into						
nucleus						

Negative regulation of	BP	4.11	0.02	0.17	4	51151
smooth muscle cell						
differentiation						
Pinocytosis	BP	4.11	0.02	0.17	4	6907
Semaphorin receptor	MF	4.11	0.02	0.17	4	17154
activity						
RNA polymerase II core	MF	4.11	0.02	0.17	4	993
binding						
Regulation of glial cell	BP	4.10	0.02	0.17	11	45685
differentiation						
Regulation of stem cell	BP	4.10	0.02	0.18	16	72091
proliferation						
Extracellular matrix	BP	4.09	0.02	0.18	43	30198
organization						
Regulation of DNA	BP	4.08	0.02	0.18	15	51101
binding						
Organic cyclic	MF	4.07	0.02	0.18	721	97159

compound binding						
Regulation of	BP	4.07	0.02	0.18	12	32642
chemokine production						
Phagocytic vesicle	CC	4.07	0.02	0.18	12	30670
membrane						
Palate development	BP	4.06	0.02	0.18	14	60021
Regulation of cyclase	BP	4.06	0.02	0.18	14	31279
activity						
Cellular biosynthetic	BP	4.06	0.02	0.18	482	44249
process						
Positive regulation of	BP	4.05	0.02	0.18	9	14015
gliogenesis						
Tissue homeostasis	BP	4.04	0.02	0.18	18	1894
Regulation of cell	BP	4.03	0.02	0.18	44	31344
projection organization						
Extracellular structure	BP	4.02	0.02	0.18	43	43062
organization						

Actin binding	MF	4.01	0.02	0.19	58	3779
Endosome membrane	CC	4.00	0.02	0.19	51	10008
Regulation of cell	BP	3.98	0.02	0.19	48	1558
growth						
Positive regulation of	BP	3.98	0.02	0.19	10	51349
lyase activity						
Response to fatty acid	BP	3.98	0.02	0.19	10	70542
Regulation of epithelial	BP	3.98	0.02	0.19	10	10717
to mesenchymal						
transition						
Tube morphogenesis	BP	3.97	0.02	0.19	19	35239
Macromolecule	BP	3.97	0.02	0.19	384	9059
biosynthetic process						
Embryo development	BP	3.97	0.02	0.19	37	9792
ending in birth or egg						
hatching						
Growth factor binding	MF	3.96	0.02	0.19	21	19838

Antigen processing and	BP	3.96	0.02	0.19	30	2478
presentation of						
exogenous peptide						
antigen						
Positive regulation of	BP	3.96	0.02	0.19	8	50766
phagocytosis						
Vesicle-mediated	BP	3.94	0.02	0.19	112	16192
transport						
Coated pit	CC	3.94	0.02	0.19	11	5905
Membrane organization	BP	3.94	0.02	0.19	64	61024
Glycosaminoglycan	BP	3.94	0.02	0.19	27	30203
metabolic process						
Divalent inorganic cation	MF	3.94	0.02	0.19	27	72509
transmembrane						
transporter activity						
Protein localization to	BP	3.93	0.02	0.19	14	72657
membrane						

Heart looping	BP	3.92	0.02	0.20	12	1947
Cell morphogenesis	BP	3.92	0.02	0.20	22	904
involved in						
differentiation						
Anchoring junction	CC	3.90	0.02	0.20	37	70161
Regulation of	BP	3.89	0.02	0.20	5	60850
transcription involved in						
cell fate commitment						
Spinal cord association	BP	3.89	0.02	0.20	5	21527
neuron differentiation						
Intrinsic to internal side	CC	3.89	0.02	0.20	5	31235
of plasma membrane						
Dorsal/ventral axis	BP	3.89	0.02	0.20	5	9950
specification						
Regulation of respiratory	BP	3.89	0.02	0.20	5	44065
system process						
Lens morphogenesis in	BP	3.89	0.02	0.20	5	2089

camera-type eye						
DNA-directed RNA	CC	3.89	0.02	0.20	5	5665
polymerase II, core						
complex						
Actin monomer binding	MF	3.89	0.02	0.20	5	3785
Central nervous system	BP	3.89	0.02	0.20	7	21954
neuron development						
Mesenchymal cell	BP	3.89	0.02	0.20	7	14031
development						
Positive regulation of	BP	3.89	0.02	0.20	7	45687
glial cell differentiation						
Zinc ion binding	MF	3.88	0.02	0.20	280	8270
Positive regulation of	BP	3.86	0.02	0.20	9	45762
adenylate cyclase						
activity						
Endosomal part	CC	3.86	0.02	0.20	53	44440
Neurotransmitter:sodium	MF	3.85	0.02	0.20	6	5328

symporter activity						
Regulation of	BP	3.85	0.02	0.20	6	2000677
transcription regulatory						
region DNA binding						
Cytoskeletal protein	MF	3.85	0.02	0.20	97	8092
binding						
Regulation of cellular	BP	3.84	0.02	0.20	458	31326
biosynthetic process						
Regulation of molecular	BP	3.84	0.02	0.21	254	65009
function						
Anion transmembrane	MF	3.83	0.02	0.21	40	8509
transporter activity						
Polysaccharide	BP	3.83	0.02	0.21	15	5976
metabolic process						
Regulation of stem cell	BP	3.83	0.02	0.21	15	2000736
differentiation						
Anatomical structure	BP	3.82	0.02	0.21	29	60249

homeostasis						
Regulation of	BP	3.81	0.02	0.21	10	32652
interleukin-1 production						
Stem cell proliferation	BP	3.81	0.02	0.21	10	72089
Antigen processing and	BP	3.81	0.02	0.21	30	19884
presentation of						
exogenous antigen						
Myosin complex	CC	3.80	0.02	0.21	14	16459
Membrane part	CC	3.80	0.02	0.21	808	44425
Voltage-gated channel	MF	3.80	0.02	0.21	31	22832
activity						
Voltage-gated ion	MF	3.80	0.02	0.21	31	5244
channel activity						
Regulation of adenylate	BP	3.79	0.02	0.21	13	45761
cyclase activity						
Regulation of cartilage	BP	3.79	0.02	0.21	11	61035
development						

Positive regulation of	BP	3.78	0.02	0.21	17	60193
lipase activity						
Organelle membrane	CC	3.78	0.02	0.21	307	31090
Positive regulation of	BP	3.78	0.02	0.21	12	46330
JNK cascade						
Regulation of cell	BP	3.78	0.02	0.21	66	2000145
motility						
Synapse	CC	3.78	0.02	0.21	34	45202
Positive regulation of	BP	3.77	0.02	0.21	21	51495
cytoskeleton						
organization						
Regulation of neuron	BP	3.77	0.02	0.21	36	10975
projection development						
Cell part morphogenesis	BP	3.77	0.02	0.21	38	32990
Cell differentiation in	BP	3.76	0.02	0.21	8	21515
spinal cord						
Face morphogenesis	BP	3.76	0.02	0.21	8	60325

Stem cell development	BP	3.76	0.02	0.21	8	48864
Activation of adenylate	BP	3.76	0.02	0.21	8	7190
cyclase activity						
Embryo development	BP	3.74	0.02	0.21	52	9790
Heterocyclic compound	MF	3.72	0.02	0.22	709	1901363
binding						
Regulation of	BP	3.72	0.02	0.22	565	60255
macromolecule						
metabolic process						
Stress-activated MAPK	BP	3.72	0.02	0.22	18	51403
cascade						
Regulation of RNA	BP	3.71	0.02	0.22	406	51252
metabolic process						
Hexose metabolic	BP	3.71	0.02	0.22	33	19318
process						
Cell projection	BP	3.70	0.02	0.22	75	30030
organization						

Positive regulation of	BP	3.70	0.02	0.22	4	45080
chemokine biosynthetic						
process						
Hematopoietic stem cell	BP	3.70	0.02	0.22	4	71425
proliferation						
Regulation of	BP	3.70	0.02	0.22	4	40034
development,						
heterochronic						
Apical protein	BP	3.70	0.02	0.22	4	45176
localization						
Ventricular trabecula	BP	3.70	0.02	0.22	4	3222
myocardium						
morphogenesis						
Positive regulation of	BP	3.70	0.02	0.22	4	33235
protein sumoylation						
Pre-mrna binding	MF	3.70	0.02	0.22	4	36002
Peripheral nervous	BP	3.70	0.02	0.22	4	48935

system neuron						
development						
Metaphase	BP	3.70	0.02	0.22	4	51323
Regulation of	BP	3.70	0.02	0.22	4	32928
superoxide anion						
generation						
Activation of	BP	3.70	0.02	0.22	4	7171
transmembrane receptor						
protein tyrosine kinase						
activity						
DNA synthesis involved	BP	3.70	0.02	0.22	4	731
in DNA repair						
Positive regulation of	BP	3.70	0.02	0.22	4	60391
SMAD protein import						
into nucleus						
Basal transcription	MF	3.70	0.02	0.22	4	1098
machinery binding						

Basal RNA polymerase	MF	3.70	0.02	0.22	4	1099
II transcription						
machinery binding						
Protein domain specific	MF	3.69	0.03	0.22	79	19904
binding						
Cellular response to	BP	3.69	0.03	0.22	100	71495
endogenous stimulus						
Heterotrimeric G-protein	CC	3.69	0.03	0.22	9	5834
complex						
Regulation of lyase	BP	3.68	0.03	0.22	14	51339
activity						
Amino acid	MF	3.68	0.03	0.22	14	15171
transmembrane						
transporter activity						
Voltage-gated cation	MF	3.68	0.03	0.22	24	22843
channel activity						
Axon part	CC	3.68	0.03	0.22	24	33267

Positive regulation of	BP	3.68	0.03	0.22	17	45732
protein catabolic						
process						
Endoderm development	BP	3.67	0.03	0.22	7	7492
Chondroitin sulfate	BP	3.67	0.03	0.22	7	30206
biosynthetic process						
Dorsal/ventral pattern	BP	3.66	0.03	0.22	13	9953
formation						
Positive regulation of	BP	3.65	0.03	0.22	25	31346
cell projection						
organization						
Recycling endosome	CC	3.64	0.03	0.22	12	55037
Response to organic	BP	3.64	0.03	0.22	261	10033
substance						
Voltage-gated	MF	3.63	0.03	0.22	16	5249
potassium channel						
activity						

Regulation of	BP	3.63	0.03	0.22	460	9889
biosynthetic process						
Regulation of	BP	3.63	0.03	0.22	20	50920
chemotaxis						
Response to acid	BP	3.62	0.03	0.22	18	1101
Wnt receptor signaling	BP	3.62	0.03	0.22	3	7223
pathway, calcium						
modulating pathway						
Adenylate cyclase	MF	3.62	0.03	0.22	3	8179
binding						
Regulation of bile acid	BP	3.62	0.03	0.22	3	70857
biosynthetic process						
Neutrophil degranulation	BP	3.62	0.03	0.22	3	43312
Coumarin metabolic	BP	3.62	0.03	0.22	3	9804
process						
N-methyl-D-aspartate	MF	3.62	0.03	0.22	3	4972
selective glutamate						

receptor activity						
FACIT collagen	CC	3.62	0.03	0.22	3	5593
WINAC complex	CC	3.62	0.03	0.22	3	71778
COPI-coated vesicle	CC	3.62	0.03	0.22	3	30137
Eukaryotic initiation	MF	3.62	0.03	0.22	3	8190
factor 4E binding						
Opioid receptor	BP	3.62	0.03	0.22	3	38003
signaling pathway						
Nerve growth factor	MF	3.62	0.03	0.22	3	48406
binding						
Somatic muscle	BP	3.62	0.03	0.22	3	7525
development						
Positive regulation of	BP	3.62	0.03	0.22	3	2000052
non-canonical Wnt						
receptor signaling						
pathway						
Positive regulation of	BP	3.62	0.03	0.22	3	10890

sequestering of						
triglyceride						
Coronary vasculature	BP	3.62	0.03	0.22	3	60977
morphogenesis						
Forebrain	BP	3.62	0.03	0.22	3	21797
anterior/posterior						
pattern specification						
Glycine transmembrane	MF	3.62	0.03	0.22	3	15187
transporter activity						
Laminin-1 binding	MF	3.62	0.03	0.22	3	43237
Growth plate cartilage	BP	3.62	0.03	0.22	3	3417
development						
Gene silencing by mirna	BP	3.62	0.03	0.22	3	35195
Retina vasculature	BP	3.62	0.03	0.22	3	61299
morphogenesis in						
camera-type eye						
Immunoglobulin	BP	3.62	0.03	0.22	3	2377

production						
JUN kinase binding	MF	3.62	0.03	0.22	3	8432
Mitotic metaphase	BP	3.62	0.03	0.22	3	89
Sensory perception of	BP	3.62	0.03	0.22	3	50916
sweet taste						
Chloride channel	MF	3.62	0.03	0.22	3	17081
regulator activity						
Collagen biosynthetic	BP	3.62	0.03	0.22	3	32964
process						
Endothelial cell	BP	3.60	0.03	0.22	6	45446
differentiation						
Activation of Rho gtpase	BP	3.60	0.03	0.22	6	32862
activity						
Intracellular ligand-	MF	3.60	0.03	0.22	6	5217
gated ion channel						
activity						
Microfilament motor	MF	3.60	0.03	0.22	6	146

activity						
Positive regulation of	BP	3.59	0.03	0.22	15	45807
endocytosis						
Carbohydrate	BP	3.59	0.03	0.22	21	33500
homeostasis						
Glucose homeostasis	BP	3.59	0.03	0.22	21	42593
Response to arsenic-	BP	3.59	0.03	0.22	5	46685
containing substance						
Alpha-mannosidase	MF	3.59	0.03	0.22	5	4559
activity						
Nua4 histone	CC	3.59	0.03	0.22	5	35267
acetyltransferase						
complex						
Axonemal dynein	CC	3.59	0.03	0.22	5	5858
complex						
Tubulin-tyrosine ligase	MF	3.59	0.03	0.22	5	4835
activity						

Extracellular region part	CC	3.58	0.03	0.22	160	44421
Negative regulation of	BP	3.57	0.03	0.22	25	30308
cell growth						
Regulation of lipid	BP	3.57	0.03	0.22	19	46890
biosynthetic process						
Tissue remodeling	BP	3.56	0.03	0.23	14	48771
Regulation of cell	BP	3.56	0.03	0.23	49	22604
morphogenesis						
Intracellular membrane-	CC	3.54	0.03	0.23	990	43231
bounded organelle						
Potassium ion	MF	3.53	0.03	0.23	23	15079
transmembrane						
transporter activity						
Metalloendopeptidase	MF	3.53	0.03	0.23	20	4222
activity						
Negative regulation of	BP	3.53	0.03	0.23	16	31345
cell projection						

organization						
Regulation of exocytosis	BP	3.53	0.03	0.23	16	17157
Endocytic vesicle	CC	3.53	0.03	0.23	16	30139
Stress-activated protein	BP	3.52	0.03	0.23	18	31098
kinase signaling						
cascade						
Adenyl ribonucleotide	MF	3.52	0.03	0.23	212	32559
binding						
Cellular response to	BP	3.52	0.03	0.23	9	71230
amino acid stimulus						
In utero embryonic	BP	3.51	0.03	0.23	35	1701
development						
Phosphotransferase	MF	3.51	0.03	0.23	107	16773
activity, alcohol group						
as acceptor						
Intracellular organelle	CC	3.51	0.03	0.23	80	70013
lumen						

Neurotrophin TRK	BP	3.51	0.03	0.23	44	48011
receptor signaling						
pathway						
Embryonic organ	BP	3.50	0.03	0.23	21	48568
development						
Regulation of lipase	BP	3.50	0.03	0.23	21	60191
activity						
Skin development	BP	3.50	0.03	0.23	10	43588
Amide biosynthetic	BP	3.50	0.03	0.23	11	43604
process						
Protein kinase activity	MF	3.48	0.03	0.24	91	4672
Regulation of cell-cell	BP	3.48	0.03	0.24	15	22407
adhesion						
Epithelial cell	BP	3.48	0.03	0.24	41	30855
differentiation						
Embryonic limb	BP	3.47	0.03	0.24	17	30326
morphogenesis						

Embryonic appendage	BP	3.47	0.03	0.24	17	35113
morphogenesis						
ATP binding	MF	3.47	0.03	0.24	208	5524
Negative regulation of	BP	3.46	0.03	0.24	7	33144
intracellular steroid						
hormone receptor						
signaling pathway						
Vesicle docking involved	BP	3.46	0.03	0.24	7	6904
in exocytosis						
Glutamate receptor	MF	3.46	0.03	0.24	7	8066
activity						
Ephrin receptor	BP	3.46	0.03	0.24	7	48013
signaling pathway						
Negative regulation of	BP	3.46	0.03	0.24	7	32873
stress-activated MAPK						
cascade						
Negative regulation of	BP	3.46	0.03	0.24	7	70303

stress-activated protein kinase signaling cascade 0.03 38179 Neurotrophin signaling BP 3.45 0.24 44 pathway Transporter activity MF 3.43 0.03 0.24 170 5215 Monosaccharide BP 3.42 0.03 0.25 16 46365 catabolic process Notch signaling pathway 3.41 0.03 0.25 7219 BP 21 Positive regulation of ΒP 3.41 0.03 0.25 13 32874 stress-activated MAPK cascade Positive regulation of ΒP 3.41 0.03 0.25 13 10811 cell-substrate adhesion Organelle localization BP 3.41 0.03 0.25 13 51640 Lysosomal membrane 0.25 26 CC 3.41 0.03 5765

0.03

0.25

212

30554

Adenyl nucleotide

MF

3.40

binding						
Proteinaceous	CC	3.39	0.03	0.25	36	5578
extracellular matrix						
Calcium ion	MF	3.39	0.03	0.25	22	15085
transmembrane						
transporter activity						
Chordate embryonic	BP	3.39	0.03	0.25	35	43009
development						
Endoplasmic reticulum	CC	3.39	0.03	0.25	29	5788
lumen						
Regulation of Ras	BP	3.39	0.03	0.25	29	32318
gtpase activity						
Antigen processing and	BP	3.38	0.03	0.25	31	48002
presentation of peptide						
antigen						
Response to amino acid	BP	3.37	0.03	0.25	15	43200
stimulus						

Positive regulation of	BP	3.37	0.03	0.25	15	10518
phospholipase activity						
Transition metal ion	MF	3.37	0.03	0.25	308	46914
binding						
Negative regulation of	BP	3.37	0.03	0.25	23	32102
response to external						
stimulus						
Regulation of gene	BP	3.37	0.03	0.25	455	10468
expression						
Regulation of embryonic	BP	3.37	0.03	0.25	17	45995
development						
Macromolecule	BP	3.37	0.03	0.25	834	43170
metabolic process						
Cellular macromolecule	BP	3.37	0.03	0.25	355	34645
biosynthetic process						
Polysaccharide	BP	3.37	0.03	0.25	6	272
catabolic process						

Adipose tissue	BP	3.37	0.03	0.25	6	60612
development						
Cardiac cell	BP	3.37	0.03	0.25	6	55006
development						
Homeostasis of number	BP	3.37	0.03	0.25	6	48873
of cells within a tissue						
Negative regulation of	BP	3.37	0.03	0.25	6	51898
protein kinase B						
signaling cascade						
Regulation of	BP	3.37	0.03	0.25	6	32673
interleukin-4 production						
Regulation of RNA	BP	3.36	0.03	0.25	395	2001141
biosynthetic process						
Negative regulation of	BP	3.36	0.03	0.25	9	51148
muscle cell						
differentiation						
Negative regulation of	BP	3.36	0.03	0.25	9	1901185

ERBB signaling						
pathway						
Negative regulation of	BP	3.36	0.03	0.25	9	42059
epidermal growth factor						
receptor signaling						
pathway						
Positive regulation of	BP	3.36	0.03	0.25	9	1901800
proteasomal protein						
catabolic process						
Integral to plasma	CC	3.35	0.04	0.25	145	5887
membrane						
MHC protein complex	CC	3.35	0.04	0.25	10	42611
RNA polymerase activity	MF	3.35	0.04	0.25	10	34062
DNA-directed RNA	MF	3.35	0.04	0.25	10	3899
polymerase activity						
Regulation of organ	BP	3.34	0.04	0.25	25	2000027
morphogenesis						

Glandular epithelial cell	BP	3.34	0.04	0.25	4	2068
development						
Negative chemotaxis	BP	3.34	0.04	0.25	4	50919
Intramolecular	MF	3.34	0.04	0.25	4	16868
transferase activity,						
phosphotransferases						
Positive regulation of	BP	3.34	0.04	0.25	4	45948
translational initiation						
Platelet alpha granule	CC	3.34	0.04	0.25	4	31091
Heart trabecula	BP	3.34	0.04	0.25	4	61384
morphogenesis						
Myelin maintenance	BP	3.34	0.04	0.25	4	43217
Regulation of Wnt	BP	3.34	0.04	0.25	4	2000095
receptor signaling						
pathway, planar cell						
polarity pathway						
Regulation of antigen	BP	3.34	0.04	0.25	4	2577

processing and						
presentation						
Dipeptidyl-peptidase	MF	3.34	0.04	0.25	4	8239
activity						
Positive regulation of	BP	3.34	0.04	0.25	4	10893
steroid biosynthetic						
process						
RNA polymerase core	MF	3.34	0.04	0.25	4	43175
enzyme binding						
Phosphorylation	BP	3.34	0.04	0.25	96	16310
Positive regulation of	BP	3.32	0.04	0.25	5	46622
organ growth						
Translation initiation	MF	3.32	0.04	0.25	5	31369
factor binding						
Nitric oxide mediated	BP	3.32	0.04	0.25	5	7263
signal transduction						
Organ formation	BP	3.32	0.04	0.25	5	48645

Vitamin D receptor	MF	3.32	0.04	0.25	5	42809
binding						
H4/H2A histone	CC	3.32	0.04	0.25	5	43189
acetyltransferase						
complex						
Negative regulation of	BP	3.32	0.04	0.25	5	34122
toll-like receptor						
signaling pathway						
Nucleotide-sugar	BP	3.32	0.04	0.25	5	9226
biosynthetic process						
Regulation of cellular	BP	3.31	0.04	0.25	426	2000112
macromolecule						
biosynthetic process						
Calcium ion binding	MF	3.29	0.04	0.25	99	5509
Positive regulation of	BP	3.29	0.04	0.25	13	70304
stress-activated protein						
kinase signaling						

cascade						
Positive regulation of	BP	3.29	0.04	0.25	13	10863
phospholipase C activity						
Monosaccharide	BP	3.27	0.04	0.25	36	5996
metabolic process						
Basement membrane	CC	3.27	0.04	0.25	15	5604
Hexose catabolic	BP	3.27	0.04	0.25	15	19320
process						
Positive regulation of	BP	3.27	0.04	0.25	7	2000179
neural precursor cell						
proliferation						
Substrate adhesion-	BP	3.27	0.04	0.25	7	34446
dependent cell						
spreading						
Calcium-dependent cell-	BP	3.27	0.04	0.25	7	16339
cell adhesion						
Enhancer sequence-	MF	3.27	0.04	0.25	7	1158

specific DNA binding						
Regulated secretory	BP	3.27	0.04	0.25	7	45055
pathway						
Autophagic vacuole	CC	3.27	0.04	0.25	7	5776
Positive regulation of	BP	3.26	0.04	0.25	12	2000648
stem cell proliferation						
Activation of	BP	3.26	0.04	0.25	12	7202
phospholipase C activity						
Chondroitin sulfate	BP	3.26	0.04	0.25	12	30204
metabolic process						
Calmodulin binding	MF	3.25	0.04	0.25	28	5516
Positive regulation of	BP	3.22	0.04	0.25	8	32755
interleukin-6 production						
Regulation of actin	BP	3.22	0.04	0.25	8	30834
filament						
depolymerization						
Negative regulation of	BP	3.22	0.04	0.25	8	10812

cell-substrate adhesion						
Cell projection	BP	3.21	0.04	0.25	35	48858
morphogenesis						
Actin cytoskeleton	BP	3.20	0.04	0.25	9	31532
reorganization						
Extracellular matrix	MF	3.20	0.04	0.25	9	50840
binding						
Ammonium	BP	3.20	0.04	0.25	2	72488
transmembrane						
transport						
Prostaglandin-E	MF	3.20	0.04	0.25	2	50220
synthase activity						
Trehalose metabolic	BP	3.20	0.04	0.25	2	5991
process						
Seminiferous tubule	BP	3.20	0.04	0.25	2	72520
development						
Cardiac atrium	BP	3.20	0.04	0.25	2	3209

morphogenesis						
Regulation of collateral	BP	3.20	0.04	0.25	2	48696
sprouting in absence of						
injury						
Spemann organizer	BP	3.20	0.04	0.25	2	60061
formation						
Regulation of exocyst	BP	3.20	0.04	0.25	2	60178
localization						
Dehydroascorbic acid	MF	3.20	0.04	0.25	2	33300
transporter activity						
D-glucose	MF	3.20	0.04	0.25	2	55056
transmembrane						
transporter activity						
Production of mirnas	BP	3.20	0.04	0.25	2	35196
involved in gene						
silencing by mirna						
Melatonin receptor	MF	3.20	0.04	0.25	2	8502

activity						
Hypothalamus cell	BP	3.20	0.04	0.25	2	21979
differentiation						
Collagen type IX	CC	3.20	0.04	0.25	2	5594
Negative regulation of	BP	3.20	0.04	0.25	2	10621
transcription by						
transcription factor						
localization						
Detection of	BP	3.20	0.04	0.25	2	9726
endogenous stimulus						
Autonomic nervous	BP	3.20	0.04	0.25	2	48483
system development						
Trophectodermal	BP	3.20	0.04	0.25	2	1831
cellular morphogenesis						
Sec61 translocon	CC	3.20	0.04	0.25	2	5784
complex						
Translocon complex	CC	3.20	0.04	0.25	2	71256

Brain segmentation	BP	3.20	0.04	0.25	2	35284
Cell migration involved	BP	3.20	0.04	0.25	2	3273
in endocardial cushion						
formation						
Transcription initiation	BP	3.20	0.04	0.25	2	6384
from RNA polymerase III						
promoter						
Calcium ion-dependent	BP	3.20	0.04	0.25	2	48791
exocytosis of						
neurotransmitter						
Dendritic growth cone	CC	3.20	0.04	0.25	2	44294
Natriuretic peptide	MF	3.20	0.04	0.25	2	16941
receptor activity						
Metencephalon	BP	3.20	0.04	0.25	2	22037
development						
Diencephalon	BP	3.20	0.04	0.25	2	48852
morphogenesis						

Cardiac vascular	BP	3.20	0.04	0.25	2	60948
smooth muscle cell						
development						
Vasoactive intestinal	MF	3.20	0.04	0.25	2	4999
polypeptide receptor						
activity						
Transcription initiation	BP	3.20	0.04	0.25	2	6391
from mitochondrial						
promoter						
G-quadruplex DNA	MF	3.20	0.04	0.25	2	51880
binding						
Butyrate metabolic	BP	3.20	0.04	0.25	2	19605
process						
Chromosome movement	BP	3.20	0.04	0.25	2	51305
towards spindle pole						
Phenylpropanoid	BP	3.20	0.04	0.25	2	46271
catabolic process						

Cerebral cortex	BP	3.20	0.04	0.25	2	21796
regionalization						
Regulation of defense	BP	3.20	0.04	0.25	2	1900424
response to bacterium						
Regulation of	BP	3.20	0.04	0.25	2	60099
phagocytosis,						
engulfment						
Positive regulation of	BP	3.20	0.04	0.25	2	60100
phagocytosis,						
engulfment						
Cellular pigmentation	BP	3.20	0.04	0.25	2	33059
Synaptic cleft	CC	3.20	0.04	0.25	2	43083
Regulation of nodal	BP	3.20	0.04	0.25	2	1900145
signaling pathway						
involved in						
determination of						
left/right asymmetry						

Regulation of nodal	BP	3.20	0.04	0.25	2	1900175
signaling pathway						
involved in						
determination of lateral						
mesoderm left/right						
asymmetry						
Establishment or	BP	3.20	0.04	0.25	2	45196
maintenance of						
neuroblast polarity						
Establishment of	BP	3.20	0.04	0.25	2	45200
neuroblast polarity						
Sensory system	BP	3.20	0.04	0.25	2	48880
development						
Cardiac muscle cell	BP	3.20	0.04	0.25	2	60379
myoblast differentiation						
Collagen V binding	MF	3.20	0.04	0.25	2	70052
Forebrain neuron fate	BP	3.20	0.04	0.25	2	21877

commitment						
Regulation of	BP	3.20	0.04	0.25	2	21912
transcription from RNA						
polymerase II promoter						
involved in spinal cord						
motor neuron fate						
specification						
Neural plate pattern	BP	3.20	0.04	0.25	2	60896
specification						
Neural plate	BP	3.20	0.04	0.25	2	60897
regionalization						
Synaptic vesicle	BP	3.20	0.04	0.25	2	97091
clustering						
Bombesin receptor	MF	3.20	0.04	0.25	2	4946
activity						
Locomotion involved in	BP	3.20	0.04	0.25	2	31987
locomotory behavior						

Regulation of	BP	3.20	0.04	0.25	2	1900094
transcription from RNA						
polymerase II promoter						
involved in						
determination of						
left/right symmetry						
Nodal signaling pathway	BP	3.20	0.04	0.25	2	1900164
involved in						
determination of lateral						
mesoderm left/right						
asymmetry						
Nodal signaling pathway	BP	3.20	0.04	0.25	2	38107
involved in						
determination of						
left/right asymmetry						
Smooth muscle cell	BP	3.20	0.04	0.25	2	48659
proliferation						

Pre-mrna intronic	MF	3.20	0.04	0.25	2	97157
binding						
Intestinal epithelial cell	BP	3.20	0.04	0.25	2	60576
development						
Negative regulation of	BP	3.20	0.04	0.25	2	45541
cholesterol biosynthetic						
process						
Negative regulation of	BP	3.20	0.04	0.25	2	90206
cholesterol metabolic						
process						
Actin polymerization-	BP	3.20	0.04	0.25	2	70358
dependent cell motility						
Protein poly-ADP-	BP	3.20	0.04	0.25	2	70212
ribosylation						
Viral assembly,	BP	3.20	0.04	0.25	2	19067
maturation, egress, and						
release						

Retinal blood vessel	BP	3.20	0.04	0.25	2	61304
morphogenesis						
Ornithine decarboxylase	MF	3.20	0.04	0.25	2	42978
activator activity						
Intestine smooth muscle	BP	3.20	0.04	0.25	2	14827
contraction						
Calcium-dependent	MF	3.20	0.04	0.25	2	4698
protein kinase C activity						
Calcium-dependent	MF	3.20	0.04	0.25	2	9931
protein serine/threonine						
kinase activity						
Laminin-11 complex	CC	3.20	0.04	0.25	2	43260
Positive regulation of	BP	3.19	0.04	0.25	69	51345
hydrolase activity						
Regulation of MAPK	BP	3.19	0.04	0.25	71	43408
cascade						
Ubiquitin protein ligase	MF	3.19	0.04	0.25	26	31625

binding						
Small conjugating	MF	3.19	0.04	0.25	26	44389
protein ligase binding						
Transcription initiation	BP	3.19	0.04	0.25	31	6367
from RNA polymerase II						
promoter						
Regulation of purine	BP	3.18	0.04	0.25	20	1900371
nucleotide biosynthetic						
process						
Regulation of nucleotide	BP	3.18	0.04	0.25	20	30808
biosynthetic process						
Transcriptional	CC	3.18	0.04	0.25	13	17053
repressor complex						
Ion transport	BP	3.17	0.04	0.25	130	6811
Regulation of	BP	3.17	0.04	0.25	391	6355
transcription, DNA-						
dependent						

Positive regulation of	BP	3.17	0.04	0.25	15	1900544
purine nucleotide						
metabolic process						
Negative regulation of	BP	3.17	0.04	0.25	15	51100
binding						
Wnt receptor signaling	BP	3.16	0.04	0.25	6	60071
pathway, planar cell						
polarity pathway						
Regulation of	BP	3.16	0.04	0.25	6	90175
establishment of planar						
polarity						
Negative regulation of	BP	3.16	0.04	0.25	6	45686
glial cell differentiation						
Motor neuron axon	BP	3.16	0.04	0.25	6	8045
guidance						
Laminin binding	MF	3.16	0.04	0.25	6	43236
Regulation of cytokine	BP	3.15	0.04	0.25	3	2739

secretion involved in						
immune response						
Galactose catabolic	BP	3.15	0.04	0.25	3	19388
process						
Filtration diaphragm	CC	3.15	0.04	0.25	3	36056
Slit diaphragm	CC	3.15	0.04	0.25	3	36057
Relaxation of smooth	BP	3.15	0.04	0.25	3	44557
muscle						
Embryonic process	BP	3.15	0.04	0.25	3	60136
involved in female						
pregnancy						
Regulation of neuron	BP	3.15	0.04	0.25	3	2001222
migration						
Cellular extravasation	BP	3.15	0.04	0.25	3	45123
Interkinetic nuclear	BP	3.15	0.04	0.25	3	22027
migration						
CD4-positive or CD8-	BP	3.15	0.04	0.25	3	43369

positive, alpha-beta T						
cell lineage commitment						
H3 histone	CC	3.15	0.04	0.25	3	70775
acetyltransferase						
complex						
MOZ/MORF histone	CC	3.15	0.04	0.25	3	70776
acetyltransferase						
complex						
Trna transcription	BP	3.15	0.04	0.25	3	9304
Neuronal cell body	CC	3.15	0.04	0.25	3	32809
membrane						
Cell body membrane	CC	3.15	0.04	0.25	3	44298
MAP-kinase scaffold	MF	3.15	0.04	0.25	3	5078
activity						
Positive regulation of	BP	3.15	0.04	0.25	3	2606
dendritic cell antigen						
processing and						

presentation						
Nuclear envelope lumen	CC	3.15	0.04	0.25	3	5641
Laminin-1 complex	CC	3.15	0.04	0.25	3	5606
Cell cortex part	CC	3.15	0.04	0.25	18	44448
Enzyme activator	MF	3.15	0.04	0.25	64	8047
activity						
Protein secretion	BP	3.14	0.04	0.26	12	9306
Regulation of cell	BP	3.13	0.04	0.26	61	30334
migration						
Regulation of	BP	3.12	0.04	0.26	16	1505
neurotransmitter levels						
Visual perception	BP	3.12	0.04	0.26	30	7601
Dendrite	CC	3.12	0.04	0.26	30	30425
Detection of external	BP	3.12	0.04	0.26	26	9581
stimulus						
Monocarboxylic acid	BP	3.12	0.04	0.26	28	72330
biosynthetic process						

Central nervous system	BP	3.10	0.04	0.26	20	7417
development						
Activating transcription	MF	3.10	0.04	0.26	11	33613
factor binding						
Negative regulation of	BP	3.09	0.05	0.27	17	90288
cellular response to						
growth factor stimulus						
G-protein coupled	MF	3.09	0.05	0.27	21	8528
peptide receptor activity						
E-box binding	MF	3.08	0.05	0.27	7	70888
Myeloid cell activation	BP	3.08	0.05	0.27	7	2275
involved in immune						
response						
Vesicle docking	BP	3.08	0.05	0.27	7	48278
Excretion	BP	3.07	0.05	0.27	10	7588
Negative regulation of	BP	3.07	0.05	0.27	5	10894
steroid biosynthetic						

process						
SMAD protein signal	BP	3.07	0.05	0.27	5	60395
transduction						
External encapsulating	CC	3.07	0.05	0.27	5	44462
structure part						
RNA polymerase II	MF	3.07	0.05	0.27	5	980
distal enhancer						
sequence-specific DNA						
binding						
Mannosidase activity	MF	3.07	0.05	0.27	5	15923
Positive regulation of	BP	3.07	0.05	0.27	5	2052
neuroblast proliferation						
Protein phosphatase 2A	MF	3.07	0.05	0.27	5	51721
binding						
Histone	MF	3.07	0.05	0.27	5	35035
acetyltransferase						
binding						

Regulation of alpha-	BP	3.07	0.05	0.27	5	2000311
amino-3-hydroxy-5-						
methyl-4-isoxazole						
propionate selective						
glutamate receptor						
activity						
Protein targeting to	BP	3.07	0.05	0.27	5	72661
plasma membrane						
Muscle cell proliferation	BP	3.07	0.05	0.27	5	33002
Positive regulation of	BP	3.07	0.05	0.27	15	45981
nucleotide metabolic						
process						
Cellular polysaccharide	BP	3.07	0.05	0.27	13	44264
metabolic process						
Potassium ion transport	BP	3.06	0.05	0.27	18	6813
Lateral plasma	CC	3.06	0.05	0.27	8	16328
membrane						

Defense response to	BP	3.06	0.05	0.27	8	50830
Gram-positive bacterium						
Positive regulation of	BP	3.06	0.05	0.27	8	45921
exocytosis						
Receptor regulator	MF	3.06	0.05	0.27	9	30545
activity						
Cardiocyte	BP	3.06	0.05	0.27	9	35051
differentiation						
Forelimb	BP	3.06	0.05	0.27	9	35136
morphogenesis						
Positive regulation of	BP	3.06	0.05	0.27	9	50772
axonogenesis						
Phagocytosis	BP	3.06	0.05	0.27	9	6909
Cell-cell adhesion	BP	3.06	0.05	0.27	54	16337
Transcription regulatory	MF	3.05	0.05	0.27	28	976
region sequence-						
specific DNA binding						

Regulation of	BP	3.05	0.05	0.27	434	10556
macromolecule						
biosynthetic process						
Chemical homeostasis	BP	3.04	0.05	0.27	107	48878
Oxidation-reduction	BP	3.04	0.05	0.27	73	55114
process						
Leydig cell	BP	3.04	0.05	0.27	4	33327
differentiation						
Trabecula	BP	3.04	0.05	0.27	4	61383
morphogenesis						
Positive regulation of	BP	3.04	0.05	0.27	4	45078
interferon-gamma						
biosynthetic process						
Positive regulation of	BP	3.04	0.05	0.27	4	43568
insulin-like growth factor						
receptor signaling						
pathway						

Mitogen-activated	MF	3.04	0.05	0.27	4	31434
protein kinase kinase						
binding						
Regulation of non-	BP	3.04	0.05	0.27	4	2000050
canonical Wnt receptor						
signaling pathway						
Chronic inflammatory	BP	3.04	0.05	0.27	4	2544
response						
Regulation of SMAD	BP	3.04	0.05	0.27	4	60390
protein import into						
nucleus						
Regulation of steroid	BP	3.04	0.05	0.27	4	90030
hormone biosynthetic						
process						
Histone H2A acetylation	BP	3.04	0.05	0.27	4	43968
Negative regulation of	BP	3.04	0.05	0.27	4	43508
JUN kinase activity						

Fibrillar collagen	CC	3.04	0.05	0.27	4	5583
Cellular response to	BP	3.03	0.05	0.27	214	70887
chemical stimulus						
Regulation of	BP	3.03	0.05	0.27	16	10517
phospholipase activity						
Positive regulation of	BP	3.02	0.05	0.27	39	40017
locomotion						
Fatty acid biosynthetic	BP	3.02	0.05	0.27	20	6633
process						
Telomere maintenance	BP	3.02	0.05	0.27	12	723
Steroid metabolic	BP	3.01	0.05	0.27	38	8202
process						
Golgi cisterna	CC	3.01	0.05	0.27	14	32580
membrane						
Peptide receptor activity	MF	3.01	0.05	0.27	21	1653
Epidermal growth factor	BP	3.00	0.05	0.27	31	7173
receptor signaling						

pathway						
Regulation of cytokine	BP	3.00	0.05	0.27	17	42035
biosynthetic process						
Positive regulation of	BP	3.00	0.05	0.27	30	43410
MAPK cascade						
Cellular chemical	BP	3.00	0.05	0.27	80	55082
homeostasis						
Nervous system	BP	2.99	0.05	0.27	44	7399
development						
Endopeptidase inhibitor	MF	2.99	0.05	0.28	28	4866
activity						
Endoplasmic reticulum-	CC	2.98	0.05	0.28	11	5793
Golgi intermediate						
compartment						
Extrinsic to internal side	CC	2.98	0.05	0.28	11	31234
of plasma membrane						
Regulation of striated	BP	2.97	0.05	0.28	15	51153

muscle cell						
differentiation						
Regulation of interferon-	BP	2.96	0.05	0.28	13	32649
gamma production						
Positive regulation of	BP	2.96	0.05	0.28	6	32731
interleukin-1 beta						
production						
Calcium activated cation	MF	2.96	0.05	0.28	6	5227
channel activity						
Regulation of	BP	2.96	0.05	0.28	6	48713
oligodendrocyte						
differentiation						
Neural tube	BP	2.96	0.05	0.28	6	21915
development						
Phosphatidylinositol 3-	MF	2.96	0.05	0.28	6	43548
kinase binding						
Middle ear	BP	2.96	0.05	0.28	6	42474


BP	2.96	0.05	0.28	6	44089
BP	2.95	0.05	0.28	91	19725
BP	2.94	0.05	0.28	31	38127
BP	2.94	0.05	0.29	36	70727
BP	2.93	0.05	0.29	21	70372
BP	2.93	0.05	0.29	110	51336
BP	2.93	0.05	0.29	101	10627
MF	2.93	0.05	0.29	22	30674
	BP BP BP	BP 2.95 BP 2.94 BP 2.93 BP 2.93 BP 2.93	BP 2.95 0.05 BP 2.94 0.05 BP 2.94 0.05 BP 2.93 0.05 BP 2.93 0.05 BP 2.93 0.05	BP 2.95 0.05 0.28 BP 2.94 0.05 0.29 BP 2.93 0.05 0.29 BP 2.93 0.05 0.29 BP 2.93 0.05 0.29	BP 2.95 0.05 0.28 91 BP 2.94 0.05 0.28 31 BP 2.94 0.05 0.29 36 BP 2.93 0.05 0.29 21 BP 2.93 0.05 0.29 110 BP 2.93 0.05 0.29 101

Response to oxygen-	BP	2.92	0.05	0.29	135	1901700
containing compound						
Ventricular cardiac	BP	2.92	0.05	0.29	9	55010
muscle tissue						
morphogenesis						
Neuron-neuron synaptic	BP	2.92	0.05	0.29	9	7270
transmission						
Hindlimb	BP	2.92	0.05	0.29	9	35137
morphogenesis						
Regulation of cellular	BP	2.92	0.05	0.29	26	90287
response to growth						
factor stimulus						
Homophilic cell	BP	2.92	0.05	0.29	24	7156
adhesion						
Cellular response to	BP	2.92	0.05	0.29	184	71310
organic substance						
Protein tyrosine kinase	MF	2.92	0.05	0.29	25	4713


activity						
Locomotory behavior	BP	2.92	0.05	0.29	25	7626
Receptor activator	MF	2.91	0.05	0.29	7	30546
activity						
Negative regulation of	BP	2.91	0.05	0.29	7	14014
gliogenesis						
Golgi-associated vesicle	CC	2.91	0.05	0.29	7	5798
Positive regulation of	BP	2.91	0.05	0.29	14	45862
proteolysis						
Extracellular ligand-	MF	2.91	0.05	0.29	14	5230
gated ion channel						
activity						
Telomere organization	BP	2.91	0.05	0.29	12	32200
Negative regulation of	BP	2.91	0.05	0.29	8	30514
BMP signaling pathway						
Frizzled binding	MF	2.91	0.05	0.29	8	5109
Delayed rectifier	MF	2.91	0.05	0.29	8	5251

potassium channel activity BP, Biological process; MF, molecular function; cellular component; #, number of; GO ID, gene ontology identification.









Ε

