



Prevalence of Malnutrition among People Living with Obesity and Hypertension

Peter M. Babangida ^{a,b,c,*}, Esegbue Peters ^a, Charles Ofili ^a
and Ezekiel U Nwose ^{a,d}

^a Department of Community and Public Health, Novena University Ogume, Delta State, Nigeria.

^b Department of Dietetics, Irrua Specialist Teaching Hospital, Irrua Edo State, Nigeria.

^c Department of Human Nutrition and Dietetics, Ambrose Alli University Ekpoma, Edo State, Nigeria.

^d School of Health and Medical Sciences, University of Southern Queensland, Australia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98864>

Systematic Review Article

Received: 15/02/2023

Accepted: 19/04/2023

Published: 25/04/2023

ABSTRACT

Background: Obesity is a growing public health problem and should be kept in mind when treating patients with hypertension because it is a strong predictor of uncontrolled hypertension. As obesity and its metabolic and vascular effects remained during the years, research has provided evidence justifying the management of metabolic components in obese individuals. The purpose of this review is to provide more evidence justifying the management of metabolic problems in obese hypertensive individuals.

Conclusion: Obesity is a multi factorial pathology and chronic diet related non-communicable disease. The most widely used parameter for its diagnosis is the body mass index (BMI) which is not suitable for assessing the body fat and other metabolic components. Obesity specifically abdominal obesity is an independent risk factor for hypertension.

*Corresponding author: E-mail: malachypeter18@gmail.com;

Recommendation: Anthropometrics like the waist circumference, waist to height ratio (WHtR), waist-hip ratio (WHR) and metabolic/biochemical components should be considered in the diagnosis of obesity.

Keywords: Prevalence; malnutrition; metabolic syndrome; obesity; hypertension.

1. INTRODUCTION

“Given the increase of sedentary lifestyle and bad diet in our life, the prevalence of obesity and hypertension has remarkably increased among most population in the last years [1]. Obesity is a growing public health problem and should be kept in mind when treating patients with hypertension because it is a strong predictor of un-controlled blood pressure” [1]. “As obesity and its metabolic and hemodynamic consequences are sustained during many years, renal injury gradually makes the hypertension more severe and more resistance to therapy” [2]. “Although weight loss is helpful in managing hypertension, many obese patients are unable to sustain adequate weight loss through lifestyle modifications and there are few available drugs that safely and effectively produce adequate long-term weight loss. Current therapeutic approaches are, therefore, aimed mainly at treating the hypertension and metabolic consequences of obesity, including diabetes mellitus, dyslipidemia, and inflammation” [3].

The obesity scourge began in the 1970s, at first in the United States and afterwards spreading globally. Roughly 60-80% of adults in many Western countries are becoming overweight or obese, which is novel in human history [4]. A recent study by Jennifer et al. reveals that obesity was independently associated with hypertension in population-based controls [5]. “There is a high rate of unrecognized hypertension and obesity among Bus Drivers from different regions which was connected with individual lifestyle and behaviors” [6]. “Improved awareness through educational and screening activities will initiate lifestyle modifications that will reduce the onset of cardio-metabolic disorders and give ideas for better disease prognosis, prevention and personalized medical care” [6]. “Hypertension is the commonest risk factor for cardiovascular diseases and it is associated with many morbidities and mortalities worldwide” [7].

“Physiological researches and epidemiological data have suggested a potential inverse

relationship between height and blood pressure in adult” [8]. “Cogent data exist connecting increased height and leg length with normotension. These findings identify shorter individuals as having higher risk with regard to hypertension and cardiovascular disease” [8,9]. Substantial epidemiological studies have revealed that high blood pressure is more than twice present in those who are obese than in lean subjects. Furthermore, the values of blood pressure and body weight have a direct association at any age of life [10,11]. The growing prevalence of chronic diseases linked to obesity endangers both human lives and economies [12]. The goal is prevention today, given the enormous expenses of obesity (both public and personal) and how challenging it is to shed weight once one is obese [13].

Additionally, there are also differences in obesity-related hypertension according to sex, race, and ethnicity. Obesity-related public health and policy developments are required to lower morbidity and mortality [14]. In obese females, hypertension and dyslipidemia usually coexisted with other illnesses. Diabetes and dyslipidemia were the two main illnesses in the co-morbidity network in obese males. Results show that managing metabolic components in obese people is justified. The findings will aid in prioritizing strategies for reducing co-morbidity as a public health objective [15].

2. METHODS

This was a review study. Journal articles and publications from 2004 to 2022 were reviewed. PubMed, Google Scholar, Science Direct, and Cochrane were some of the search engines used. Search terms included prevalence, malnutrition, metabolic syndrome, obesity and hypertension. A total of 150 papers were retrieved during the search; 51 were included in this review and 41 of the papers were on nutrition-related issues.

Global: Malnutrition is a global problem that no nation can afford to ignore. 39% of the world's adults are obese (2018 Global Nutrition Report).

Beyond health, the social and economic growth of nations is being impacted by the poor progress on malnutrition. According to estimates, all types of malnutrition might cost society up to US\$3.5 trillion annually, with obesity alone costing US\$500 billion [16]. According to the 2018 Global Nutrition Report, more illnesses are caused by malnutrition than by any other factor [16]. According to WHO, “an energy imbalance between calories taken and calories expended is the primary cause of obesity” [17]. “Globally, there has been a rise in the consumption of calorie-dense foods that are high in fat and sugar, as well as a rise in physical inactivity as a result of the sedentary nature of many occupations, evolving transportation options, and escalating urbanization. Body mass index (BMI) is a straightforward measure of weight in relation to height that is frequently used to categorize adult overweight and obesity. It is determined by dividing the individual's weight in kilograms by the square of his or her height in meters (kg/m²). WHO defines obesity in adults as having a BMI of 30 or higher” [17]. “As BMI rises, the risk for these non-communicable diseases also rises” [17]. “An estimated four million people die each year as a result of obesity and hypertension's negative health effects” [18].

individuals on earth reside in nations where obesity kills more people than underweight [17]. “The leading cause of lost years of life with a handicap in the world is hypertension. According to the American Heart Association (AHA), hypertension affects 86 million adults in the US, or 34% of the population. The AHA defines hypertension as having a systolic blood pressure (SBP) of 140 mm Hg or higher, or a diastolic blood pressure (DBP) of 90 mm Hg or higher, taking antihypertensive medication, or being identified as having hypertension by a doctor at least twice” [20]. Hypertension affects most people over 60 in the population [21]. The allocation of healthcare resources for the condition and associated co-morbidities is predicted to approach \$150 billion annually in the United States, making obesity and hypertension the next big epidemiologic challenge facing today's clinicians [22]. According to a different study by Omair and Travis, “obesity is a major global public health issue and is intrinsically connected to poor cardiovascular outcomes” [23]. “It is commonly known that an increase in blood pressure is correlated with an increase in body fat, and that 65-78% of primary hypertension cases are thought to be attributable to obesity” [23]. An estimated one billion individuals worldwide are affected by high blood pressure, which continues to be a significant contributor to the burden of non-communicable diseases and death internationally [24,25].

Since 1975, the global rate of obesity has nearly tripled. Over 650 million persons, or 13% of adults, were obese in 2016. The majority of




Fig. 1. Shows the prevalence of hypertension and hypertension caused by obesity in various age groups [19]

At a median follow-up of 9.5 years in a study on cardiovascular disease that evaluated intakes of white rice, whole grains, and refined grains, 9.2% of the participants experienced a primary outcome that was a composite of death and major cardiovascular events (cardiovascular death, nonfatal myocardial infarction, heart failure, or stroke). A higher risk was linked to consuming more refined grains [26].

“Unhealthy weight gain is a major contributor to hypertension, accounting for 65% to 75% of the risk for human primary (essential) hypertension, especially when coupled with increasing visceral adiposity. Pressure natriuresis is impaired by increased renal tubular sodium re-absorption, which is a major factor in the development of obesity-related hypertension” [27].

Limiting total fat and sugar intake, increasing fruit and vegetable consumption, as well as legume, whole grain, and nut consumption, and engaging in regular physical activity (60 minutes per day for children, and 150 minutes spread throughout the week for adults) are all effective ways to prevent disease on an individual level. This is made feasible by the persistent use of population-based, evidence-based policies that enable everyone, including the poorest people, to have regular access to healthy eating and physical activity. One such measure is a charge on beverages with added sugars [17].

“Reduced fat, sugar, and salt content in processed foods, availability of affordable, healthy food options for all consumers, restrictions on marketing of foods high in sugar, salt, and fats, especially those marketed to children and teenagers, availability of healthy food options, and encouragement of regular physical activity in the workplace are all ways that the industry can prevent obesity and hypertension” [17,28].

Africa: Obesity and hypertension are significant public health issues that have a significant impact on the rising global trend in morbidity and early mortality [29]. From 4.5% in 2000 to 7% in 2010 and up to 29.2% (28.8%-29.7%) by 2025, the worldwide burden of disease attributed to hypertension has dramatically increased. Africa has the highest prevalence of hypertension compared to other WHO regions, with a 46% total prevalence in persons aged 25 and older [29]. The risk of hypertension and other cardiovascular illnesses is significantly increased by obesity. Its prevalence is increasing

worldwide, and current estimates indicate that 20% to 50% of metropolitan African residents are either overweight or obese. Consequently, two of the most significant risk factors for morbidity and mortality in Africa are hypertension and obesity [29].

Age-standardized prevalence of hypertension in rural Nigeria was 19.3% (95% CI: 17.3-21.3), in rural Kenya it was 21.4% (19.8-23.0), in urban Tanzania it was 23.7% (21.3-26.2), and in urban Namibia it was 38.0% (35.9-40.1). These findings come from research by Akpa et al [29]. “The percentage of people with grade 2 or grade 3 hypertension (180/110 mmHg) ranged from 29.2% (Namibia) to 43.3% (Nigeria) in those with hypertension” [31]. Hendriks et al. reported that “the prevalence of hypertension control varied from 2.6% in Kenya to 17.8% in Namibia. Obesity prevalence (BMI 30) ranged from 6.1% in Nigeria to 17.4% in Tanzania, and BMI independently predicted blood pressure level in all study populations along with age and gender” [31].

The two most significant modifiable risk factors for cardiovascular illnesses, obesity and hypertension, according to Jennifer et al., are not well understood in Africa [5]. The leading cause of death in 2012 was cardiovascular disease, which is mostly heart disease and stroke. Obesity and hypertension are significant risk factors for cardiovascular illnesses. Hypertension is a serious public health concern globally [32]. Once uncommon in traditional African civilizations, hypertension has become a significant public health issue due to an increase in its risk factors [33]; being the main reason for admissions and deaths in Ghana, it is a serious health issue there [34]. In Ghana, hypertension affects more than one in four persons. Both rural and urban populations share this high incidence, which has persisted for decades. If Ghana is to attain the worldwide target for hypertension, additional investments in cardiovascular health are necessary given the low awareness and poor control of hypertension [34]. Currently, 46% of the population in low- and middle-income nations and sub-Saharan Africa (SSA) suffers from hypertension. Subclinical disease, often known as inadequate health, is more concerning. This condition is marked by a sense of physical weakness and a lack of vitality, and is viewed as a transitional state between health and chronic diseases like hypertension [6]. Africans who were obese had a more than twofold increased risk of having hypertension, and the risk rose with age.

The prevalence of hypertension is rising globally, particularly in Sub-Saharan African nations. In addition to being the largest risk factor for mortality from cardiovascular diseases globally, hypertension is also the leading risk factor for premature death and disability [35]. among Eastern Sudan, hypertension is prevalent, especially among older and obese people. Prevention strategies, like dietary changes, should be used [35].


Southeast Asian nations have seen a sharp rise in the global prevalence of hypertension during the past 20 years. A thorough analysis was done to determine the prevalence of hypertension and associated risk factors among Southeast Asian metropolitan populations. It was discovered that

male gender, ethnicity, socioeconomic status, education, waist circumference, smoking, and dyslipidemia were common risk factors. Activities for primary and secondary prevention are urgently needed, according to the study. To enhance the general health outcomes of all populations in all Southeast Asian countries, a multisectoral and intersectoral approach and collaboration should be used [36]. It is necessary to view obesity as a true disorder. One of the main global public health problems linked to higher morbidity and mortality rates is obesity. Unfortunately, despite various educational initiatives, the battle against obesity does not appear to be successful: in recent years, the incidence kept rising [37].



Regional patterns and the association between obesity and hypertension in Africa: the H3Africa CHAIR study




- Regional differences in age-adjusted proportions
- Relatively higher in western Africa




- Proportion of hypertension (≥ 140/90 mmHg) across BMI
- Proportion increases with increasing BMI



- Adjusted odds of hypertension (≥ 140/90mmHg (def1) and ≥130/80mmHg (def2)) in obesity and across age categories
- Obesity doubles the odds of hypertension
- Adjusted odds of hypertension in obesity increases with increasing age



- Regional differences in age-adjusted proportions
- Relatively higher in Southern Africa



- Proportion of hypertension (≥ 130/80mmHg) across BMI
- Proportion increases with increasing BMI

Fig. 2. Obesity and hypertension in Africa: regional patterns and association [30]




Fig. 3. Shows the prevalence of hypertension in a rural population in the state of Edo in relation to body mass index [40]

Nigeria and Edo State: According to a study by Chukwuonye et al., obesity and overweight are currently very common among urban adults in Nigeria. The study also provides solid data that may be used to highlight the urgent need to reinforce and put preventive measures into place in order to shift the paradigm [38]. “The combined crude prevalence rates of obesity and overweight in Nigeria were 25.0% (95% confidence interval, CI: 20.4-29.6) and 14.3% (95% CI: 12.0-15.5), respectively, based on 35 studies (n = 52,816). Women were more likely than men to be overweight or obese, with prevalence rates of 25.5% (95% CI: 17.1-34.0) versus 25.2% (95% CI: 18.0-32.4) and 19.8% (95% CI: 3.9-25.6) versus 12.9% (95% CI: 9.1-16.7), respectively. Waist circumference and body mass index (BMI) were combined to yield mean values of 25.6 kg/m² and 86.5 cm, respectively. In 2020, we predicted that there will be 21 million and 12 million overweight and obese people in Nigeria, representing an age-adjusted prevalence of 20.3% and 11.6%, respectively. Overweight and obesity were more prevalent than usual rates” [39].

A higher prevalence of obesity and hypertension is associated with negative cardiovascular outcomes [41]. “A secondary analysis of data was conducted for 5135 participants aged ≥ 16 years from community-based hypertension prevalence study to determine the prevalence of obesity and association between multiple anthropometric indices and blood pressure (BP).

The indices were waist circumference (WC), body mass index (BMI), waist-to-height ratio (WHtR), waist-to-hip ratio (WHR), a body shape index (ABSI), abdominal volume index (AVI), body adiposity index (BAI), body roundness index (BRI), visceral adiposity index (VAI) and conicity index (CI). A statistical analysis was performed to determine the association, predictive ability, cutoff values and independent determinants of hypertension. Crude prevalence of obesity was 136 per 1000 (95% confidence interval 126–146). BMI had the strongest correlation with systolic and diastolic BP ($r = 0.260$ and 0.264 , respectively). Indices of central adiposity (AVI, WC, WHtR, BRI) were the strongest predictors of hypertension ($\geq 140/90$ mmHg), and their cut-off values were generally higher in females than males. WHR, age, BMI and CI were independent determinants of hypertension ≥ 140 mmHg ($p < 0.05$). The study concluded that measures of central adiposity are the strongest predictors and independent determinants of hypertension in the studied population” [41].

Obesity and overweight were prevalent at 13.8% and 9.4%, respectively. In obese versus normal BMI females, the prevalence of hypertensive range systolic blood pressure was 16% versus 23% ($p=0.00$), while in men, it was 12.1% versus 6.4% ($p=0.27$). In obese versus normal BMI females, the prevalence of hypertensive range diastolic BP was 12% versus 1.4% ($p=0.00$), while in men, it was 15.2% versus 3.5%

($p=0.01$). The pre-hypertension and hypertensive range systolic blood pressure in overweight ($P = 0.01$, $P = 0.002$) and obese ($P = 0.00$, $P = 0.00$) patients, as well as the hypertensive range diastolic blood pressure in only obese subjects, were significantly correlated with BMI in group B. Obesity and pre-hypertensive range diastolic blood pressure had the only statistically significant link in group A ($P = 0.00$) [42].

In rural areas, women and men's cardiovascular risk factors for hypertension and obesity differ significantly [43]. Nigeria's targeted public health response to obesity is comparatively underdeveloped because of the lack of epidemiologic knowledge [44]. The prevalence of hypertension was 33.1% overall (males 36.8%, females 31.1%). 11.1% of people self-reported having high blood pressure, whereas 5.1% were currently taking anti-hypertensive medication. The average age of the responders was 38.8 15.6 years. In the categories of underweight, normal, overweight, and obese, the respondents' body mass indices were 5.2%, 52.0%, 29.5%, and 13.3%, respectively. 11.5% and 3.2%, respectively, reported using alcohol or cigarettes. Binary logistic regression analysis showed a significant association between hypertension and age groups 30-49 years (OR 2.258, 95% CI: 1.311-3.884), 50 years (OR 7.145, 95% CI: 3.644-14.011), and being overweight or obese (OR 2.281, 95% CI: 1.022-5.088). Being underweight was negatively correlated with having hypertension (OR 0.537, 95% CI: 0.395-0.832) [45].

The study also reveals that anthropometric indices of central adiposity and BMI are both significant risk factors for the development of hypertension. It was suggested that clinical treatment should routinely use these anthropometric measures of central adiposity (WHR, CI, AVI, WC, WHtR) rather than too relying on BMI alone because they may be more accurate predictors of hypertension. The findings also emphasized the relative imprecision of cutoffs and cautioned against using population-specific ranges or cutoff values as a baseline for implementing lifestyle changes in practice rather than population-specific ranges or cutoff values. Therefore, it was accepted that additional confirmation of the findings in the context of population [38].

3. RESULTS

The result from this review revealed that the incidence of obesity has served as the primary

impetus for study into the processes that underlie this illness and the ailments that it is associated with. As the situation gets worse, obesity-related diseases including diabetes, heart disease, and cancer start to appear more frequently. Individuals who had considerably higher values for their weight, body mass index, waist to height ratio, hip circumference, fat mass, and fat-free mass also had hypertension compared to those with normotension, which is proof that obesity and hypertension are related. It was observed that in developing nations, hypertension and excess body weight are two key risk factors for cardiovascular morbidity and mortality. Age, regular alcohol use, physical activity level, obesity, stress, and fasting blood glucose concentration in blood are some risk factors linked to the development of hypertension in adults.

4. DISCUSSION

The majority of economically developed nations have experienced a steady and rapid rise in obesity rates over the past ten years, which has served as the main driver for study into the pathophysiological mechanisms behind this disease and its associated illnesses [37]. Lower health costs for primary and secondary prevention of the most common degenerative diseases linked to obesity will result from the consequences of an accurate and early identification of the condition [37]. As a result, the fundamental risk is directly proportionate to the absolute therapeutic benefit. In order to minimize the negative effects of under- and overdiagnosis, there is increasing interest worldwide in early obesity diagnosis. As a result, the condition will worsen and an increase in obesity-related pathologies like diabetes, cardiovascular disease, and cancer will occur [37].

The clustering of obesity and other metabolic syndrome symptoms may have significant preventative implications, particularly in relation to whether visceral obesity-specific therapies might reduce cardiovascular and renal morbidity [46]. Incidence and prevalence of idiopathic intracranial hypertension are rising sharply in tandem with rises in body mass index in the population. Given the comorbidities, difficulties, and higher use of healthcare services linked to idiopathic intracranial hypertension, this has significant implications for health care providers and policy makers [47].

Estimating how much obesity is increasing in the population and how that affects incident

hypertension has significant consequences for public health policy [14]. The direct correlation between obesity and hypertension was supported by participants who had hypertension compared to those with normotension and had significantly greater weight, body mass index, waist, hip, and fat mass values [48].

Major risk factors for cardiovascular morbidity and mortality in emerging nations include hypertension and excess body weight. "According to study, obesity is very common in Nigeria. This is especially noticeable in metropolitan Nigeria and among women, which may be partly explained by a generalized sedentary lifestyle and an increase in fast food restaurants, which is a general tendency in many African settings" [44]. "According to all research, Umuahia, Abia State, South-east Nigeria has the highest prevalence of obesity in 2012, at 33.7% [49], with Kano, North-West Nigeria, recording the lowest rate in 2013 at 0.84%" [50]. Age, regular alcohol use, amount of physical activity, obesity, stress, and fasting blood glucose levels are some risk factors linked to the development of hypertension (HTN) in adults. More than 60% of study participants practiced healthy eating habits, exercise, and understanding of hypertension [33]. Similar to reports from other suburban communities, Irrua's obesity rate is slightly higher than anticipated for the area [12]. This can be the result of residents' lifestyles gradually changing. The study's obesity indicators revealed a positive relationship between co-morbidities like hypertension and stroke, the world's top cause of death and disability and one in which hypertension is the most common risk factor [51,12]. According to this study, obesity and its co-morbidities are becoming a bigger issue in this area [12].

5. CONCLUSION

Obesity is a multi-factorial pathology and chronic diet related non-communicable disease. Body mass index (BMI), the most frequently used metric for its diagnosis, is ineffective for determining body fat. Numerous researches show that using BMI alone to determine obesity is ineffective. Obesity, particularly abdominal obesity, increases the risk of developing and progressing hypertension independently. The majority of studies on the relationship between body mass index (BMI) and hypertension, a prevalent health problem, focused on this relationship.

The regulation of blood pressure is positively impacted by increased physical activity, quitting smoking and drinking, eating more fruits and vegetables, and consuming less red and assorted meats, fats, highly processed and refined foods.

6. RECOMMENDATION

Health talks on diet, frequent exercise, and screening are advised.

Sensitization of the population on hypertension and its complications is important.

Along with recommendations to lose weight, there should be specific guidelines for treating obesity-related hypertension.

Weight-reduction interventions are essential for reducing the prevalence of hypertension.

Other parameters like the waist-hip ratio (WHR), waist-height ratio (WHtR) and waist circumference should be considered in the diagnosis of obesity which has been found to have a strong association with systolic and diastolic blood pressures.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hammami R, Bahloul A, Triki S, Charfeddine S, Triki F, Zakhama L, et al. Impact of overweight and obesity on blood pressure control among hypertensive patients (results from nature Htn). *J Hypertens*. 2021;39;Suppl 1:339.
DOI:10.1097/01.hjh.0000748444.35042.87
2. Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity, kidney dysfunction and hypertension: mechanistic links. *Nat Rev Nephrol*. 2019;15(6):367-85.
DOI: 10.1038/s41581-019-0145-4, PMID 31015582, PMCID PMC7278043.

3. Cohen JB, Gadde KM. Weight loss medications in the treatment of obesity and hypertension. *Curr Hypertens Rep.* 2019; 21(2):16.
DOI: 10.1007/s11906-019-0915-1, PMID 30747357, PMCID PMC6415530.
4. Agness A, Ibrahim A. Should obesity be recognized as a disease? *Br Med J.* 2019;366:l4258.
5. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, et al. Global disparities of hypertension prevalence and control: A systematic analysis of population-based studies from 90 countries. *Circulation.* 2016;134(6):441-50.
DOI: 10.1161/CIRCULATIONAHA.115.018912, PMID 27502908.
6. Anto EO, Owiredu WKBA, Adua E, Obirikorang C, Fondjo LA, Annani-Akollor ME, et al. Prevalence and lifestyle-related risk factors of obesity and unrecognized hypertension among bus drivers in Ghana. *Heliyon.* 2020;6(1):e03147.
DOI: 10.1016/j.heliyon.2019.e03147, PMID 32042945.
7. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol.* 2020;16(4):223-37.
DOI: 10.1038/s41581-019-0244-2, PMID 32024986, PMCID PMC7998524.
8. Cochran JM, Siebert VR, Bates J, Butulija D, Kolpakchi A, Kadiyala H, et al. The relationship between adult height and blood pressure. *Cardiology.* 2021;146(3):345-50.
Doi: 10.1159/000514205, PMID 33721862.
9. Islam MT, Siraj MS, Hassan MZ, Nayem M, Chandra Nag D, Islam MA, et al. Influence of height on blood pressure and hypertension among Bangladeshi adults. *Int J Cardiol Hypertens.* 2020;5:100028.
DOI: 10.1016/j.ijch.2020.100028, PMID 33447757, PMCID PMC7803027.
10. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol.* 2020;16(4):223-37.
DOI: 10.1038/s41581-019-0244-2, PMID 32024986, PMCID PMC7998524.
11. Grassi G. How to treat hypertension in the obese. *European Society of Cardiology.* 2013;12:2-18.
12. Oghomen Onwaha. Prevalence and correlates of obesity in Irrua, Edo State, Nigeria, national postgraduate medical college of Nigeria (NPMCN), article section; 2019.
13. Chan RS, Woo J. Prevention of overweight and obesity: How effective is the current public health approach. *Int J Environ Res Public Health.* 2010;7(3):765-83.
DOI: 10.3390/ijerph7030765, PMID 20617002, PMCID PMC2872299.
14. Cameron NA, Petito LC, McCabe M, Allen NB, O'Brien MJ, Carnethon MR, et al. Quantifying the sex-race/ethnicity-specific burden of obesity on incident diabetes mellitus in the United States, 2001 to 2016: MESA and NHANES. *J Am Heart Assoc.* 2021;10(4):e018799.
DOI: 10.1161/JAHA.120.018799, PMID 33563002.
15. Lee HA, Park H. Comorbidity network analysis related to obesity in middle-aged and older adults: Findings from Korean population-based survey data. *Epidemiol Health.* 2021;43:e2021018.
DOI: 10.4178/epih.e2021018, PMID 33677857.
16. UNICEF. Global nutrition report. Sourced on the 25th March, 2023; 2018.
Available:<https://www.unicef.org/press-releases/2018-global-nutrition-report-reveals-malnutrition-unacceptably-high-and-affects>
17. World Health Organization. Key facts Obesity and overweight; 2021. Sourced on 29th March, 2023.
Available:<https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
18. Jiang SZ, Lu W, Zong XF, Ruan HY, Liu Y. Obesity and hypertension. *Exp Ther Med.* 2016;12(4):2395-9.
DOI: 10.3892/etm.2016.3667, PMID 27703502, PMCID PMC5038894.
19. Zhang Y, Hou LS, Tang WW, Xu F, Xu RH, Liu X, et al. High prevalence of obesity-related hypertension among adults aged 40 to 79 years in Southwest China. *Sci Rep.* 2019;9(1):15838.
DOI: 10.1038/s41598-019-52132-6, PMID 31676873.
Zhang Y, Hou LS, Tang WW, Xu F, Xu RH, Liu X, et al. High prevalence of obesity-related hypertension among adults aged 40 to 79 years in Southwest China. *Sci Rep.* 2019;9(1):15838.

- DOI: 10.1038/s41598-019-52132-6, PMID 31676873.
20. Whelton PK, Carey RM, Aronow WS, et al. ACC/AHA/AAPA/ABC/ACPM/AGS. Hypertension; 71(6):13 - 115/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/ American Heart Association Task Force on Clinical Practice Guidelines; 2017.
 21. Clark CE, McDonagh STJ, McManus RJ, Martin U. COVID-19 and hypertension: Risks and management. A scientific statement on behalf of the British and Irish Hypertension Society. J Hum Hypertens. 2021;35(4):304-7.
DOI: 10.1038/s41371-020-00451-x, PMID 33483621.
 22. Ryan Richard M, Netta Weinstein J, Bernstein J, Brown KW, Mistretta L, Gagné M, Kirk Warren Brown, Louis Mistretta and Marylene Gagne. Vitalizing effects of being outdoors and in nature. J Environ Psychol. 2010;30:159-68.
DOI: 10.1016/j.jenvp.2009.10.009
 23. Shariq OA, McKenzie TJ. Obesity-related hypertension: A review of pathophysiology, management, and the role of metabolic surgery. Gland Surg. 2020;9(1):80-93.
DOI: 10.21037/gs.2019.12.03, PMID 32206601.
 24. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. Nat Rev Nephrol. 2020;16(4):223-37.
DOI: 10.1038/s41581-019-0244-2, PMID 32024986, PMCID PMC7998524.
 25. World Health Organization. Cardiovascular diseases (CVDs): Key facts; 2021. Sourced on the 16 April, 2023.
Available: [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)).
 26. Swaminathan S, Dehghan M, Raj JM, Thomas T, Rangarajan S, Jenkins D, et al. Associations of cereal grains intake with cardiovascular disease and mortality across 21 countries in Prospective Urban and Rural Epidemiology study: Prospective cohort study. Br Med J. 2021;372:m4948.
DOI: 10.1136/bmj.m4948, PMID 33536317.
 27. Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity-induced hypertension: Interaction of neurohumoral and renal mechanisms. Circ Res. 2015;116(6):991-1006.
DOI: 10.1161/CIRCRESAHA.116.305697, PMID 25767285.
 28. Onyeaka H, Nwaiwu O, Oibileke KC, Miri T, Al-Sharif ZT. Global nutritional challenges of reformulated food: a review. Food Science & Nutrition. 2023. doi: 10.1002/fsn3.3286.
 29. Akpa OM, Made F, Ojo A, Ovbiagele B, Adu D, Motala AA, et al. Regional patterns and association between obesity and hypertension in Africa: Evidence From the H3Africa CHAIR Study. Hypertension. 2020;75(5):1167-78.
DOI: 10.1161/HYPERTENSIONAHA.119.14147, PMID 32172619.
 30. Akpa OM, Made F, Ojo A, Ovbiagele B, Adu D, Motala AA, et al. Regional patterns and association between obesity and hypertension in Africa: Evidence from the H3Africa CHAIR study. Hypertension. 2020;75(5):1167-78.
DOI: 10.1161/HYPERTENSIONAHA.119.14147, PMID 32172619.
 31. Hendriks ME, Wit FWNM, Roos MTL, Brewster LM, Akande TM, de Beer IH, et al. Hypertension in sub-Saharan Africa: cross-sectional surveys in four rural and urban communities. PLoS One. 2012;7(3):e32638.
DOI: 10.1371/journal.pone.0032638, PMID 22427857.
 32. Ajayi IO, Sowemimo IO, Akpa Onoja M, Ossai NE. Prevalence of hypertension and associated factors among residents of Ibadan-North Local Government Area of Nigeria. Niger J Cardiol. 2016;13(1):67-75.
DOI: 10.4103/0189-7969.165168
 33. Princewale F, Cumber SN, Kimbi J, Anchang, Nkfusai CN, Keka EI, Viyoff VZ, et al. Fa la Bede, Joyce Mahlako Tsoka-Gwegweni, Eric Achidi akum. Pan Afr Med J. Prevalence and risk factors associated with hypertension among adults in a rural setting: The case of Ombe, Cameroon. 2019;34:147.
 34. Bosu WK, Bosu DK. Prevalence, awareness and control of hypertension in Ghana: A systematic review and meta-analysis. PLOS ONE. 2021;16(3):e0248137.
DOI: 10.1371/journal.pone.0248137, PMID 33667277.

35. Omar SM, Musa IR, Osman OE, Adam I. Prevalence and associated factors of hypertension among adults in Gadarif in eastern Sudan: a community-based study. *BMC Public Health*. 2020;20(1):291. DOI: 10.1186/s12889-020-8386-5, PMID 32138715.
36. Nawi AM, Mohammad Z, Jetly K, Razak MAA, Ramli NS, Ibadullah WAHannan W, et al. The prevalence and risk factors of hypertension among the urban population in Southeast Asian countries: A systematic review and meta-analysis. *Int J Hypertens*. 2021;14.
37. De Lorenzo A, Gratteri S, Gualtieri P, Cammarano A, Bertucci P, Di Renzo L. Why primary obesity is a disease? *J Transl Med*. 2019;17(1):169. DOI: 10.1186/s12967-019-1919-y, PMID 31118060.
38. Chukwuonye II, Chuku A, John C, Ohagwu KA, Imoh ME, Isa SE, et al. Prevalence of overweight and obesity in adult Nigerians – a systematic review, *Diabetes, Metabolic Syndrome and Obesity*. 2013;6:43-7.
39. Adeloye D, Ige-Elegbede JO, Ezejimofor M, Owolabi EO, Ezeigwe N, Omoyele C, et al. Estimating the prevalence of overweight and obesity in Nigeria in 2020: a systematic review and meta-analysis. *Ann Med*. 2021;53(1):495-507. DOI: 10.1080/07853890.2021.1897665, PMID 33783281.
40. Egbi OG, Ahmed SD, Madubuko R. Prevalence and biosocial determinants of hypertension in a rural population in Edo State, Southern Nigeria. *Afr J Prim Health Care Fam Med*. 2021;13(1):e1-7. DOI: 10.4102/phcfm.v13i1.2833, PMID 34212744, PMCID PMC8335784.
41. Adegoke O, Ozoh OB, Odeniyi IA, Bello BT, Akinkugbe AO, Ojo OO, et al. Prevalence of obesity and an interrogation of the correlation between anthropometric indices and blood pressures in urban Lagos, Nigeria. *Sci Rep*. 2021;11(1):3522. DOI: 10.1038/s41598-021-83055-w, PMID 33568712.
42. Oduwole AA, Ladapo TA, Fajolu IB, Ekure EN, Adeniyi OF. Obesity and elevated blood pressure among adolescents in Lagos, Nigeria: a cross-sectional study. *BMC Public Health*. 2012;12:616. DOI: 10.1186/1471-2458-12-616, PMID 22867531.
43. Malaza A, Mossong J, Bärnighausen T, Newell ML. Hypertension and obesity in adults living in a high HIV prevalence rural Area in South Africa. *PLoS One*. 2012;7(10):e47761. DOI: 10.1371/journal.pone.0047761, PMID 23082211.
44. Adeloye D, Ige-Elegbede JO, Ezejimofor M, Owolabi EO, Ezeigwe N, Omoyele C, et al. Estimating the prevalence of overweight and obesity in Nigeria in 2020: A systematic review and meta-analysis. *Ann Med*. 2021;53(1):495-507. DOI: 10.1080/07853890.2021.1897665, PMID 33783281.
45. Ajayi IO, Sowemimo IO, Akpa OnojaM, Ossai NE. Prevalence of hypertension and associated factors among residents of Ibadan-North Local Government Area of Nigeria. *Niger J Cardiol*. 2021;13(1): 67-75. PMID 21105455.
46. Gaspar L, Poliak P, Makovník M, Hlinšťáková S, Krahulec B, Pisková T, et al. Obezita a artérová hypertenzia [Obesity and arterial hypertension]. *Vnitr Lek*. 2010;56(10):1074-7. PMID 21105455.
47. Miah L, Strafford H, Fonferko-Shadrach B, Hollinghurst J, Sawhney IMS, Hadjikutis S, et al. Incidence, prevalence, and health care outcomes in idiopathic intracranial hypertension A population study. 2021;96(8):1251-61.
48. Abolhasani M, Maghbouli N, Karbalai Saleh S, Aghsaeifar Z, Sazgara F, Tahmasebi M, et al. Which anthropometric and metabolic index is superior in hypertension prediction among overweight/obese adults? *Integr Blood Press Control*. 2021;14:153-61. DOI: 10.2147/IBPC.S340664, PMID 34795521, PMCID PMC8593692.
49. Ogah OS, Madukwe OO, Onyeonoro UU, Chukwuonye II, Ukegbu AU, Akhimien MO, et al. Cardiovascular risk factors and non-communicable diseases in Abia state, Nigeria: report of a community-based survey. *Int J Med Biomed Res*. 2013;2(1):57-68. DOI: 10.14194/ijmbr.2110
50. Yusuf Shehu, Muhammad M, Baba M, Ibrahim G, Andrew U. Metabolic syndrome overweight and obesity among adolescents in Kano, Nigeria. *J Metab Syndr*. 2013;2(21).

51. Okokhere PO, Bankole IA, Erohubie CA. Characteristics, risk factors and case fatality rate of stroke in hospitalized patients in semi-urban South–South Nigeria. *SAGE Open Med.* 2013;1: 2050312113516112. DOI: 10.1177/2050312113516112, PMID 26770696.

© 2023 Babangida et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98864>