

2950 Niles Road, St. Joseph, MI 49085-9659, USA 269.429.0300 fax 269.429.3852 hq@asabe.org www.asabe.org

An ASABE Meeting Presentation DOI: 10.13031/aim.20162460831 Paper Number: 162460831

An investigation into the fertilizer potential of slaughterhouse cattle paunch

Bernadette K. McCabe¹, Diogenes L. Antille^{1,*}, Henry W. G. Birt¹, Jennifer E. Spence¹, Jamal M. Fernana¹, Wilmer van der Spek², Craig P. Baillie¹

¹ University of Southern Queensland, National Centre for Engineering in Agriculture, Toowoomba, QLD, Australia.

²Wageningen University and Research Centre, Department of Soil Physics and Land Management, Wageningen, The Netherlands.

*Corresponding author: E: Dio.Antille@usq.edu.au, Ph: +61 7 46312948.

Written for presentation at the 2016 ASABE Annual International Meeting Sponsored by ASABE Orlando, Florida July 17-20, 2016

Abstract. In Australia, the red meat processing industry actively seeks approaches to improve the management of solid waste from processing operations and enhance the environmental performance. Recycling of paunch waste to farmland could be a cost-effective and practicable environmental option. However, little is known about the agronomic value of fresh and composted paunch, and the associated requirements for land application. Therefore, a short-term experimental work was undertaken to assess potential risks due to weed seed contamination and determine the agronomic response of ryegrass (Lolium perenne L.) to soil incorporation of paunch. The risk of weed contamination from soil application of paunch appeared to be low; however, methods that account for viability of seeds may be required to fully discard such a risk. Soil application of paunch at field equivalent rates of 150-300 kg ha⁻¹ of N increased dry matter yield by ≈30% on average compared with untreated grass, but was approximately 35% lower than a mineral fertilizer treatment applied at the same rates. Dry matter yield of paunch-treated grass was between 2000 and 3000 kg per ha over four consecutive cuts at 25-day intervals. Nitrogen use-efficiency of paunch was approximately 10% (range: 3% to 20%, depending on paunch type), and total N in harvested plant material showed values, which were between 2% and 3%. Overall, there appears to be potential for paunch-derived products to be used as a source of carbon and nutrients in crop production. Areas that merit a research priority within this space are also outlined in this paper. Such work is required to inform soil-, climate- and crop-specific land application rates, optimize agronomic performance, and minimize environmental concerns. There is also a requirement for the value proposition to industry to be determined, including reduced cost of disposal of material via gate fees and fertilizer replacement value.

Keywords. Byproducts, Dry matter yield, Compost, Fertilizer replacement value, Nutrient recovery, Nutrient use-efficiency, Recycling of organic waste, Vertisol.

The authors are solely responsible for the content of this meeting presentation. The presentation does not necessarily reflect the official position of the American Society of Agricultural and Biological Engineers (ASABE), and its printing and distribution does not constitute an endorsement of views which may be expressed. Meeting presentations are not subject to the formal peer review process by ASABE editorial committees; therefore, they are not to be presented as refereed publications. Citation: McCabe, B. K., Antille, D. L., Birt, H. W. G., Spence, J. E., Fernana, J. M., van der Spek, W., Baillie, C. P. (2016). An investigation into the fertilizer potential of slaughterhouse cattle paunch. ASABE Paper No.: 16-2460831. St. Joseph, Mich.: ASABE. DOI:10.13031/aim.202460831. For information about securing permission to reprint or reproduce a meeting presentation, please contact ASABE at rutter@asabe.org or 269-932-7004 (2950 Niles Road, St. Joseph, MI 49085-9659 USA).

Introduction

Australia produced approximately 2.6 million tonnes of red meat in 2015 (≈5% of global production) and is the seventh largest beef producer in the world (FAO, 2015). This production results in a significant amount of waste, which requires careful management and disposal to minimize environmental problems, social inconvenience and economic cost (Jayathilakan, et al., 2012; Petrovic, et al., 2015). A large proportion of the waste generated by the red meat industry is produced during slaughter (Arvanitoyannis and Ladas, 2008). Paunch waste is the partially digested feed from the rumen of cattle that is removed during the processing of carcasses, and is a significant contributor to the waste stream of Australian abattoirs (Australian Bureau of Statistics, 2016). For example, a medium-sized abattoir in Australia produces between 60 and 90 m³ of paunch per week, which equates to about 800 tonnes of paunch (dry matter basis) per year (Spence, 2012). Current disposal methods incur significant costs to abattoirs and are increasingly regarded as non-environmentally friendly options. By contrast, land application of paunch is relatively less expensive compared with traditional disposal methods, and similar to other organic materials, is considered to be the best practicable environmental option in most circumstances, as it supports the waste management hierarchy (Liu and Haynes, 2013; Ksheem et al., 2015; Six et al., 2016). Abattoirs currently undertake one of several options to manage paunch waste, including: (a) removal of paunch and other solids off-site, (b) composting material on-site and use on-site, and (c) composting material on-site and use off-site. For house-locked abattoirs, where land space is not available, composting or processing of paunch cannot be performed on-site. In these situations, paunch is removed off-site to a licensed premise, which incurs significant costs in gate fees and transport. Paunch is subsequently mixed with other bulking agents and sold as either bulk soil conditioner or bagged compost. Consequently, the benefits of the nutrient-rich material are lost to the farming community because the end use is primarily for domestic gardens.

At present, there are no guidelines or industry standards available regarding best management practice of paunch, which poses the challenge of how to recycle this material for beneficial use on farms at an affordable cost to the industry. Abattoirs in southern Queensland (Australia) have proposed one option, which includes firstly holding the paunch and other yard solids on-site for 1-2 weeks, and subsequently transporting the material to a suitable property for up to 3 months before it is used as a soil amendment. Beneficial Use Agreements (BUA) have been developed; however, further work to qualify particulars surrounding the requirements of paunch management and handling is required as well as to determine whether there is sufficient information to satisfy the conditions of BUA for handling of paunch. Furthermore, there is a need for fundamental work to be undertaken on the following key issues: (a) validating the criteria for paunch stabilization, and (b) determine application rates for on-farm use. There appears to be a paucity of information available in the scientific literature on the beneficial use of paunch as a fertilizer material (e.g., Hansen, 1992). Little is also known about the agronomic efficiency of paunch applied to soil in various degrees of decomposition, from fresh through to fully composted (Wilson, 1992; Fleming and MacAlpine, 2004).

This study aims to determine optimal agronomic and environmentally-safe land application rates to assist the development of best management practices for the use of paunch as a fertilizer in crop production. Such work is required to maximize nutrient recovery in crop biomass and minimize environmental concerns. This preliminary work forms the basis of a broader scope of research which will establish criteria for measuring the stability of paunch, assess potential risks from on-site storage and on- or off-site farm use due to the risk of pathogens and weed seeds contamination and will quantify greenhouse gas emissions (N₂O, CH₄, CO₂) following soil incorporation. The work reported in this paper summarizes preliminary findings derived from a short-term experimental study, which was conducted to: (1) assess potential risks due to weed seed contamination, and (2) determine the agronomic response of ryegrass to soil incorporation of paunch. Two sets of experiments were established under controlled environmental conditions as discussed here, and included laboratory and glasshouse studies, respectively.

Materials and Methods

Two experiments were conducted under controlled conditions of temperature and soil moisture in the laboratory and glasshouse, which included a germination test and a pot trial. A characterization of the soil used in the pot experiment (Table 1) and the range of paunch materials (Table 2) used in the study were also conducted to assist the interpretation of soil nutrient dynamics and soil × plant interactions. These materials included fresh and composted paunch at various degrees of stabilization. Composted paunch is referred to in the text based on its composting age, which is given in weeks (from 2 to 16, respectively). The soil used in the pot experiment was a Black Vertosol, which is common in southern Queensland (Australia) (Isbell, 2002), and was therefore representative of the area of interest for this work where paunch is likely to be applied.

 Table 1: Characterisation of the Black Vertosol used in the glasshouse study. Description of analyses conducted prior to the experiment to establish baseline levels. SD is standard deviation, n=3.

Determination	Value ± SD	Unit	Method	
Sand (>20 μm)	9.3±0.58	% (w w ⁻¹)		
Silt (2-20 μm)	20.1±2.00	% (w w ⁻¹)	Bouyoucos (1962)	
Clay (<2 μm)	70.6±2.08	% (w w ⁻¹)		
Field capacity	40.4±3.11	% (w w⁻¹) at ⅓ bar	Cassel and Nielsen (1986)	
Soil bulk density	1040±85	kg m ⁻³	Blake and Hartge (1986)	
Soil pH	8.55±0.071		Rayment and Lyons (2011)	
Electrical conductivity	0.121±0.008	dS m ⁻¹	Rayment and Lyons (2011)	
Soil organic C	1.95±0.07	% (w w ⁻¹)	Walkley and Black (1934)	
Total N in soil	0.15±0.01	% (w w-1)	MAFF (1986, Method No.: 49)	
Soil mineral N	2.4±0.65	mg kg ⁻¹	MAFF (1986, Method No.: 53)	
Soil extractable P	20.3±3.89	mg kg ⁻¹	Colwell (1963)	

Table 2: Characterisation of paunch and mineral fertilizers used in the glasshouse study. SD is standard deviation (n=3, except when not shown). Determination of total N and total P are based on MAFF (1986, Method No.: 49), and BS7755 (1998), respectively. SSP is single superphosphate, and number of weeks refers to composting time.

······································									
Determination	Total N	Total P	Moisture content	Bulk Density (kg m ⁻³)					
Material	(%)	(%)	(%, w w ⁻¹)	Untapped	Tapped				
2 weeks	1.09±0.07	0.11±0.01	95.6	594±15.1	780±15.6				
4 weeks	0.72±0.00	0.11±0.02	81.6	350±20.6	593±37.9				
6 weeks	1.41±0.03	0.36±0.07	66.6	379±12.7	530±10.1				
12 weeks	1.89±0.02	0.21±0.01	76.6	176±4.9	249±7.2				
16 weeks	0.96±0.03	0.28±0.02	52.9	407±0.9	490±25.9				
Urea	46			743±6.3**					
SSP		18 [*]		1120					

*Total P as P₂O₅ (DEFRA, 2010), ** after Antille et al. (2015a).

Germination Test

The germination test was conducted to assess the risk of weed contamination and the likely introduction of weed species following soil incorporation of paunch. The five types of paunch described in Table 2 (from fresh through to fully composted) were mixed with pure sand using a ratio of 100 g (moist-basis) of organic material-to-400 g of oven-dried sand, and placed in plastic containers (dimensions: 200×100×50 mm). Water was subsequently added to the mixture to reach near-saturated conditions (≈90% of saturation) and was maintained throughout the experiment for a period of three weeks. All treatments including controls (pure sand with no addition of paunch) were replicated five times and plantule counts recorded daily. Figure 1 shows the experimental setup. Paunch pH and electrical conductivity were measured based on MAFF (1986, Methods No.: 24 and 33, respectively) to determine potential effects of these two factors on seed germination.



Figure 1: Overview of the germination test, (a) paunch treatments from 2 to 16 week-old composts, and (b) close-up of a germination tray containing fresh paunch (2 week-old compost).

Glasshouse Study

The glasshouse study was conducted to determine the agronomic efficiency of paunch applied to a ryegrass (*Lolium perenne* L.) grown in pots under controlled environmental conditions of temperature and soil moisture. The agronomic efficiency of the organic materials was assessed by determining: (1) aboveground biomass referred to in the text as field equivalent dry matter yield (DMY) per ha, (2) nitrogen uptake, and (3) nitrogen use-efficiency (NUE). This assessment was conducted by comparing five types of paunch with mineral fertilizers of known performance, and controls (zero-amendment), respectively. Controls were used to quantify the soil's residual nutrient contribution to biomass yield and nitrogen uptake, and to enable NUE from fertilizer- and paunch-treated grass to be estimated using the difference method (Baligar et al., 2001). The glasshouse experiment will inform the design of future full-scale field experimentation and the development of preliminary guidelines for land application rates.

The experiment used the same materials described for the germination test, which were sourced from local abattoirs in southern Queensland (Australia (Table 1). The experiment included treatment based on urea (46% N) and single superphosphate (0:18:0) to evaluate N application rates. Field equivalent application rates included control (zero-amendment), 150 and 300 kg ha⁻¹ of N, respectively. All treatments, including controls were replicated three times (n=3). Preparation of pots and mixing of fertilizer materials with soil were based on the approach used by Antille et al. (2014a). Dry matter yield (DMY) was determined by hand-cutting the grass to a height of 40 mm, at regular intervals of 25 days after emergence, which was recorded at day 8 after seeding. The time interval for cutting of the grass was consistent with recommendations reported in earlier studies for maximum yield (e.g., Fulkerson et al., 1993; Antille et al., 2015b). At the time this report was produced, a total of four cuts were completed, however the experiment will continue to include a total of six cuts. Total fresh weight was determined and a subsample was taken for determination of dry matter, and total N content in plant material, which was determined based on MAFF (1986, Method No.: 48). From this, nitrogen uptake was derived and used to determine NUE, as follows (Equation 1, after Baligar et al., 2001):

$$NUE = \left(\frac{N_F - N_{F=0}}{N_{Rate}}\right) \tag{1}$$

Where: *NUE* is nitrogen use-efficiency (kg kg⁻¹), N_F and $N_{F=0}$ are N uptakes (kg ha⁻¹) corresponding to fertilizer or organic amendment treatments and control (zero-amendment), respectively, and N_{Rate} is field equivalent N application rate (kg ha⁻¹). Data for N content in plant material was available for the first three cuts only. Pots were placed in a glasshouse facility available at USQ and maintained near-field capacity conditions at 25±3°C throughout the experiment. The experimental setup in the glasshouse is shown in Figure 3.



Figure 3: Overview of the pots experiment conducted in a glasshouse under controlled conditions of temperature and soil moisture, (a) preparation of pots, (b) close-up of a pot (note germination of plants), and (c) established grass in pots eight days after germination.

Fertilizer Replacement Value of Paunch

Nitrogen fertilizer replacement value (NFRP) of paunch was estimated based on the approach reported in Lalor et al. (2011) for cattle slurry, which uses the dry matter yield-to-fertilizer nitrogen response curve to derive NFRV.

Statistical Analyses

Statistical analyses were undertaken using GenStat release 16th Edition (VSN, 2013) and involved analysis of variance (ANOVA), and the least significant differences (LSD) to compare means using a probability level of 5%. Repeated measurement of ANOVA was employed to compare measured crop attributes between-cuts, using the same probability level. Dry matter yield-to-nitrogen responses were investigated by means of simple (linear) regression analyses. Quadratic functions were also fitted to the data and results are discussed. The analyses conducted were graphically verified by means of residual plots. Normalization of the data was not required.

Results and Discussion

Germination Test

The main results derived from this test are summarized below:

- The number of plantules germinated after three weeks reported a value of zero in all daily observations and treatments. This suggested low (or no) risk of weed contamination in soil amended with paunch whether fresh or at varying degrees of composting,
- However, given weed species often exhibit dormancy (temporary absence of germination capacity) even under satisfactory conditions for germination (Vivian et al., 2008), the risk of weed contamination in soils receiving paunch may not be completely discarded. Therefore, application of alternative methods (e.g., Grabe, 1970; Tekrony, 1983; Tompkins et al., 1998) to determine risk of weed contamination warrants further investigation,
- Given values of paunch pH (range: 7.18 to 7.87) and EC (range: 0.40 to 0.69 dS m⁻¹) encountered in paunch samples, there was not sufficient evidence to suggest that weed seed germination was inhibited by these factors. No seeds germination may be attributed to the temperature range developed within the composting paunch windrow (≈50-60°C), which may have reduced seeds' viability as mentioned in related studies (Eghball and Lesoing, 2000).

Glasshouse Study

Dry matter yield (DMY) recorded over four cuts is shown in Figure 4. Overall, there were significant differences in DMY between treated and non-treated grass (P<0.001). There were also significant differences between amendment type, nitrogen (N) application rate, and interaction amendment × N application rate on DMY (P-values <0.05). The amendment type effect was mainly due to differences in DMY between urea and all types of paunch, particularly at cuts 1 and 2.

However, these differences were relatively smaller at cuts 3 and 4, and non-significant except for higher (≈10%)

DMY in grass treated with the 6 week-old material compared with other treatments (LSD 5% level: 95.7). Dry matter yield recorded in treated grass at cuts 3 and 4 was marginally higher but not statistically different than controls (LSD 5% level: 117.2), which suggested low mineral N supply rates from the organic materials. Despite this, less stabilization of organic matter in fresh (<2 week-old) and semi-composted (6 week-old) materials suggested that both C and N fractions were more readily available, and therefore N supply to the plant was affected to lesser extent compared with fully composted paunch. Figure 4 also shows that overall differences in DMY between-paunch treatments applied at field equivalent rates of 150 and 300 kg ha⁻¹ of N were less than 10% on average, which supports the above statement and denotes little agronomic benefit of paunch applied at the higher rate used in this study. For urea, there was a 22% difference in cumulative DMY over four cuts when applied at 150 and 300 kg ha⁻¹ of N, respectively.



Figure 4: Dry matter yield recorded over four consecutive cuts at 25-day intervals, and after a single application of mineral fertilizers and paunch to ryegrass. Number of weeks denotes compost age, followed by field equivalent N application rate in kg per ha. SD is standard deviation, SSP is single superphosphate.

Table 3 shows dry matter yield-to-nitrogen responses observed after a single application of paunch or mineral fertilizer, and over four consecutive cuts. Responses were essentially linear for the range of N application rates investigated (P-values <0.05), which agrees with earlier studies dealing with synthetic N sources (e.g., Reid, 1985; Antille et al., 2015b), and organic materials used in ryegrass (e.g., Antille et al., 2014a). Despite this, nonlinear responses were also fitted to the data, but estimate of parameters for the square term were not significant (p-values >0.05), which was observed for all amendments used in this study. Therefore, dry matter yield-to-nitrogen responses were better explained by linear functions, which showed significance to the linear term (P-values <0.05). This was expected because of the range of N application rates investigated; however, nonlinear functions may be possible with a more complete dataset (e.g., Sparrow, 1979; Morrison et al., 1980; Antille et al., 2013a-b). Linear regression analyses conducted for each paunch type explained, individually, relatively more variation than it did a common slope (P<0.05, R² =0.56) because of the amendment × N rate effect (P<0.05), with the exception of the moderately-composted material (12 week-old), which was sourced from a different abattoir ($R^2 = 0.40$). This observation suggested significant differences in the quality of the paunch material, linked to the composting process (time and pile management), and the type of cattle feed (e.g., grainor grass-fed or both). The slopes derived from the set of linear functions presented in Table 3 denote the agronomic efficiency of N applied as fertilizer (kg DMY kg⁻¹ N). Responses were higher with 6 week- and 16 week-old composts (≈3.1 kg kg⁻¹) compared with other materials, but about 50% lower than that obtained with urea (≈6.5 kg kg⁻¹). Nitrogen responses encountered in this study (range: 1.0-6.5 kg DMY kg⁻¹ N) were generally lower than the range (10-30 kg DMY kg⁻¹ N) reported in the literature for synthetic N fertilizers applied to ryegrass grown in subtropical environments (e.g., McKenzie, 1996; Callow et al., 2003). However, it is acknowledged the fact that our study comprised only four cuts over a period of 100 days, and that it was conducted during the autumn without accounting for seasonal patterns of grass growth (Fulkerson et al., 1998; Cullen et al., 2008; Vogeler et al., 2016).

Differences in responses observed between mineral fertilizer and organic amendment treatments are also

explained by $N \times P$ effect on DMY (Mouat and Nes, 1983; Fageria, 2001). As shown in Figure 5, relatively higher availability of single superphosphate-P (Charleston, 1984) compared with paunch-P is mentioned as a contributing factor to enhanced N uptake and DM partitioning of urea-N compared with the organic material.

Material	Response	P-value	R ²
2 weeks	DMY = 2183 + 1.78N	<0.05	0.51
4 weeks	DMY = 2131 + 2.02N	<0.05	0.70
6 weeks	DMY = 2200 + 3.03N	<0.05	0.74
12 weeks	DMY = 2127 + 1.02N	<0.10	0.40
16 weeks	DMY = 2147 + 3.18N	<0.05	0.86
Urea + single superphosphate	DMY = 2152 + 6.45N	<0.001	0.93

Table 3: Dry matter yield-to-nitrogen responses recorded over four cuts after a single application of mineral fertilizer and paunch to ryegrass. Number of weeks denotes compost age.

Nitrogen in plant material, uptake and use-efficiency

Figure 5 shows distribution of data corresponding to N in plant material recorded over the first three cuts for control, mineral fertilizer- and paunch-treated grass, respectively. Note the median value of grass treated with the 6-week old compost was approximately equivalent to that of the mineral fertilizer treatment ($\approx 2.6\%$ N). For the mineral fertilizer treatment, a relatively wide range of values (Q₁ = 2.1% N, Q₃ = 3.6% N) compared to all types of paunch (Q₁: 2.3-2.5% N, Q₃: 2.4-2.7% N) reflects significant effect of nitrogen application rate on total N in plant material (P<0.05), which was not observed in paunch-treated grass (P>0.05). Low N supply to grass crops compromises growth rate, tiller density and therefore biomass production, and may also reduce N concentration in plant (Wilman et al., 1976; Delagarde et al., 1997). For example, high-producing dairy cows require N contents in grass in the range of 2.2% to 2.7% to sustain satisfactory production levels and milk quality (Delaby et al., 1996; Peyraud and Astigarraga, 1998). Values of total N in plant material encountered within this study for both paunch and mineral fertilizers were within the range reported in the literature for ryegrass fertilized with synthetic N sources (e.g., Wilman and Mohamed, 1980; Lowe et al., 1999). For paunch-treated grass values within that critical range of N contents may be explained by relatively low biomass production (no dilution effect), and the frequency of cutting (Aavola and Kärner, 2008; Reyes et al., 2015).



Figure 5: A box-plot comparing total nitrogen content in plant material over three consecutive cuts at 25-day intervals, and after a single application of mineral fertilizers and paunch to ryegrass. Box-plots show: Min, Q₁, Med, Q₃, and Max, respectively. Use n=9 (control), n=18 (treatments), number of weeks denotes compost age. SSP is single superphosphate.

Figure 6 shows nitrogen uptake by grass over three consecutive cuts. Overall, N uptake was significantly higher (P<0.05) in paunch and fertilizer-treated grass compared with controls (by $\approx 60\%$ on average over the first three cuts). There were also significant amendment type and nitrogen application rate effects on nitrogen uptake, which were observed in all cuts (P-values <0.05). Despite this, the N application rate effect was mainly due to differences (≈35%) in N uptake in the mineral fertilizer treatment applied at 150 and 300 kg per ha of N, respectively. Overall differences in cumulative N uptake between-treatments were in the order: mineral fertilizer >6 weeks = 16 weeks >2 weeks = 4 weeks >12 weeks, respectively. However, N uptake in cut 3 reduced by approximately 50% across all treatments compared with the first two cuts, and approximated N uptake levels observed in controls (≈13 kg ha⁻¹ of N). Higher uptake in 16 week-old compared with 12 week-old composts may be attributed to qualitative differences in carbon (C) fractions (labile vs. non-labile) of the organic material affecting nutrient release, and therefore plant uptake (Fontaine et al., 2003; Culman et al., 2013; Antille et al., 2014b). These differences in compost-C may be due to the actual composting process (note that the 12 and 16 week-old composts were sourced from different abattoirs) as well as the quality of the animal feed influencing C dynamics during the stabilization process. These observations require further investigation including qualitative assessment of C fractions in composted paunch (e.g., Schnitzer, 1982; Garcia et al., 1991; Lucas and Weil, 2012).



Figure 6: Nitrogen uptake recorded over three consecutive cuts at 25-day intervals, and after a single application of mineral fertilizers and paunch to ryegrass. Number of weeks denotes compost age, followed by N application rate in kg per ha (field equivalent rate). SSP is single superphosphate.

Figure 7 shows nitrogen use efficiency (NUE, Equation 1) derived from cumulative N uptake over the first three cuts. Overall, there were significant effects of treatments and N application rates on NUE. On average, NUE decreased in the order: mineral fertilizer >6 weeks \approx 16 weeks >2 weeks \approx 4 weeks >12 weeks, respectively. Values of NUE relative to the mineral fertilizer treatment were approximately 40% in the 6 and 16 week-old composts, but reduced to 25% or less in all other paunch materials. On average across all type of amendments, NUE was approximately 50% lower when applied at 300 kg ha⁻¹ of N than at 150 kg ha⁻¹ of N. As shown in Figure 4, differences in DMY for paunch-treated grass at 150 and 300 kg ha⁻¹ of N were small. These results suggested that there was little agronomic benefit in applying paunch at field equivalent rates higher than 150 kg ha⁻¹ of N. and the same was true in terms of grass quality as N content in plant material varied only within a narrow range (Figure 5). From the range of paunch materials used in this study, it appears that highest agronomic efficiencies can be achieved with 6 and 16 week-old composts applied at 150 kg ha⁻¹ of N, respectively. A 6-week composting period has operational as well as financial advantages compared with 16 weeks. Such a practice may therefore be justified, which will also result in higher agronomic performance compared with either shorter or longer composting periods. However, the implications of using relatively less stabilized paunch materials on soil C stocks and soil C dynamics are not well understood, and cannot be inferred from this short-term study. This is an aspect that requires investigation, particularly in low C soils (e.g., SOC <1%).



Figure 7: Nitrogen use efficiency (NUE) recorded over three consecutive cuts at 25-day intervals, and after a single application of mineral fertilizers and paunch to ryegrass. Number of weeks denotes compost age, followed by N application rate in kg per ha (field equivalent). Superimposed figure shows NUE of paunch relative to NUE of mineral fertilizer treatment (averaged NUE over the two N application rates used in this study). SSP is single superphosphate.

Fertilizer Replacement Value of Paunch

Mean values (±SD) of N fertilizer replacement value (NFRV) of paunch applied to grass were: 39±19.8% (2 weeks), 35±15.5% (4 weeks), 56±19.3% (6 weeks), 19±17.5% (12 weeks), and 52±13.6% (16 weeks), respectively. Value range of NFRV of paunch within this study was generally higher than those reported in the literature for organic materials applied to grass under field conditions; for example, farm yard manure (range: 37-50%, Pikula et al., 2016), cattle slurry (range: 10-39%, Lalor et al., 2011), and composted household waste (<10%, Petersen, 2003). Field-scale studies are required to validate these preliminary results obtained under controlled conditions in the glasshouse, and determine how timing and method of application (e.g., soil incorporated or surface-applied) influence NFRV of paunch. Based on related studies (e.g., Misselbrook et al., 1996; Laws et al., 2002; Lalor et al., 2014) this is mentioned as an important practical consideration for optimizing nutrient recovery in crop and minimizing nitrogen losses following soil application of paunch.

Summary

The main findings derived from this work are summarized below:

- 1. Risk of weed contamination from soil application of paunch appears to be low. Methods that may enable accounting for dormancy of weed seeds, and viability of seeds may be required to fully discard such a risk,
- Soil application of paunch increased dry matter yield of ryegrass by ≈30% on average compared with untreated grass, but was approximately 35% lower than the mineral fertilizer treatment. Dry matter yieldto-nitrogen responses were linear within the N application rates investigated, which was observed for all amendments. Responses varied from 1.1 to 3.2 kg DM kg⁻¹ N for paunch- and about 6.5 kg DM kg⁻¹ N for urea-treated grass, respectively,
- Nitrogen use-efficiency (NUE) in paunch was approximately 10% on average (range: 3% to 20%), and between 0.15 and 0.40×NUE_{UREA} depending on paunch type and rate. The quality of harvested plant material in paunch-treated grass was satisfactory (total N_{PLANT} ranged between 2% and 3%) based on reported nutritional requirements for high-producing cattle,
- 4. The nitrogen fertilizer replacement value of paunch, relative to urea-N, reported an average value of 40±17.1% (single application over three consecutive cuts).

Future Research Requirements

The research areas highlighted here are required to inform soil-, climate- and crop-specific land application rates, optimize agronomic performance and minimize environmental concerns associated with the use of paunch in crop production:

- 1. Investigate the effect of paunch on soil carbon dynamics and carbon sequestration in soil, and associated effects on nutrient dynamics, particularly nitrogen, including modelling approaches that capture such effects on soil responses and crop productivity,
- Investigate the effect of paunch on greenhouse gas emissions, including surface application (likely grass based and arable cropping systems under zero-tillage) and shallow incorporation (conventional and minimum tillage cropping systems), respectively, and associated effects on seasonal patterns of nitrogen supply as affected by method of application,
- Determine the optimum timing of application for both grassland and arable cropping systems, and number of splits within a calendar year or growing season. There is also a requirement to review available methods for land application with a view to improving field operating efficiency, energy requirements of such operation, and minimize impacts on soil, particularly traffic compaction,
- 4. Conduct detailed economic analyses assessed against alternative options available for disposal of paunch. Such analyses should be undertaken as a cost reduction (avoidance) strategy, and may require the composted product to be partially subsidized to encourage uptake by farmers,
- 5. The nitrogen fertilizer replacement value of paunch has been derived from a short-term glasshouse experiment using pots, and therefore requires validation in longer-term trials under field conditions.

Acknowledgements

The work described in this paper was fully supported by the Australian Meat Processor Corporation (AMPC) and Meat and Livestock Australia (MLA). The authors are grateful to the National Centre for Engineering in Agriculture at the University of Southern Queensland (Toowoomba, QLD, Australia) for financial and operational support to conduct this research. Assistance received from Mr. Mike Spence, and USQ students Ms. Jamie Mcintyre and Ms. Seonmi Lee is gratefully acknowledged. Chemical analyses were conducted by Mr. Ian Grant (Agricultural Chemistry Pty Ltd., Ipswich, Australia).

References

- Aavola, R., Kärner, M. (2008). Nitrogen uptake at various fertilization levels and cutting frequencies of *Lolium* species. *Agronomy Research* 6(1): 5-14.
- Antille, D. L., Gallar, L., Miller, P. C. H., Godwin, R. J. (2015a). An investigation into the fertilizer particle dynamics off-the-disc. *Applied Engineering in Agriculture* 31(1): 49-60. DOI: 10.13031/aea.31.10729.
- Antille, D. L., Hoekstra, N. J., Lalor, S. T. J. (2015b). Field-scale evaluation of calcium ammonium nitrate, urea, and urea treated with N-(n-butyl) thiophosphoric triamide applied to grassland in Ireland. *Communications in Soil Science and Plant Analysis* 46(11): 1345-1361. DOI: 10.1080/00103624.2015.1033540.
- Antille, D. L., Sakrabani, R., Godwin, R. J. (2013a). Field-scale evaluation of biosolids-derived organomineral fertilisers applied to ryegrass (Lolium perenne L.) in England. *Applied and Environmental Soil Science* Vol.: 2013, Article ID: 960629, 9 pp. DOI: 10.1155/2013/960629.
- Antille, D. L., Sakrabani, R., Godwin, R. J. (2013b). Field studies on the value of organomineral fertilisers as amendments for perennial ryegrass (*Lolium perenne* L). ASABE Paper No.: 131620220. St. Joseph, Mich.: ASABE. DOI: 10.13031/aim.20131620220.
- Antille, D. L., Sakrabani, R., Godwin, R. J. (2014a). Effects of biosolids-derived organomineral fertilizers, urea and biosolids granules on crop and soil established with ryegrass (*Lolium perenne* L.). Communications in Soil Science and Plant Analysis 45(12): 1605-1621. DOI: 10.1080/00103624.2013.875205.
- Antille, D. L., Sakrabani, R., Godwin, R. J. (2014b). Nitrogen release characteristics from biosolids-derived organomineral fertilisers. *Communications in Soil Science and Plant Analysis, 45*(12): 1687-1698. DOI: 10.1080/00103624.2014.907915.
- Arvanitoyannis, I. S., Ladas, D. (2008). Meat waste management: treatment methods and potential uses of treated wastes. In: Waste Management for the Food Industries (1st Edition). Burlington, MA., USA: Elsevier Academic Press, 765-800 pp.
- Australian Bureau of Statistics. (2016). *Livestock and meat*. Canberra, ACT, Australia: Commonwealth of Australia. Retrieved from: http://www.abs.gov.au/ausstats/abs@.nsf/mf/7218.0.55.001.
- Baligar, V. C., Fageria, N. K., He, Z. L. (2001). Nutrient use efficiency in plants. Communications in Soil Science and Plant Analysis 32(7-8): 921-950. DOI: 10.1081/CSS-100104098.
- Blake, G. R., Hartge, K. H. (1986). Bulk density. In: Klute, A. (Editor). Methods of soil analysis. Part 1: Physical and mineralogical methods (2nd Edition). Agronomy Monograph No.: 9, pp. 363-382. Madison, WI: American Society of Agronomy.

British Standard 7755 Section 3.13. (1998). Soil quality. Chemical methods. Determination of cadmium, chromium, cobalt,

copper, lead, manganese, nickel and zinc in aqua regia extracts of soil. Flame and electrothermal atomic absorption spectrometric methods. Equivalent to ISO 11047:1998. London: The British Standards Institution.

- Callow, M. N., Lowe, K. F., Bowdler, T. M., Lowe, S. A., Gobius, N. R. (2003). Dry matter yield, forage quality and persistence of tall fescue (*Festuca arundinacea*) cultivars compared with perennial ryegrass (*Lolium perenne*) in a subtropical environment. *Australian Journal of Experimental Agriculture* 43(9): 1093-1099. DOI: 10.1071/EA02001.
- Cassel, D. K., Nielsen, D. R. (1986). Field capacity and available water capacity. *In*: Klute, A. (Editor). Methods of soil analysis, Part I: Physical and mineralogical methods (2nd Edition). Agronomy Monograph No.: 9, pp. 901-926. Madison, WI: American Society of Agronomy.
- Charleston, A. G. (1984). Solubilities of single superphosphate components in water, 2% citric acid, and neutral ammonium citrate solution. *New Zealand Journal of Science* 27(3): 269-277.
- Colwell, J. D. (1963). The estimation of the phosphorus fertilizer requirements of wheat in southern New South Wales by soil analysis. *Australian Journal of Experimental Agriculture* 3(10): 190-197. DOI: 10.1071/EA9630190.
- Cullen, B. R., Eckard, R. J., Callow, M. N., Johnson, I. R., Chapman, D. F., Rawnsley, R. P., Garcia, S. C., White, T., Snow, V. O. (2008). Simulating pasture growth rates in Australian and New Zealand grazing systems. *Australian Journal of Agricultural Research* 59(8): 761-768. DOI: 10.1071/AR07371.
- Culman, S. W., Snapp, S. S., Green, J. M., Gentry, L. E. (2013). Short- and long-term labile soil carbon and nitrogen dynamics reflect management and predict corn agronomic performance. *Agronomy Journal* 105(2): 493-502. DOI: 10.2134/agronj2012.0382.
- Delaby, L., Peyraud, J. L., Vérité, R., Marquis, B. (1996). Effect of protein content in the concentrate and level of nitrogen fertilization on the performance of dairy cows in pasture. *Animal Research* 45(4): 327-341. DOI: 10.1051/animres:19960405.
- Delagarde, R., Peyraud, J. L., Delaby, L. (1997). The effect of nitrogen fertilization level and protein supplementation on herbage intake, feeding behaviour and digestion in grazing dairy cows. *Animal Feed Science and Technology* 66(1-4): 165-180. DOI: 10.1016/S0377-8401(96)01098-X.
- Department for Environment, Food and Rural Affairs. (2010). *Fertiliser manual*. Reference Book No.: 209, (8th Edition). London: The Stationery Office.
- Eghball, B., Lesoing, G. W. (2000). Viability of weed seeds following manure windrow composting. *Compost Science and Utilization* 8(1): 46-53. DOI: 10.1080/1065657X.2000.10701749.
- Fageria, V. D. (2001). Nutrient interactions in crop plants. Journal of Plant Nutrition 24(8): 1269-1290. DOI: 10.1081/pln-100106981.
- Fleming, R., MacAlpine, M. (2004). Composting paunch manure with solid cattle manure. Final Report, 13 pp. Ridgetown, ON, Canada: Ridgetown College, University of Guelph.
- Fontaine, S., Mariotti, A., Abbadie, L. (2003). The priming effect of organic matter: a question of microbial competition? *Soil Biology and Biochemistry* 35(6): 837-843. DOI: 10.1016/s0038-0717(03)00123-8.
- Food and Agricultural Organization of the United Nations. (2015). *Production quantity of cattle and indegenous cattle.* Retrieved from: http://faostat3.fao.org/.
- Fulkerson, W. J., Slack, K., Hennessy, D. W., Hough, G. M. (1998). Nutrients in ryegrass (*Lolium spp.*), white clover (*Trifolium repens*) and kikuyu (*Pennisetum clandestinum*) pastures in relation to season and stage of regrowth in a subtropical environment. *Australian Journal of Experimental Agriculture* 38(3): 227-240. DOI: 10.1071/EA97161.
- Fulkerson, W. J., Slack, K., Moore, K., Rolfe, C. (1993). Management of *Lolium perenne/Trifolium repens* pastures in the subtropics. I. Effect of defoliation interval, seeding rate and application of N and lime. *Australian Journal of Agricultural Research* 44(8): 1947-1958. DOI: 10.1071/AR9931947.
- Garcia, C., Hernandez, T., Costa, F. (1991). Changes in carbon fractions during composting and maturation of organic wastes. *Environmental Management* 15(3): 433-439. DOI: 10.1007/bf02393889.
- Grabe, D. F. (1970). Tetrazolium Testing Handbook, pp. 62. Contribution No.: 29, Handbook on Seed Testing. Washington DC, USA: Association of Official Seed Analysts.
- Hansen, C. L. (1992). Land application of paunch manure and blood. In: Pearson, A. M., Dutson, T. R. (Editors). Inedible meat by-products, Vol.: 8, pp. 359-373. Dordrecht: Springer Netherlands. DOI: 10.1007/978-94-011-7933-1_15.
- Isbell, R. F. (2002). *The Australian Soil Classification*. Melbourne, Australia: CSIRO Publishing.
- Jayathilakan, K., Sultana, K., Radhakrishna, K., Bawa, A. (2012). Utilization of byproducts and waste materials from meat, poultry and fish processing industries: a review. *Journal of Food Science and Technology* 49(3): 278-293. DOI: 10.1007/s13197-011-0290-7.
- Ksheem, A. M., Bennett, J. McL., Antille, D. L., Raine, S. R. (2015). Towards a method for optimized extraction of soluble nutrients from fresh and composted chicken manures. *Waste Management* 45: 76-90. DOI:10.1016/j.wasman.2015.02.011.
- Lalor, S. T. J., Schröder, J. J., Lantinga, E. A., Oenema, O., Kirwan, L., Schulte, R. P. O. (2011). Nitrogen fertilizer replacement value of cattle slurry in grassland as affected by method and timing of application. *Journal of Environmental Quality* 40(2): 362-373. DOI: 10.2134/jeq2010.0038.
- Lalor, S. T. J., Schröder, J. J., Lantinga, E. A., Schulte, R. P. O. (2014). Effect of application timing and grass height on the nitrogen fertilizer replacement value of cattle slurry applied with a trailing-shoe application system. *Grass and Forage Science*

69(3): 488-501. DOI: 10.1111/gfs.12051.

- Laws, J. A., Smith, K. A., Jackson, D. R., Pain, B. F. (2002). Effects of slurry application method and timing on grass silage quality. *The Journal of Agricultural Science* 139(4): 371-384. DOI: 10.1017/s0021859602002708.
- Liu, Y.-Y., Haynes, R. J. (2013). Effect of disposal of effluent and paunch from a meat processing factory on soil chemical and microbial properties. *Water, Air, and Soil Pollution* 224(9): 1-14. DOI: 10.1007/s11270-013-1655-5.
- Lowe, K. F., Bowdler, T. M., Casey, N. D., Moss, R. J. (1999). Performance of temperate perennial pastures in the Australian subtropics 1. Yield, persistence and pasture quality. *Australian Journal of Experimental Agriculture* 39(6): 663-676. DOI: 10.1071/EA98021.
- Lucas, S. T., Weil, R. R. (2012). Can a labile carbon test be used to predict crop responses to improve soil organic matter management? *Agronomy Journal* 104(4): 1160-1170. DOI: 10.2134/agronj2011.0415.
- McKenzie, F. R. (1996). The influence of applied nitrogen on herbage yield and quality of *Lolium perenne* L. pastures during the establishment year under subtropical conditions. *South African Journal of Plant and Soil* 13(1): 22-26. DOI: 10.1080/02571862.1996.10634369.
- Ministry of Agriculture, Fisheries Food. (1986). *The analysis of agricultural materials*. Reference Book No.: 427 (3rd Edition), London: The Stationery Office.
- Misselbrook, T. H., Laws, J. A., Pain, B. F. (1996). Surface application and shallow injection of cattle slurry on grassland: Nitrogen losses, herbage yields and nitrogen recoveries. *Grass and Forage Science* 51(3): 270-277. DOI: 10.1111/j.1365-2494.1996.tb02062.x.
- Morrison, J., Jackson, M. V., Sparrow, P. E. (1980). *The response of perennial ryegrass to fertilizer nitrogen in relation to climate and soil.* Joint ADAS/GRI Grassland Manuring Trial GM20, Technical Report No.: 27. Berkshire, U.K.: Grassland Research Institute.
- Mouat, M. C. H., Nes, P. (1983). Effect of the interaction of nitrogen and phosphorus on the growth of ryegrass. *New Zealand Journal of Agricultural Research* 26(3): 333-336. DOI: 10.1080/00288233.1983.10427039.
- Petersen, J. (2003). Nitrogen fertilizer replacement value of sewage sludge, composted household waste and farmyard manure. *The Journal of Agricultural Science* 140(2): 169-182. DOI: 10.1017/S0021859603003010.
- Petrovic, Z., Djordjevic, V., Milicevic, D., Nastasijevic, I., Parunovic, N. (2015). Meat production and consumption: environmental consequences. *Procedia Food Science* 5: 235-238. DOI: 10.1016/j.profoo.2015.09.041.
- Peyraud, J. L., Astigarraga, L. (1998). Review of the effect of nitrogen fertilization on the chemical composition, intake, digestion and nutritive value of fresh herbage: consequences on animal nutrition and N balance. *Animal Feed Science and Technology* 72(3-4): 235-259. DOI: 10.1016/S0377-8401(97)00191-0.
- Pikula, D., Ten Berge, H. F. M., Goedhart, P. W., Schröder, J. J. (2016). Apparent nitrogen fertilizer replacement value of grassclover leys and of farmyard manure in an arable rotation. Part II: Farmyard manure. *Soil Use and Management* 32: 20-31. DOI: 10.1111/sum.12245.
- Rayment, G. E., Lyons, D. J. (2011). Soil chemical methods Australasia. Collingwood, VIC, Australia: CSIRO Publishing.
- Reid, D. (1985). A comparison of the yield responses of four grasses to a wide range of nitrogen application rates. *The Journal of Agricultural Science* 105(2): 381-387. DOI: 10.1017/s0021859600056434.
- Reyes, J., Schellberg, J., Siebert, S., Elsaesser, M., Adam, J., Ewert, F. (2015). Improved estimation of nitrogen uptake in grasslands using the nitrogen dilution curve. *Agronomy for Sustainable Development* 35(4): 1561-1570. DOI: 10.1007/s13593-015-0321-2.
- Schnitzer, M. (1982). Organic matter characterization. *In*: Page, A. L. (Ed.). Methods of soil analysis: chemical and microbiological properties, Part 2 (2nd Edition). Agronomy Monograph No.: 9, pp. 581-594. Madison, WI: American Society of Agronomy.
- Six, L., Velghe, F., Verstichel, S., De Meester, S. (2016). Chapter 11: Sustainability considerations on the valorization of organic waste. In: Poltronieri, P, D'Urso, O. F. (Editors). Biotransformation of agricultural waste and by-products: The food, feed, fibre, fuel (4F) economy, pp. 287-307. Elsevier B. V. DOI: 10.1016/b978-0-12-803622-8.00011-2.
- Sparrow, P. E. (1979). The comparison of five response curves for representing the relationship between the annual dry-matter yield of grass herbage and fertilizer nitrogen. *The Journal of Agricultural Science* 93(3): 513-520. DOI: 10.1017/S0021859600038910.
- Spence, J. (2012). Renewable energy in the Australian red meat processing industry and the viability of paunch as a biofuel. Master of Science Thesis. Toowoomba, QLD, Australia: University of Southern Queensland.
- Tekrony, D. M. (1983). Seed vigor testing 1982. Journal of Seed Technology 8: 55-60.
- Tompkins, D. K., Chaw, D., Abiola, A. T. (1998). Effect of windrow composting on weed seed germination and viability. *Compost Science and Utilization* 6(1): 30-34. DOI: 10.1080/1065657X.1998.10701906.
- Vivian, R., Silva, A. A., Gimenes Jr., M., Fagan, E. B., Ruiz, S. T., Labonia, V. (2008). Dormência em sementes de plantas daninhas como mecanismo de sobrevivência: breve revisão. *Planta Daninha* 26(3): 695-706. DOI: 10.1590/S0100-83582008000300026.
- Vogeler, I., Mackay, A., Vibart, R., Rendel, J., Beautrais, J., Dennis, S. (2016). Effect of inter-annual variability in pasture growth and irrigation response on farm productivity and profitability based on biophysical and farm systems modelling. *Science of the Total Environment* 565: 564-575. DOI: 10.1016/j.scitotenv.2016.05.006.

VSN (2013). GenStat release 16th Edition Reference Manual. Hemel Hempstead, U.K.: VSN International.

- Walkley, A., Black, I. A. (1934). An examination of the degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science* 37(1): 29-38.
- Wilman, D., Koocheki, A., Lwoga, A. B., Droushiotis, D., Shim, J. S. (1976). The effect of interval between harvests and nitrogen application on the numbers and weights of tillers and leaves in four ryegrass varieties. *The Journal of Agricultural Science* 87(1): 45-57. DOI: 10.1017/s0021859600026575.
- Wilman, D., Mohamed, A. A. (1980). Response to nitrogen application and interval between harvests in five grasses. I. Drymatter yield, nitrogen content and yield, numbers and weights of tillers, and proportion of crop fractions. *Fertilizer Research* 1(4): 245-263. DOI: 10.1007/bf01074197.
- Wilson, D. (1992). Methods of disposal of paunch contents with emphasis on composting. In: Pearson, A. M., Dutson, T. R. (Editors). Inedible meat by-products, Vol.: 8, pp. 265-281. Dordrecht: Springer Netherlands. DOI: 10.1007/978-94-011-7933-1_11.