

# DYNAMICS AND PREDICTABILITY OF EL NIÑO–SOUTHERN OSCILLATION

An Australian Perspective on Progress and Challenges

AGUS SANTOSO, HARRY HENDON, ANDREW WATKINS, SCOTT POWER, DIETMAR DOMMENGET, MATTHEW H. ENGLAND, LEEA FRANKCOMBE, NEIL J. HOLBROOK, RYAN HOLMES, PANDORA HOPE, EUN-PA LIM, JING-JIA LUO, SHAYNE MCGREGOR, SONJA NESKE, HANH NGUYEN, ACACIA PEPLER, HARUN RASHID, ALEX SEN GUPTA, ANDRÉA S. TASCHETTO, GUOMIN WANG, ESTEBAN ABELLÁN, ARNOLD SULLIVAN, MAURICE F. HUGUENIN, FELICITY GAMBLE, AND FRANCOIS DELAGE

Many scientific challenges remain for managing the risk of future ENSO impacts in countries like Australia that are strongly affected by ENSO event diversity.

**MOTIVATION.** *The Australian context.* El Niño–Southern Oscillation (ENSO) has long been recognized to strongly influence global and regional climate. Australian climate is particularly impacted by ENSO (e.g., McBride and Nicholls 1983; Ropelewski and Halpert 1987; Power et al. 1998). The associated changes in circulation, rainfall, and temperatures are strong enough to impact its terrestrial and marine ecosystems (e.g., Nicholls 1985, 1991; Norman and Nicholls 1991; Holbrook et al. 2009). Although the impact can vary markedly from decade to decade (Power et al. 1999), bushfires, heat waves, and droughts generally tend to be more severe during El Niño years (e.g., Williams and Karoly 1999; Loughran et al. 2017), while the frequency of tropical cyclones across the north and flooding throughout much of the east tend to be enhanced during La Niña (e.g., Werner and Holbrook 2011; Power and Callaghan 2016). To provide timely information on the likelihood of upcoming disruptions of climate, ENSO Outlooks have been issued routinely by the

Australian Bureau of Meteorology (BOM) since 2000.

The complex dynamics of ENSO manifest in diverse spatial and temporal evolution across events that lead to differing regional impacts (e.g., Power et al. 1999; Ashok et al. 2007; Wang and Hendon 2007; Capotondi et al. 2015a). For instance, in Australia, the magnitude of an El Niño event alone does not provide clear guidance on its impacts (Power et al. 2006; Wang and Hendon 2007; Chung and Power 2017). For example, the impact of the 1997/98 extreme El Niño was limited to the southeastern region and Tasmania, but much more severe and widespread drought occurred during the moderate 2002/03 El Niño (Wang and Hendon 2007; Taschetto and England 2009; Lim and Hendon 2015), leading to a massive 25% drop in agricultural output (Lu and Hedley 2004).

The 2002/03 event was not only notably weaker in intensity than the 1997/98 extreme event, but also exhibited a characteristically different pattern of sea surface temperature (SST) anomalies. The 1997/98 El

Niño had SST anomalies ( $\sim+3^{\circ}\text{C}$ ) that peaked toward South America while those during the 2002/03 event peaked in the central Pacific ( $\sim+1^{\circ}\text{C}$ ). The contrast in spatial patterns fits the notion of two archetype structures of ENSO: eastern Pacific (EP) and central Pacific (CP) events (e.g., Ashok et al. 2007; Kao and Yu 2009) following an earlier assessment of ENSO SST patterns by Trenberth and Stepaniak (2001). This is illustrated in Fig. 1, which also shows that an event may not necessarily fall into either category: a pattern that is a mix of EP and CP types is possible as part of the ENSO continuum arising from nonlinear dynamics, stochasticity, and remote forcing, which can also give rise to temporal evolution diversity (e.g., Takahashi et al. 2011; Dommenges et al. 2013; Lee et al. 2014; Takahashi and Dewitte 2016). This intrinsic ENSO complexity was recently summarized by Timmermann et al. (2018). As such, a clear classification of certain ENSO events into EP and CP can be difficult and can be sensitive to the choice of index (e.g., Capotondi et al. 2015a).

The typical surface temperature and rainfall anomaly patterns associated with EP and CP ENSO are different over Australia, as well as other regions across the globe (Fig. 2). Specifically, CP events tend to be associated with larger and more widespread rainfall and temperature changes in Australia than EP events—analogueous to the difference in impacts between the 2002/03 and 1997/98 events, even though EP El Niños tend to be stronger than CP events (Fig. 1; see also Capotondi et al. 2015a). Investigating the cause for the differing impacts between EP and CP events is still an open area of research, but based on

investigating the difference between 2002/03 and 1997/98 events, the contrast may be due to the forcing center of CP events that is closer to Australia (Wang and Hendon 2007; Lim and Hendon 2015). This could also illustrate the more general result, that impacts in Australia are more tightly linked to the magnitude of La Niña than they are to the magnitude of El Niño (Power et al. 2006), as stronger La Niña events tend to be of a CP type (Fig. 1). As such, predicting the spatial structure of ENSO events is important for impact preparedness over regions like Australia (Hendon et al. 2009). Apart from classification of individual events, other factors such as event precursors, local processes (e.g., antecedent soil moisture, anomalies in regional seas), and random disturbances, as well as other modes of climate variability, also matter in determining impacts.

The Australian continent extends from the tropics to the midlatitudes and is surrounded by warm tropical Indo-Pacific oceans to the north and the Southern Ocean to the south. Thus, it is not only affected by direct tropical impacts of ENSO via the Southern Oscillation but also by extratropical teleconnections due to ENSO-induced changes in tropical convection. Furthermore, Australian climate is affected by a rich interplay between ENSO and other climatic events such as the Indian Ocean dipole (IOD) and the southern annular mode (SAM; e.g., Hendon et al. 2007; Meyers et al. 2007; Risbey et al. 2009; Cai et al. 2011; Taschetto et al. 2011; Pui et al. 2012; Lim and Hendon 2015). Ocean surface temperature variations surrounding northern Australia, which tend to covary with ENSO, also exert strong influence on

**AFFILIATIONS:** SANTOSO—Climate Change Research Centre, and ARC Centre of Excellence for Climate System Science, University of New South Wales, Sydney, New South Wales, and Centre for Southern Hemisphere Oceans Research, CSIRO Oceans and Atmosphere, Hobart, Tasmania, Australia; HENDON, WATKINS, POWER, HOPE, LIM, NGUYEN, PEPLER, WANG, GAMBLE, AND DELAGE—Australian Bureau of Meteorology, Melbourne, Victoria, Australia; DOMMENGET, MCGREGOR, AND NESKE—School of Earth, Atmosphere, and Environment, and ARC Centre of Excellence for Climate System Science, Monash University, Melbourne, Victoria, Australia; ENGLAND, FRANKCOMBE, SEN GUPTA, AND TASCETTO—Climate Change Research Centre, and ARC Centre of Excellence for Climate Extremes, University of New South Wales, Sydney, New South Wales, Australia; HOLBROOK—Institute for Marine and Antarctic Studies, and ARC Centre of Excellence for Climate Extremes, University of Tasmania, Hobart, Tasmania, Australia; HOLMES—Climate Change Research Centre, and ARC Centre of Excellence for Climate Extremes, and School of Mathematics and Statistics, University of New South Wales, Sydney, New South Wales, Australia; LUO—Australian

Bureau of Meteorology, Melbourne, Victoria, Australia, and Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters/KLME/ILCEC, Nanjing University of Information Science and Technology, Nanjing, China; RASHID AND SULLIVAN—CSIRO Oceans and Atmosphere, Aspendale, Melbourne, Victoria, Australia; ABELLÁN—Climate Change Research Centre, and ARC Centre of Excellence for Climate System Science, University of New South Wales, Sydney, New South Wales, Australia; HUGUENIN—Institute for Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland  
**CORRESPONDING AUTHOR:** Agus Santoso, a.santoso@unsw.edu.au

*The abstract for this article can be found in this issue, following the table of contents.*

DOI:10.1175/BAMS-D-18-0057.1

In final form 16 September 2018


©2019 American Meteorological Society

For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](#).

Australian climate (e.g., Drosowsky and Chambers 2001; Hendon et al. 2012; Ummenhofer et al. 2015) and may even affect development of ENSO itself (Nicholls 1984).

This complexity of impacts and interactions combined with the uniqueness of every ENSO event poses grand challenges for predicting Australian climate. For example, unlike the extreme 1997/98

event, the extreme 1982/83 El Niño, which is also classified as an EP event, had a particularly strong impact on Australia, likely due to a relatively strong cold sea surface anomaly to the north-northeast of Australia (van Rensch et al. 2015). Severe drought gripped the eastern half of the country, marked by the historically catastrophic “Ash Wednesday” bushfires in the southeast (Voice and Gauntlett 1984). For these



**FIG. 1. ENSO diversity over 1980–2017.** (a) EP ENSO index (EPI) vs CP index (CPI) averaged over DJF when ENSO events typically peak, where circle size corresponds to ENSO amplitude and the color indicates the type (EP, CP, EP/CP). (b)–(e) Composite of DJF SST anomalies for each type of ENSO event. (f) SST anomaly over the equatorial Pacific (averaged over 5°S–5°N) marked by different colors that signify event types in (a). The EPI and CPI are based on those of Sullivan et al. (2016), defined as  $\text{Niño-3} - 0.5 \times \text{Niño-4}$  and  $\text{Niño-4} - 0.5 \times \text{Niño-3}$ , respectively (where the Niño indices are first normalized). Niño-3 and Niño-4 indices are SST anomalies averaged over (5°S–5°N, 150°–90°W) and (5°S–5°N, 160°E–150°W), respectively. An arbitrary threshold (Thr) can be applied to the indices to classify each year into EP, CP, or a mix (EP/CP). In this case, 0.7 of the index standard deviation (sdev) is used (dotted lines), with the 1982/83 and 1997/98 extreme El Niños being classified as EP events (dark red) in which  $\text{EPI} > \text{EP Thr}$  and  $\text{CPI} < \text{CP Thr}$ . In this way, the 2015/16 and 1991/92 El Niños can be classified as both EP and CP (red), and the events in yellow are CP El Niños ( $\text{CPI} > \text{CP Thr}$ ,  $\text{EPI} < \text{EP Thr}$ ). The same applies for La Niñas but using negative thresholds. Note how the event classification can change with subtle shift in the thresholds. The size of the circles corresponds to the magnitude of the Niño-3.4 anomaly: large circles for  $|\text{Niño-3.4}| > 1.8$  sdev, medium circles for  $1 \text{ sdev} < |\text{Niño-3.4}| < 1.8$  sdev, and small circles for  $0.5 \text{ sdev} < |\text{Niño-3.4}| < 1 \text{ sdev}$ . Gray circles are considered as neutral years ( $|\text{Niño-3.4}| < 0.5$  sdev). The NOAA Extended Reconstructed SST version 5 (ERSSTv5; Huang et al. 2017) is used in this analysis with linear trends removed.

reasons, and given Australia's susceptibility to future climate change, ENSO is at the forefront of climate research in Australia.


**Unexpected turns of events.** Following the extreme El Niño events in 1982/83 and 1997/98, the most recent major El Niño event occurred in 2015/16 (Blunden and Arndt 2016; Xue and Kumar 2017; L'Heureux et al. 2017). This first extreme El Niño of the twenty-first century (Santoso et al. 2017) followed a "false alarm" in 2014. In 2014, the equatorial Pacific warm water volume (WWV) increased rapidly during the austral autumn following a strong westerly wind burst (WWB) event, reaching a level not seen since 1997 (McPhaden 2015). Increased WWV and increased activity of WWBs are typical precursors for an El Niño (see Table 1 for WWV and WWB definitions).

However, the much anticipated big El Niño did not emerge at the end of 2014 (Hannam 2014), but it did instead in 2015.

The "roller coaster" evolution of the 2014–16 events and their prediction are illustrated in Fig. 3, using the ENSO Outlook indicator from the BOM. The BOM raised its ENSO Outlook to "watch" in February and March 2014, and subsequently elevated to "alert" in April–July, indicating the increasing possibility of an El Niño (of any magnitude) later in the year. This coincided with a spike in WWV that often precedes El Niño events, in line with the ENSO recharge oscillator theory (Jin 1997). However, the outlook status was downgraded back to "watch" in August–October, and then elevated again to "alert" in November–January. This outlook variation appears to be in line with WWV decline since the previous April before a slight increase again around July.

A strong El Niño never materialized, which was later shown to be due to a combination of impeding factors, such as muted WWB activity (Menkes et al. 2014) and an occurrence of intense easterly wind burst (Hu and Fedorov 2016), as well as a mean state associated with the negative phase of the interdecadal Pacific oscillation (IPO) that is less favorable for Bjerknes feedbacks at the root of El Niño growth (Wang and Hendon 2017) and an anomalously warm Indian Ocean surface (Dong and McPhaden 2018)—both factors are associated with stronger Pacific Walker circulation. The tropical Pacific was nonetheless left anomalously warm, but fell short to being considered an El Niño condition (e.g., Santoso et al. 2017).

In early 2015 clear signs of an emerging El Niño were detected, and the outlook status was raised to "event" in May 2015 (Watkins 2015). A strong El Niño developed in the



**FIG. 2.** Surface air temperature (SAT) and rainfall anomaly patterns associated with (a) EP and (b) CP ENSO shown as the regression of SAT (color shading) and rainfall anomalies (contours) against EP and CP ENSO indices (see Fig. 1 caption). Units are in  $^{\circ}\text{C}$  and  $\text{mm day}^{-1}$  per standard deviation. The analysis uses monthly data of the NCEP–NCAR reanalysis (Kalnay et al. 1996) for SAT, CPC Merged Analysis of Precipitation (CMAP; Xie and Arkin 1997), and ERSSTv5 (Huang et al. 2017) to calculate the ENSO indices from 1980 to 2016, with monthly climatology and long-term trends removed. Stippling indicates rainfall regression coefficients that are statistically significant above the 95% level.

**TABLE 1. Definitions of terms discussed in this article.**

Terminology	Definition
Bjerknes feedback (Bjerknes 1969)	Positive air–sea coupled feedback along the equator in which a positive SST anomaly during an El Niño growth phase induces westerly wind anomalies that deepen the thermocline, thereby reinforcing the positive SST anomaly, and the cycle continues taking El Niño to its peak. The converse occurs during a La Niña.
Interdecadal Pacific oscillation (IPO; Power et al. 1999; Newman et al. 2016; Henley et al. 2015)	Ocean–atmosphere variability in the Pacific Ocean operating on 10–30-yr time scales with a near-global pattern resembling that of El Niño and La Niña during its positive/warm and negative/cold phases, respectively. The IPO could be a long-term integration of various processes, including interannual and decadal components of ENSO variability, stochastic air–sea fluxes, and other sources of low-frequency variability.
Indian Ocean dipole (IOD; Saji et al. 1999)	Year-to-year climate variability in the Indian Ocean that peaks in austral spring, with its positive phase exhibiting a pool of anomalously cold sea surface off Java–Sumatra and anomalously warm sea surface off Africa. Such a pattern is associated with weaker Walker circulation over the Indian Ocean and often coincides, but not always, with a developing El Niño. The converse occurs during the negative phase and La Niña.
Southern annular mode (SAM; Thompson and Wallace 2000)	Vacillations in atmospheric pressure over the extratropical Southern Hemisphere to Antarctica associated with stronger westerly wind at high latitudes and weaker westerlies at midlatitudes during its positive phase, and conversely during the negative phase.
Walker circulation (Bjerknes 1969)	Large-scale zonal atmospheric circulation in the tropical Pacific marked by easterly winds blowing from the colder eastern Pacific toward the western Pacific warm pool where warm air rises and moves eastward as it loses moisture before eventually descending in the eastern Pacific. The Walker circulation weakens during an El Niño and strengthens during a La Niña, and the Southern Oscillation refers to the associated vacillation of atmospheric pressure in the tropical western and eastern Pacific.
Warm water volume (WWV; Meinen and McPhaden 2000)	Volume of water above 20°C isotherm across the equatorial Pacific (5°S–5°N, 120°E–80°W) as a proxy of upper equatorial Pacific Ocean heat content. Anomalously high WWV around austral autumn is a necessary condition for an El Niño at the end of the year. The opposite is true for La Niña.
Westerly wind burst (WWB; Vecchi and Harrison 2000; Puy et al. 2016)	Sustained west-to-east winds over the western and central equatorial Pacific, typically exceeding a certain threshold (e.g., 2 m s <sup>-1</sup> ) and lasting more than a few days.


latter half of 2015. The tropical Pacific then cooled, with a borderline La Niña developing in austral summer of 2016/17 declared by some agencies (but not the BOM) followed by the BOM's official declaration of a weak La Niña in December 2017.

Amid the widespread speculation in early 2014 of a strong El Niño that year, the BOM's coupled model, Predictive Ocean Atmosphere Model for Australia (POAMA), predicted only a weak event when initialized in austral autumn 2014 (Wang and Hendon 2017, see their Fig. 3a). In contrast, other models surveyed by the BOM predicted a strong El Niño for 2014 ([www.bom.gov.au/climate/ahead/archive/models/201405-ms.shtml](http://www.bom.gov.au/climate/ahead/archive/models/201405-ms.shtml)). On the other hand, while issuing a stronger forecast for El Niño in 2015 than in 2014, POAMA initially underestimated the strength of the 2015/16 El Niño [see Fig. 3b of Wang and Hendon (2017); the observed Niño-3 falls outside the forecast 5%–95% uncertainty range] and was weaker than the other surveyed models, until POAMA was

initialized with late austral winter conditions (see the BOM's 2015 archive; e.g., [www.bom.gov.au/climate/ahead/archive/models/201508-ms.shtml](http://www.bom.gov.au/climate/ahead/archive/models/201508-ms.shtml)). This is not surprising though, as predicting the magnitude of ENSO is challenging, more so than predicting the phase, especially early in the year when signal-to-noise ratio is low.

The challenge in anticipating and predicting the evolution and magnitude of the 2014–16 chain of events led to much retrospection about the state of our understanding of ENSO dynamics beyond the classical recharge–discharge oscillator theory, as well as the current state-of-the-art climate models used to make predictions.

At an international ENSO workshop held in Sydney, Australia, in 2015, key aspects of ENSO extremes and the associated open questions were discussed (Santoso et al. 2015). However, at that meeting, our knowledge of extreme El Niño was largely based on the 1982/83 and 1997/98



**FIG. 3. ENSO Outlook of the BOM along with the anomaly of temperature averaged over the top 300 m in the equatorial Pacific Ocean as a proxy of WWV (black) and the Niño-3.4 index (green) from Jan 2013 to Feb 2018.** The outlook is produced based on a survey of BOM's own coupled seasonal forecast model and seven others from leading international climate agencies (generally WMO Global Producing Centres of Long Range Forecasts). The explanation for the outlook can be found in [www.bom.gov.au/climate/enso/outlook/#tabs=ENSO-Outlook-history](http://www.bom.gov.au/climate/enso/outlook/#tabs=ENSO-Outlook-history). The WWV data can be accessed from [www.pmel.noaa.gov/tao/wwv/data/](http://www.pmel.noaa.gov/tao/wwv/data/). The Niño-3.4 index is an average of SST over (5°S–5°N, 170°–120°W) calculated based on the ERSSTv5 dataset ([www.esrl.noaa.gov/psd/data/gridded/data.noaa.ersst.v5.html](http://www.esrl.noaa.gov/psd/data/gridded/data.noaa.ersst.v5.html)).

events—the only two extreme El Niño events in the modern instrumental record that showed distinct characteristics from other ENSO events. These characteristics include i) intense WWB activity in the western/central Pacific during event onset and development phases; ii) a dramatic eastward and equatorward shift of atmospheric convection as El Niño emerges and matures, thereby inducing unusually high rainfall in the climatologically dry and cold eastern equatorial Pacific; and iii) prominent eastward propagation of anomalous SSTs along the equatorial Pacific Ocean over event onset to decay phase.<sup>1</sup> However, the latter two properties, which had previously been thought to typify an extreme El Niño, were less apparent during the 2015/16 El Niño (Santoso et al. 2017). In particular, while the 2015/16 El Niño did produce heavy rainfall over the eastern equatorial Pacific with December–February (DJF) average rainfall in the Niño-3 region (5°S–5°N, 150°–90°W) close to 5 mm day<sup>-1</sup>, a threshold used by Cai et al. (2014) to define an extreme El Niño, it exhibited record-breaking rainfall over the central Pacific, in stark contrast to the relatively weak El Niño-related rainfall in 1982/83 and 1997/98 events (Santoso et al. 2017).

The peculiarity of the 2015/16 event, the volatile 2014/15 ENSO outlook (Fig. 3), and the complex

ENSO behavior and its impacts over Australia (see “The Australian context” section) motivated a second workshop on ENSO dynamics and prediction that was held in Sydney in November 2017 and involved 25 Australian ENSO researchers ([www.climate-science.org.au/content/1182-enso-dynamics-workshop](http://www.climate-science.org.au/content/1182-enso-dynamics-workshop)). Here we present the outcomes of this workshop, outlining recent research progress, knowledge gaps, impediments, and recommendations to further advance ENSO research and prediction systems and service. These issues are discussed within each of five themes outlined in the “Discussions” section, based largely on the studies presented by the workshop participants as referenced therein. A summary is provided in the last section along with closing remarks on infrastructures and synergy behind ENSO research in Australia. Definitions of some terminologies discussed in the paper are provided in Table 1.

#### **DISCUSSIONS. Insights from the 2015/16 El Niño.**

The emergence of the strong 2015/16 El Niño showed that an extreme El Niño does not necessarily exhibit SST anomalies peaking toward the far eastern Pacific as in the 1982/83 and 1997/98 events, which have until recently been used as a benchmark for defining an extreme El Niño [see Santoso et al. (2017) for a review]. The global climate context for the 2015/16 El Niño was different from that for the 1982/83 and 1997/98 events (see also Newman et al. 2018). For instance, there was a much more significant and persistent signature of extratropical influence in the 2015/16 El Niño, with warm SST anomalies extending from the

<sup>1</sup> The propagation signature is diagnosed as the time–longitude slope of maximum equatorial SST anomaly (5°N–5°S average) over 160°E–80°W from May of the El Niño development year to the following May when the event subsides, following Santoso et al. (2013).

northeastern Pacific to the central Pacific associated with the North Pacific meridional mode, than in the 1982/83 and 1997/98 events (Santoso et al. 2017; Paek et al. 2017).

Unlike the previous two extremes, the large amplitude of the 2015/16 El Niño was built upon an already abnormally warm tropical Pacific from 2014, rather than relying solely on a vigorous Bjerknes feedback (Abellán et al. 2018). The weak 2014/15 El Niño-like condition prevented a large discharge of warm water out of the equatorial Pacific (Levine and McPhaden 2016) that would normally occur following a strong El Niño. This allowed anomalous equatorial warming to persist into 2015, priming the ocean for the subsequent El Niño. The suite of processes leading up to the 2015/16 El Niño, along with a buildup of ocean heat content in the off-equatorial western Pacific over the previous decade, may have triggered a shift in the phase of the IPO, from negative before to positive after 2014 (Meehl et al. 2016). This can be invoked to partially explain the difference in amplitude between the 2014/15 and 2015/16 events (Wang and Hendon 2017), as a mean state associated with a positive IPO is more conducive to the Bjerknes positive feedbacks for El Niño development in the eastern Pacific (e.g., Zhao et al. 2016)—although the IPO itself is, in turn, partially a long-term imprint of ENSO variability (Power and Colman 2006; Newman et al. 2016). The subsequent decay of the 2015/16 event was also different from the 1982/83 and 1997/98 events in that it had persistent warm SST anomalies near the date line, which lingered right through the austral fall of 2016 and significantly delayed the Bjerknes feedback required for the development of a following La Niña (Lim and Hendon 2017).

The distinctive characteristics of the 2015/16 extreme El Niño demonstrate that our observational record is still too short to fully sample the diversity of ENSO characteristics. Furthermore, there are uncertainties in observed SST data prior to the satellite era, especially before the 1950s when observations were sparse and ship recording practices were not homogeneous (e.g., Ishii et al. 2005). This is particularly the case for meteorological variables over the ocean, meaning that comparisons to past events using a single index based on SST, and other variables for that matter, may not be accurate in terms of relative strength or variability. These issues mean that multiple observational products and indices beyond the commonly used ENSO metrics in operational forecasting (e.g., Niño-3.4, WWV, WWB, Southern Oscillation index) are required to capture the diversity

of ENSO extremes. Refinement of existing indices is also needed to better describe ENSO event diversity (e.g., Sullivan et al. 2016). In addition, the background climate upon which ENSO evolves is changing due to greenhouse warming and internal multidecadal variability. To detect these long-term changes and the impact on ENSO characteristics, continuous high-quality observations are critical, and so are reliable paleoreconstructions for resolving characteristics of past ENSO events.

*ENSO predictability.* According to conventional ENSO theory (e.g., Jin 1997), WWV is a key precursor and hence predictor for ENSO. Consequently, WWV or the associated subsurface information is utilized in initializing forecast models to help alleviate the drop in ENSO prediction skill in austral autumn—widely known as the “(boreal) spring predictability barrier” (Webster and Yang 1992). However, anomalous WWV, while necessary, is not the only requirement for development of ENSO events, as demonstrated by the 2014 case. El Niño is typically triggered by a series of WWBs in the western/central Pacific (Vecchi and Harrison 2000). On the other hand, a negative WWV anomaly during austral autumn appears to be a better predictor for strong La Niña events than any other type of ENSO event (Santoso et al. 2017). A better understanding of the relationship between WWV, WWB, and ENSO is important in ENSO prediction.

On this front, Neske and McGregor (2018) showed that WWBs themselves can create a significant WWV response that can be decomposed into two components: the “adjusted response” (which relies on slow ocean dynamics associated with Rossby wave reflection as depicted by the recharge oscillator theory) and the “instantaneous response” (which represents surface Ekman transport in response to WWB). The adjusted response is identified as the source of predictability, and it has weakened since the start of the twenty-first century—the reason for this decline remains unclear but may be associated with a shift in the background climate toward the cold phase of the IPO (Zhao et al. 2016) through modulation of governing ENSO processes (WWBs, WWV, etc.). The instantaneous response, which has increased in prominence in recent decades, emphasizes that ENSO is event-like rather than cyclical. This research highlights that the balance of the two components can vary on decadal time scales, giving rise to decadal modulation of ENSO predictability.

ENSO predictability does not lie solely within the tropical Pacific. Climate variability in other oceanic basins also plays a role. Remote climate anomalies

induce atmospheric changes that are transmitted into the tropical Pacific through atmospheric planetary waves or changes in the Walker circulation, thereby affecting ENSO evolution. This remote influence is highlighted by two recent studies that investigated the predictability of recent La Niña events. Using two coupled models, Luo et al. (2017) showed that the Indian and Atlantic Ocean warming contributed to the two-year lead predictability of the 2010–12 series of La Niña events. The multiyear surface warming in these oceans enhanced the trade winds over the central–western Pacific that tend to favor La Niña development. The role of the Indian Ocean was emphasized by Lim and Hendon (2017), who showed that the appearance of La Niña in 2016, although weak, was promoted by earlier-than-normal development of a record strong negative IOD, which tends to enhance convection over the Indo-Pacific warm pool, thereby strengthening the Pacific trade winds. They showed that using realistic ocean conditions in the Indian Ocean in late April 2016 was sufficient to produce the strong negative IOD during austral winter and spring and was necessary for delivering an improved La Niña forecast.


**Response to greenhouse forcing.** While the Pacific Ocean is projected to warm in the future, and that the anthropogenic warming is already evident in the western part of the basin (Wang et al. 2016), there is still uncertainty around whether ENSO events (typically measured through equatorial Pacific SST anomalies) will change in terms of their spatial patterns, amplitude, and frequency. Climate models produce contrasting projections due to the different relative importance of ENSO feedback processes, different patterns of changes in the mean climate, and different depictions of decadal variability (e.g., Collins et al. 2010; DiNezio et al. 2012; Kim et al. 2014a; Chen et al. 2017). The power spectra of ENSO SST variability also exhibit large discrepancies across models and paleoreconstructions (Hope et al. 2017). However, there is a better intermodel consensus on a general increase in ENSO-driven tropical rainfall in the equatorial central and eastern Pacific in response to global warming, reflecting the consensus for more mean warming in the eastern equatorial Pacific (Power et al. 2013; Chung et al. 2014).

According to phase 5 of the Coupled Model Intercomparison Project (CMIP5) scenario simulations, the projected weakening of the Walker circulation in the twenty-first century (e.g., Vecchi et al. 2006; Kociuba and Power 2015) with faster warming in the eastern equatorial Pacific than in

the surrounding oceans is expected to shift atmospheric convection into this usually cold and dry region, resulting in more occurrences of heavy precipitation that characterize an extreme El Niño (Cai et al. 2014). The projection does not arise from the climatological increase in mean rainfall and is robust when using atmospheric vertical velocity (Cai et al. 2017). The associated weakening of the climatological equatorial ocean currents is expected to promote eastward-propagating SST anomalies—a characteristic of the 1982/83 and 1997/98 extreme El Niño events (Santoso et al. 2013). On the other hand, increased occurrences of extreme La Niña events characterized by anomalous surface cooling in the central Pacific could also arise due to faster warming of the Maritime Continent and eastern Pacific than the central Pacific (Cai et al. 2015b). The warming background climate can enhance ENSO teleconnection and thus its impact, even if the SST anomalies themselves do not intensify (Cai et al. 2015a; Power and Delage 2018). Even if global warming is kept below 1.5° or 2°C, the risk associated with increased extreme El Niño frequency and ENSO-related major rainfall disruptions is likely to persist or even increase (Wang et al. 2017; Power et al. 2017a). In fact, global warming might have already made ENSO events more disruptive (Power et al. 2017a), enhancing ENSO-driven variability in many regions around the world (Bonfils et al. 2015; Power and Delage 2018).

These projections may be sensitive to model deficiencies in simulating ENSO. As shown by Vijayeta and Dommenges (2018) using the recharge oscillator framework, models tend to underestimate ENSO feedback processes. Realistic simulation of ENSO behavior may stem from error compensation rather than from correct simulation of the governing feedback processes. Confidence in projections is also reduced by the inability of climate models to capture decadal La Niña–like trends as strong as those observed in the Pacific in recent decades (e.g., Kociuba and Power 2015; England et al. 2014), and a tendency to overestimate Pacific warming over the past 50 years (Power et al. 2017b). Possible reasons include model underestimation of internal variability (Kociuba and Power 2015; Power et al. 2017b), biases in the upper-ocean thermal stratification (Kohyama et al. 2017), and biases in the interbasin warming contrast across the three oceans and in the SST–cloud forcing feedback (Luo et al. 2018). To reduce uncertainty in ENSO future projections, it is clear that much work needs to be done to improve climate models.





**FIG. 4. Climate model bias in SST and ENSO seasonality as part of the many challenges facing ENSO researchers. (a) Difference in SST between multimodel ensemble mean and observed (ERSSTv5) exhibiting the classical cold tongue bias (color shading). The climatological observed and multimodel mean SSTs are shown in black and red contours, respectively. Twenty CMIP5 models are utilized, with stippling indicating 18 or more models exhibiting the same sign in the bias. (b) Standard deviation of the Niño-3.4 index in the CMIP5 models with annual mean of the standard deviation removed. Observed and multimodel mean seasonal climatologies are shown in thick black and red dashed lines, respectively; both indicate a peak of ENSO variability around austral summer and lowest variability in austral autumn. The most biased models in terms of the annual cycle are highlighted in thick colored lines.**

*ENSO modeling.* Poor simulation of ENSO is often linked to the persistent “cold tongue” bias in which the cold upwelled water in the eastern equatorial Pacific extends too far west toward the Maritime Continent (Fig. 4a). A double intertropical convergence zone (ITCZ) is also associated with this cold tongue bias. These biases can affect ENSO simulation through misrepresentation of air–sea feedbacks (Kim et al. 2014b; Wengel et al. 2018). Graham et al. (2017) showed that the cold tongue bias leads to a propensity for occurrences of spatially double-peaked ENSO SST anomalies (peaking concurrently in both the eastern and central Pacific), which are not apparent in historical observations.

Another important ENSO feature is its synchronization to the annual cycle, with peak SST anomalies typically occurring during austral summer. However, many models still do not represent this accurately (Fig. 4b). Worse, in some models (such as those highlighted in Fig. 4b) the seasonality is completely reversed (Taschetto et al. 2014). A suite of important processes shapes ENSO seasonality, and the incorrect seasonality indicates that the underlying ENSO dynamics (e.g., SST–cloud and thermocline feedbacks) in the models are unlikely to be correct (Rashid and Hirst 2016). This, together with the bias in anomaly patterns, has ramifications for determining ENSO teleconnections and predicting ENSO impacts on rainfall, for instance.

How these unrealistic ENSO features affect future projections needs to be carefully considered and investigated through detailed analysis of the underlying coupled feedbacks (e.g., Guilyardi et al. 2016; Capotondi et al. 2015b), which are also dependent upon the mean state. As an example, Rashid et al. (2016) found that the strengths of the zonal wind stress forcing and wind–convection coupling simulated by the CMIP5 models largely determine whether the ENSO amplitude will increase or decrease under global warming in those models. Importantly, improvement of the model’s mean state is critical. For instance, by taking into account the effect of ocean currents on the momentum transfer to the atmosphere (Pacanowski 1987), Luo et al. (2005) found a notable reduction in the cold tongue bias in their climate model.

The equatorial Pacific cold tongue is also affected by small-scale oceanic processes such as tropical instability waves (TIWs). TIWs are not well resolved by current state-of-the-art climate models, which are still run at relatively coarse ocean resolution. TIWs heat the Pacific cold tongue at a rate comparable to atmospheric heating (up to  $1^{\circ}\text{C month}^{-1}$ ; e.g., Menkes et al. 2006) and can potentially be an important nonlinear negative feedback on ENSO (An and Jin 2004). Research into TIWs’ influence on ENSO is still limited. Holmes et al. (2018) address how TIWs modify the response of the CP and EP to WWBs, providing a first estimate of TIWs contribution to ENSO irregularity, with implications for ENSO prediction (also see Ham and Kang 2011).

In addition, the Pacific cold tongue is a region of strong atmospheric heat uptake and vigorous turbulent mixing that transports this heat into the ocean interior. Recent research by Holmes et al. (2019) has highlighted the global significance of air–sea fluxes and turbulent mixing in this region for modulating ocean heat uptake. However, the parameterization of this vertical mixing in climate models remains a difficult task, and improvements are required in order to reduce model biases (e.g., Sasaki et al. 2013; Zhu and Zhang 2018).

*Low-frequency variability.* ENSO properties vary on decadal and longer time scales (e.g., Holbrook et al. 2014; Wittenberg 2015; Power and Smith 2007) through changes in air–sea feedbacks linked to noise, chaotic dynamics, and slow variations in the mean state (e.g., Wittenberg 2009; Newman et al. 2011; Wittenberg et al. 2014; Zhao et al. 2016). While the mean-state variations are thought to be in part a rectified effect of changes in ENSO (e.g., Power and Colman 2006; Ogata et al. 2013), they also affect

the interactions between ENSO and other modes of variability such as the SAM (e.g., Lim et al. 2016), as well as their eventual impact, such as on atmospheric cyclones and anticyclones. It is therefore crucial to understand the processes governing mean-state changes and their impact on ENSO.

The impact of mean-state change was highlighted by the 1970s climate shift that saw the Pacific climate system transition into a positive IPO that lasted until the late 1990s. Since then, the issue has been reignited by the recent short-lived slowdown in global surface warming that coincided with a negative IPO phase (approximately 1999–2012). In contrast to the preceding positive IPO, this negative IPO period was marked by a lack of a strong El Niño, more frequent CP El Niños, more prominent La Niñas (e.g., see Fig. 3 of Santoso et al. 2017), and reduced seasonal predictability (Zhao et al. 2016). This change in ENSO characteristics and predictability was attributed by Zhao et al. (2016) to reduction in the strength of the Bjerknes feedback in the central and eastern Pacific as a result of the IPO-related colder surface temperatures and enhanced mean Walker circulation.

The IPO can be partially explained as the accumulated response to interdecadal variability in ENSO activity (Power et al. 2006; Newman et al. 2016). However, other factors may be involved, and the extent to which the IPO influences ENSO activity requires further study. More research is needed into the processes governing decadal variability, including those during the slowdown period as well as periods of more rapid warming. For instance, the cause for the unprecedented strength of the Pacific trade winds during the global warming hiatus period (England et al. 2014) needs to be better understood, including the roles of interbasin warming contrast, radiative forcing, and ENSO rectification onto the mean state. Modeling studies have shown that these factors can influence the strength of Pacific climate change (e.g., Luo et al. 2018; Kohyama et al. 2017).

Recent studies indicate that warming trends in other ocean basins could be the cause for the record strong Pacific trade winds during 1999–2012. In particular, the role of the Atlantic warming trend appears to be important (e.g., McGregor et al. 2014; Luo et al. 2017). Models tend to underestimate the recent Pacific wind acceleration and to have strong climatological biases in the Atlantic Ocean. Indeed, there is an intermodel relationship between these two aspects (Kajtar et al. 2018; McGregor et al. 2018), highlighting the need to account for other basins in studying Pacific decadal variability.


In addition, using an eddy-permitting ocean model, Maher et al. (2018) showed that the recent Pacific trade wind acceleration can explain heat content trends in the Pacific Ocean, and in the Indian Ocean via the Indonesian Throughflow. They also showed that these heat content anomalies do not entirely dissipate with the abatement of the Pacific winds, leaving residual heat in the ocean. The impact of low-frequency subduction of heat on ENSO processes, particularly over decadal time scales, warrants further investigation.

**SUMMARY AND CLOSING REMARKS.** The following key issues were raised during the workshop:

- *Characteristics of ENSO extremes:* The 2015/16 El Niño demonstrated greater potential diversity in extreme El Niño events than previously realized. It highlights the need for a better understanding of the causes and predictability of extreme ENSO events, supported by high-quality observational data and metrics toward better monitoring, predicting, and analyzing future events.
- *ENSO predictability:* There is scope to improve our understanding of ENSO development and ENSO representation in climate models, toward better operational predictive capability. Other than WWV and WWBs, climate variability and trends outside the tropical Pacific are also important factors. Decadal variability and long-term changes of these factors have important implications for the decadal predictability of ENSO.
- *Response to greenhouse forcing:* There is an indication that extreme ENSO events may become more frequent in a warmer future, and the present global warming may already be great enough to exacerbate the ENSO-induced rainfall disruption in the Pacific. Given existing model biases, there is a need to continually revise these assessments with improved models (e.g., CMIP6).

- *Model bias:* Current state-of-the-art climate models still have problems simulating the detailed nuances of ENSO and its seasonality, and these problems may be linked to the persistent cold tongue bias and its broader effects (e.g., double ITCZ bias), as well as to biases in other basins that can impact ENSO through atmospheric and oceanic teleconnections.
- *Synoptic-scale oceanic processes:* Tropical instability waves, which are not well resolved in climate models, are an underrepresented source of ocean mixing. Our understanding of ENSO irregularity would benefit from quantifying the relative effects of ocean noise and atmospheric stochasticity.
- *Mechanisms for decadal variability and trends:* Further research is needed to clarify the manner in which the IPO and other aspects of decadal variability interact with ENSO, and the extent to which the IPO is attributed to red noise associated with ENSO and other factors including external forcing and the role of remote oceanic basins.

The key elements of our discussions are summarized in Fig. 5, which highlights areas of challenges and the ramifications associated with understanding ENSO dynamics and predictability. While ENSO predictability and the regional impacts can vary across events and decades (e.g., Power et al. 1999;



**FIG. 5. Factors affecting ENSO and the societal implications. ENSO characteristics are influenced by many factors, including coupled feedback processes, atmospheric and oceanic noise, and climate forcing from other oceanic basins, as well as the basic mean state that evolves on long time scales. All of these components interact with one another and are influenced by external forcing (e.g., greenhouse gases, aerosols, solar variability), which in turn influences the predictability and impacts of ENSO.**

Barnston et al. 2012; Karamperidou et al. 2014; Zhao et al. 2016) and the ENSO predictability limit is not yet known (e.g., Newman and Sardeshmukh 2017), improved understanding of the interaction of these processes and their depiction in climate models could ultimately improve seasonal forecasts, decadal predictions, and future projections of ENSO. This should also lead to more accurate identification of ENSO impacts, with longer lead times and potential benefits to improved climate risk management. These are especially important for Australia given the pronounced and complex ENSO impacts on its regional climate (see “Motivation” section), but this should also be applicable to all other ENSO-affected countries. The way forward is to sustain and expand ENSO research through further advances in modeling, observations, theoretical frameworks, forecasts, analysis techniques, and development of new metrics/indices, coordinated across a collaborative international research environment.

Such endeavors have been ongoing in Australia and have been nurtured by various government initiatives, organizations, universities, and industries. Research collaboration and student training across institutions are fostered through, for example, the currently active National Environmental Science Program ([www.environment.gov.au/science/nesp](http://www.environment.gov.au/science/nesp)) and the Australian Research Council (ARC) Centre of Excellence for Climate Extremes (<https://climateextremes.org.au>), both of which integrate various ENSO-related topics within their overarching research programs. The ARC Centre has extended the collaborative network to include several international institutions. ENSO is a core research element in the recently established Centre for Southern Hemisphere Oceans Research (CSHOR; <https://cshor.csiro.au>), which is a partnership between the CSIRO and Qingdao National Laboratory for Marine Science and Technology (QNLN), with the University of New South Wales (UNSW) and the University of Tasmania as partners.

Australia has the National Computational Infrastructure (NCI) that provides high-performance computing and data-intensive services to researchers, supporting model development such as Australia’s national climate model, the Australian Community Climate and Earth System Simulator (ACCESS), that contributes to CMIP and the Intergovernmental Panel on Climate Change (IPCC). The development of ACCESS has involved close international partnerships, particularly with the Met Office (UKMO) in the United Kingdom and the Geophysical Fluid Dynamics Laboratory (GFDL) in the United States.

The seasonal forecast system that is used to produce the ENSO outlook continues to advance at the BOM, with POAMA having been replaced in 2018 by the seasonal forecast version of ACCESS (ACCESS-S), which is of a higher resolution and better than POAMA in distinguishing between EP and CP El Niño events, particularly at increasing lead times (Hudson et al. 2017). Development and improvement of the BOM’s coupled model predictions systems rely heavily on the partnership with the UKMO, but have also been and continue to be strongly supported by various agricultural research and development corporations, with matching funding from the Australian Government through the Managing Climate Variability Program (<http://managingclimate.gov.au>). Some of the key focuses of the BOM’s research directions toward improved seasonal predictions include tackling key model biases that are affecting the simulation and prediction of ENSO diversity, improving data assimilation techniques for more accurate forecast model initialization, extending ENSO prediction lead times, and exploring the potential for multiyear prediction of ENSO.

Observing the tropical Pacific for ENSO, of which the Tropical Atmosphere Ocean (TAO)/Triangle Trans-Ocean Buoy Network (TRITON) mooring arrays ([www.pmel.noaa.gov/gtmba](http://www.pmel.noaa.gov/gtmba)) have been instrumental, is an integral part of international efforts in ocean observations under the auspices of the Global Ocean Observing System (GOOS). While not a contributor to TAO/TRITON, Australia is a major contributor to GOOS through deployment of Argo floats operated by the Integrated Marine Observing System (IMOS; <http://imos.org.au>). The global array of Argo profiling floats provides real-time data of temperature, salinity, and currents up to 2,000-m depth over the global oceans, including the tropical Pacific. The BOM relies on Argo subsurface temperature and salinity data for initializing their seasonal prediction models.

Australian ENSO researchers have also actively participated in international initiatives and organizations, such as the World Climate Research Programme (WCRP), Tropical Pacific Observing System (TPOS) 2020, and IPCC, among many others, that in turn foster and enrich ENSO research activities. The future success of ENSO research and prediction in Australia will certainly benefit from sustained cross-institutional synergies in a conducive environment of the multinational network.

**ACKNOWLEDGMENTS.** The authors thank the Australian Research Council (ARC) Centre of Excellence

for Climate Extremes for supporting the workshop. The workshop stemmed in part from discussions at a meeting of the National Environmental Science Program (NESP) Earth System and Climate Change Hub Project 2.2 science team. Andrew Wittenberg and two anonymous reviewers provided constructive comments and suggestions that improved the manuscript. The writing of this essay is partly supported by NESP, the ARC, and the Centre for Southern Hemisphere Oceans Research (CSHOR), a joint research center for Southern Hemisphere ocean research between QNLM and CSIRO. This paper contributes to the goals of CLIVAR of WCRP and the International Commission on Climate of IAMAS/IUGG.

## REFERENCES

- Abellán, E., S. McGregor, M. H. England, and A. Santoso, 2018: Distinctive role of ocean advection anomalies in the development of the extreme 2015–16 El Niño. *Climate Dyn.*, **51**, 2191–2208, <https://doi.org/10.1007/s00382-017-4007-0>.
- An, S., and F.-F. Jin, 2004: Nonlinearity and asymmetry of ENSO. *J. Climate*, **17**, 2399–2412, [https://doi.org/10.1175/1520-0442\(2004\)017<2399:NAOE>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2399:NAOE>2.0.CO;2).
- Ashok, K., S. K. Behera, S. A. Rao, H. Weng, and T. Yamagata, 2007: El Niño Modoki and its possible teleconnection. *J. Geophys. Res.*, **112**, C11007, <https://doi.org/10.1029/2006JC003798>.
- Barnston, A. G., M. K. Tippett, M. L. L'Heureux, S. Li, and D. G. DeWitt, 2012: Skill of real-time seasonal ENSO model predictions during 2002–11: Is our capability increasing? *Bull. Amer. Meteor. Soc.*, **93**, 631–651, <https://doi.org/10.1175/BAMS-D-11-00111.1>.
- Bjerknes, J., 1969: Atmospheric teleconnections from the equatorial Pacific. *Mon. Wea. Rev.*, **97**, 163–172, [https://doi.org/10.1175/1520-0493\(1969\)097<0163:ATFTEP>2.3.CO;2](https://doi.org/10.1175/1520-0493(1969)097<0163:ATFTEP>2.3.CO;2).
- Blunden, J., and D. S. Arndt, Eds., 2016: State of the Climate in 2016. *Bull. Amer. Meteor. Soc.*, **98** (8), S1–S280, <https://doi.org/10.1175/2017BAMSStateoftheClimate.1>.
- Bonfils, C. J., B. D. Santer, T. J. Phillips, K. Marvel, L. R. Leung, C. Doutriaux, and A. Capotondi, 2015: Relative contributions of mean-state shifts and ENSO-driven variability to precipitation changes in a warming climate. *J. Climate*, **28**, 9997–102013, <https://doi.org/10.1175/JCLI-D-15-0341.1>.
- Cai, W., P. van Rensch, T. Cowan, and H. H. Hendon, 2011: Teleconnection pathways of ENSO and the IOD and the mechanisms for impacts on Australian rainfall. *J. Climate*, **24**, 3910–3923, <https://doi.org/10.1175/2011JCLI4129.1>.
- , and Coauthors, 2014: Increasing frequency of extreme El Niño events due to greenhouse warming. *Nat. Climate Change*, **4**, 111–116, <https://doi.org/10.1038/nclimate2100>.
- , and Coauthors, 2015a: ENSO and greenhouse warming. *Nat. Climate Change*, **5**, 849–859, <https://doi.org/10.1038/nclimate2743>.
- , and Coauthors, 2015b: Increased frequency of extreme La Niña events under greenhouse warming. *Nat. Climate Change*, **5**, 132–137, <https://doi.org/10.1038/nclimate2492>.
- , G. Wang, A. Santoso, X. Lin, and L. Wu, 2017: Definition of extreme El Niño and its impact on projected increase in extreme El Niño frequency. *Geophys. Res. Lett.*, **44**, 11 184–11 190, <https://doi.org/10.1002/2017GL075635>.
- Capotondi, A., and Coauthors, 2015a: Understanding ENSO diversity. *Bull. Amer. Meteor. Soc.*, **96**, 921–938, <https://doi.org/10.1175/BAMS-D-13-00117.1>.
- , Y.-G. Ham, A. Wittenberg, and J.-S. Kug, 2015b: Climate model biases and El Niño Southern Oscillation (ENSO) simulation. *US CLIVAR Variations*, Vol. 13, No. 1, US CLIVAR Program, Washington, DC, 21–25, [https://usclivar.org/sites/default/files/documents/2015/Variations2015Winter\\_0.pdf](https://usclivar.org/sites/default/files/documents/2015/Variations2015Winter_0.pdf).
- Chen, C., M.A. Cane, A.T. Wittenberg, and D. Chen, 2017: ENSO in the CMIP5 simulations: Life cycles, diversity, and responses to climate change. *J. Climate*, **30**, 775–801, <https://doi.org/10.1175/JCLI-D-15-0901.1>.
- Chung, C. Y. Y., and S. B. Power, 2017: The non-linear impact of El Niño, La Niña and the Southern Oscillation on seasonal and regional Australian precipitation. *J. South. Hemisphere Earth Syst. Sci.*, **67**, 25–45, <https://doi.org/10.22499/3.6701.003>.
- , —, J. M. Arblaster, H. A. Rashid, and G. L. Roff, 2014: Nonlinear precipitation response to El Niño and global warming in the Indo-Pacific. *Climate Dyn.*, **42**, 1837–1856, <https://doi.org/10.1007/s00382-013-1892-8>.
- Collins, M., and Coauthors, 2010: The impact of global warming on the tropical Pacific Ocean and El Niño. *Nat. Geosci.*, **3**, 391–397, <https://doi.org/10.1038/ngeo868>.
- DiNezio, P., B. P. Kirtman, A. C. Clement, S. Lee, G. A. Vecchi, and A. Wittenberg, 2012: Mean climate controls on the simulated response of ENSO to increasing greenhouse gases. *J. Climate*, **25**, 7399–7420, <https://doi.org/10.1175/JCLI-D-11-00494.1>.
- Dommengat, D., T. Bayr, and C. Frauen, 2013: Analysis of the non-linearity in the pattern and time evolution of El Niño Southern Oscillation. *Climate Dyn.*, **40**, 2825–2847, <https://doi.org/10.1007/s00382-012-1475-0>.

- Dong, L., and M. J. McPhaden, 2018: Unusually warm Indian Ocean sea surface temperatures help to arrest development of El Niño in 2014. *Sci. Rep.*, **8**, 2249, <https://doi.org/10.1038/s41598-018-20294-4>.
- Drosowsky, W., and L. E. Chambers, 2001: Near-global sea surface temperature anomalies as predictors of Australian seasonal rainfall. *J. Climate*, **14**, 1677–1687, [https://doi.org/10.1175/1520-0442\(2001\)014<1677:NA CNGS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<1677:NA CNGS>2.0.CO;2).
- England, M. H., and Coauthors, 2014: Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus. *Nat. Climate Change*, **4**, 222–227, <https://doi.org/10.1038/nclimate2106>.
- Graham, F. S., J. N. Brown, A. T. Wittenberg, S. J. Marsland, and N. J. Holbrook, 2017: Understanding the double peaked El Niño in coupled GCMs. *Climate Dyn.*, **48**, 2045–2063, <https://doi.org/10.1007/s00382-016-3189-1>.
- Guilyardi, E., A. Wittenberg, M. Balmaseda, W. Cai, M. Collins, M. J. McPhaden, M. Watanabe, and S. Yeh, 2016: Fourth CLIVAR Workshop on the Evaluation of ENSO Processes in Climate Models: ENSO in a changing climate. *Bull. Amer. Meteor. Soc.*, **97**, 817–820, <https://doi.org/10.1175/BAMS-D-15-00287.1>.
- Ham, Y.-G., and I.-S. Kang, 2011: Improvement of seasonal forecasts with inclusion of tropical instability waves on initial conditions. *Climate Dyn.*, **36**, 1277–1290, <https://doi.org/10.1007/s00382-010-0743-0>.
- Hannam, P., 2014: ‘Most-watched’ El Niño gathers pace in the Pacific. *Sydney Morning Herald*, 17 June, [www.smh.com.au/environment/weather/mostwatched-el-nino-gathers-pace-in-pacific-20140617-zsaw6.html](http://www.smh.com.au/environment/weather/mostwatched-el-nino-gathers-pace-in-pacific-20140617-zsaw6.html).
- Hendon, H., D. Thompson, and M. Wheeler, 2007: Australian rainfall and surface temperature variations associated with the southern annular mode. *J. Climate*, **20**, 2452–2467, <https://doi.org/10.1175/JCLI4134.1>.
- , E.-P. Lim, G. Wang, O. Alves, and D. Hudson, 2009: Prospects for predicting two flavors of El Niño. *Geophys. Res. Lett.*, **36**, L19713, <https://doi.org/10.1029/2009GL040100>.
- , —, and G. Liu, 2012: The role of air–sea interaction for prediction of Australian summer monsoon rainfall. *J. Climate*, **25**, 1278–1290, <https://doi.org/10.1175/JCLI-D-11-00125.1>.
- Henley, B., J. Gergis, D. J. Karoly, S. Power, J. Kennedy, and C. K. Folland, 2015: A tripole index for the Interdecadal Pacific Oscillation. *Climate Dyn.*, **45**, 3077–3090, <https://doi.org/10.1007/s00382-015-2525-1>.
- Holbrook, N. J., J. Davidson, M. Feng, A. J. Hobday, J. M. Lough, S. McGregor, and J. S. Risbey, 2009: El Niño–Southern Oscillation. Report card of marine climate change for Australia: Detailed scientific assessment, E. S. Poloczanska, A. J. Hobday, and A. J. Richardson, Eds., NCCARF Publ. 05/09, 29–51.
- , and Coauthors, 2014: Decadal climate variability and cross-scale interactions: ICCL 2013 Expert Assessment Workshop. *Bull. Amer. Meteor. Soc.*, **95**, ES155–ES158, <https://doi.org/10.1175/BAMS-D-13-00201.1>.
- Holmes, R. M., S. McGregor, A. Santoso, and M. H. England, 2018: Contribution of tropical instability waves to ENSO irregularity. *Climate Dyn.*, <https://doi.org/10.1007/s00382-018-4217-0>.
- , J. D. Zika, and M. H. England, 2019: Diathermal heat transport in a global ocean model. *J. Phys. Oceanogr.*, **49**, 141–161, <https://doi.org/10.1175/JPO-D-18-0098.1>.
- Hope, P., B. J. Henley, J. Gergis, J. Brown, and H. Ye, 2017: Time-varying spectral characteristics of ENSO over the Last Millennium. *Climate Dyn.*, **49**, 1705–1727, <https://doi.org/10.1007/s00382-016-3393-z>.
- Hu, S., and A. V. Fedorov, 2016: Exceptionally strong easterly wind burst stalling El Niño of 2014. *Proc. Natl. Acad. Sci. USA*, **113**, 2005–2010, <https://doi.org/10.1073/pnas.1514182113>.
- Huang, B., and Coauthors, 2017: Extended reconstructed sea surface temperature version 5 (ERSSTv5): Upgrades, validations, and intercomparisons. *J. Climate*, **30**, 8179–8205, <https://doi.org/10.1175/JCLI-D-16-0836.1>.
- Hudson, D., and Coauthors, 2017: ACCESS-S1: The new Bureau of Meteorology multi-week to seasonal prediction system. *J. South. Hemisphere Earth Syst. Sci.*, **67**, 132–159, [www.bom.gov.au/jshess/docs/2017/Hudson.pdf](http://www.bom.gov.au/jshess/docs/2017/Hudson.pdf).
- Ishii, M., A. Shouji, S. Sugimoto, and T. Matsumoto, 2005: Objective analyses of sea-surface temperature and marine meteorological variables for the 20th century using ICOADS and the Kobe collection. *Int. J. Climatol.*, **25**, 865–879, <https://doi.org/10.1002/joc.1169>.
- Jin, F.-F., 1997: An equatorial ocean recharge paradigm for ENSO. Part I: Conceptual model. *J. Atmos. Sci.*, **54**, 811–829, [https://doi.org/10.1175/1520-0469\(1997\)054<0811:AEORPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<0811:AEORPF>2.0.CO;2).
- Kajtar, J. B., A. Santoso, S. McGregor, M. H. England, and Z. Baille, 2018: Model under-representation of decadal Pacific trade wind trends and its link to tropical Atlantic bias. *Climate Dyn.*, **50**, 1471–1484, <https://doi.org/10.1007/s00382-017-3699-5>.
- Kalnay, E., and Coauthors, 1996: The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Amer. Meteor. Soc.*, **77**, 437–471, [https://doi.org/10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2).

- Kao, H.-Y., and J.-Y. Yu, 2009: Contrasting eastern Pacific and central Pacific types of El Niño. *J. Climate*, **22**, 615–632, <https://doi.org/10.1175/2008JCLI2309.1>.
- Karamperidou, C., M. A. Cane, U. Lall, and A. T. Wittenberg, 2014: Intrinsic modulation of ENSO predictability viewed through a local Lyapunov lens. *Climate Dyn.*, **42**, 253–270, <https://doi.org/10.1007/s00382-013-1759-z>.
- Kim, S.-T., W. Cai, F.-F. Jin, A. Santoso, L. Wu, E. Guilyardi, and S.-I. An, 2014a: Response of El Niño sea surface temperature variability to greenhouse warming. *Nat. Climate Change*, **4**, 786–790, <https://doi.org/10.1038/nclimate2326>.
- , —, —, and J.-Y. Yu, 2014b: ENSO stability in coupled climate models and its association with mean state. *Climate Dyn.*, **42**, 3313–3321, <https://doi.org/10.1007/s00382-013-1833-6>.
- Kociuba, G., and S. B. Power, 2015: Inability of CMIP5 models to simulate recent strengthening of the Walker circulation: Implications for projections. *J. Climate*, **28**, 20–35, <https://doi.org/10.1175/JCLI-D-13-00752.1>.
- Kohyama, T., D. L. Hartmann, and D. S. Battisti, 2017: La Niña-like mean-state response to global warming and potential oceanic roles. *J. Climate*, **30**, 4207–4225, <https://doi.org/10.1175/JCLI-D-16-0441.1>.
- Lee, S.-K., P. N. DiNezio, E.-S. Chung, S.-W. Yeh, A. T. Wittenberg, and C. Wang, 2014: Spring persistence, transition and resurgence of El Niño. *Geophys. Res. Lett.*, **41**, 8578–8585, <https://doi.org/10.1002/2014GL062484>.
- Levine, A. F. Z., and M. J. McPhaden, 2016: How the July 2014 easterly wind burst gave the 2015–2016 El Niño a head start. *Geophys. Res. Lett.*, **43**, 6503–6510, <https://doi.org/10.1002/2016GL069204>.
- L’Heureux, M. L., and Coauthors, 2017: Observing and predicting the 2015/16 El Niño. *Bull. Amer. Meteor. Soc.*, **98**, 1363–1382, <https://doi.org/10.1175/BAMS-D-16-0009.1>.
- Lim, E.-P., and H. H. Hendon, 2015: Understanding the contrast of Australian springtime rainfall of 1997 and 2002 in the frame of two flavors of El Niño. *J. Climate*, **28**, 2804–2822, <https://doi.org/10.1175/JCLI-D-14-00582.1>.
- , and —, 2017: Causes and predictability of the negative Indian Ocean Dipole and its impact on La Niña during 2016. *Sci. Rep.*, **7**, 12619, <https://doi.org/10.1038/s41598-017-12674-z>.
- , —, J. M. Arblaster, C. Chung, A. F. Moise, P. Hope, G. Young, and M. Zhao, 2016: Interaction of the recent 50 year SST trend and La Niña 2010: Amplification of the Southern Annular Mode and Australian springtime rainfall. *Climate Dyn.*, **47**, 2273–2291, <https://doi.org/10.1007/s00382-015-2963-9>.
- Loughran, T. F., S. E. Perkins-Kirkpatrick, and L. V. Alexander, 2017: Understanding the spatio-temporal influence of climate variability on Australian heatwaves. *Int. J. Climatol.*, **37**, 3963–3975, <https://doi.org/10.1002/joc.4971>.
- Lu, L., and D. Hedley, 2004: The impact of the 2002 drought on the economy and agricultural employment. Economic Roundup Autumn 2004, Australian Government, <https://treasury.gov.au/publication/economic-roundup-autumn-2004/the-impact-of-the-2002-drought-on-the-economy-and-agricultural-employment/>.
- Luo, J.-J., S. Masson, E. Roeckner, G. Madec, and T. Yamagata, 2005: Reducing climatology bias in an ocean–atmosphere CGCM with improved coupling physics. *J. Climate*, **18**, 2344–2360, <https://doi.org/10.1175/JCLI3404.1>.
- , G. Liu, H. Hendon, O. Alves, and T. Yamagata, 2017: Inter-basin sources for two-year predictability of the multi-year La Niña event in 2010–2012. *Sci. Rep.*, **7**, 2276, <https://doi.org/10.1038/s41598-017-01479-9>.
- , G. Wang, and D. Dommenges, 2018: May common model biases reduce CMIP5’s ability to simulate the recent Pacific La Niña-like cooling? *Climate Dyn.*, **50**, 1335, <https://doi.org/10.1007/s00382-017-3688-8>.
- Maher, N., M. H. England, A. Sen Gupta, and P. Spence, 2018: Role of Pacific trade winds in driving ocean temperatures during the recent slowdown and projections under a wind trend reversal. *Climate Dyn.*, **51**, 321–336, <https://doi.org/10.1007/s00382-017-3923-3>.
- McBride, J. L., and N. Nicholls, 1983: Seasonal relationship between Australian rainfall and Southern Oscillation. *Mon. Wea. Rev.*, **111**, 1998–2004, [https://doi.org/10.1175/1520-0493\(1983\)111<1998:SRBARA>2.0.CO;2](https://doi.org/10.1175/1520-0493(1983)111<1998:SRBARA>2.0.CO;2).
- McGregor, S., A. Timmermann, M. F. Stuecker, M. H. England, M. Merrifield, F.-F. Jin, and Y. Chikamoto, 2014: Recent Walker circulation strengthening and Pacific cooling amplified by Atlantic warming. *Nat. Climate Change*, **4**, 888–892, <https://doi.org/10.1038/nclimate2330>.
- , M. F. Stuecker, J. B. Kajtar, M. H. England, and M. Collins, 2018: Model tropical Atlantic biases underpin diminished Pacific decadal variability. *Nat. Climate Change*, **8**, 493–498, <https://doi.org/10.1038/s41558-018-0163-4>.
- McPhaden, M. J., 2015: Playing hide and seek with El Niño. *Nat. Climate Change*, **5**, 791–795, <https://doi.org/10.1038/nclimate2775>.
- Meehl, G. A., A. Hu, and H. Teng, 2016: Initialized decadal prediction for transition to positive phase of

- the Interdecadal Pacific Oscillation. *Nature Commun.*, **7**, 11718, <https://doi.org/10.1038/ncomms11718>.
- Meinen, C. S., and M. J. McPhaden, 2000: Observations of warm water volume changes in the equatorial Pacific and their relationship to El Niño and La Niña. *J. Climate*, **13**, 3551–3559, [https://doi.org/10.1175/1520-0442\(2000\)013<3551:OOWWVC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<3551:OOWWVC>2.0.CO;2).
- Menkes, C., J. G. Vialard, S. C. Kennan, J. Boulanger, and G. V. Madec, 2006: A modeling study of the impact of tropical instability waves on the heat budget of the eastern equatorial Pacific. *J. Phys. Oceanogr.*, **36**, 847–865, <https://doi.org/10.1175/JPO2904.1>.
- , M. Lengaigne, J. Vialard, M. Puy, P. Marchesiello, S. Cravatte, and G. Cambon, 2014: About the role of the westerly wind events in the possible development of an El Niño in 2014. *Geophys. Res. Lett.*, **41**, 6476–6483, <https://doi.org/10.1002/2014GL061186>.
- Meyers, G., P. McIntosh, L. Pigot, and M. Pook, 2007: The years of El Niño, La Niña, and interactions with the tropical Indian Ocean. *J. Climate*, **20**, 2872–2880, <https://doi.org/10.1175/JCLI4152.1>.
- Neske, S., and S. McGregor, 2018: Understanding the warm water volume precursor of ENSO events and its interdecadal variation. *Geophys. Res. Lett.*, **45**, 1577–1585, <https://doi.org/10.1002/2017GL076439>.
- Newman, M., and P. D. Sardeshmukh, 2017: Are we near the predictability limit of tropical Indo-Pacific sea surface temperatures? *Geophys. Res. Lett.*, **44**, 8520–8529, <https://doi.org/10.1002/2017GL074088>.
- , S.-I. Shin, and M. A. Alexander, 2011: Natural variation in ENSO flavors. *Geophys. Res. Lett.*, **38**, L14705, <https://doi.org/10.1029/2011GL047658>.
- , and Coauthors, 2016: The Pacific decadal oscillation, revisited. *J. Climate*, **29**, 4399–4427, <https://doi.org/10.1175/JCLI-D-15-0508.1>.
- , A. T. Wittenberg, L. Cheng, G. P. Compo, and C. A. Smith, 2018: The extreme 2015/16 El Niño, in the context of historical climate variability and change [in “Explaining Extreme Events of 2016 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **99**, S16–S20, <https://doi.org/10.1175/BAMS-D-17-0116.1>.
- Nicholls, N., 1984: The Southern Oscillation and Indonesian sea surface temperature. *Mon. Wea. Rev.*, **112**, 424–432, [https://doi.org/10.1175/1520-0493\(1984\)112<0424:TSAIS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1984)112<0424:TSAIS>2.0.CO;2).
- , 1985: Impact of the Southern Oscillation on Australian crops. *J. Climatol.*, **5**, 553–560, <https://doi.org/10.1002/joc.3370050508>.
- , 1991: The El Niño/Southern Oscillation and Australian vegetation. *Vegetatio*, **91**, 23–36, <https://doi.org/10.1007/BF00036045>.
- Norman, F. I., and N. Nicholls, 1991: The Southern Oscillation and variations in waterfowl abundance in southeastern Australia. *Aust. J. Ecol.*, **16**, 485–490, <https://doi.org/10.1111/j.1442-9993.1991.tb01077.x>.
- Ogata, T., S. Xie, A. Wittenberg, and D. Sun, 2013: Interdecadal amplitude modulation of El Niño–Southern Oscillation and its impacts on tropical Pacific decadal variability. *J. Climate*, **26**, 7280–7297, <https://doi.org/10.1175/JCLI-D-12-00415.1>.
- Pacanowski, R., 1987: Effect of equatorial currents on surface stress. *J. Phys. Oceanogr.*, **17**, 833–838, [https://doi.org/10.1175/1520-0485\(1987\)017<0833:EOECOS>2.0.CO;2](https://doi.org/10.1175/1520-0485(1987)017<0833:EOECOS>2.0.CO;2).
- Paek, H., J. Y. Yu, and C. Qian, 2017: Why were the 2015/2016 and 1997/1998 extreme El Niños different? *Geophys. Res. Lett.*, **44**, 1848–1856, <https://doi.org/10.1002/2016GL071515>.
- Power, S., and R. Colman, 2006: Multi-year predictability in a coupled general circulation model. *Climate Dyn.*, **26**, 247–272, <https://doi.org/10.1007/s00382-005-0055-y>.
- , and I. N. Smith, 2007: Weakening of the Walker Circulation and apparent dominance of El Niño both reach record levels, but has ENSO really changed? *Geophys. Res. Lett.*, **34**, L18702, <https://doi.org/10.1029/2007GL030854>.
- , and J. Callaghan, 2016: Variability in severe coastal flooding, associated storms, and death tolls in southeastern Australia since the mid-nineteenth century. *J. Appl. Meteor. Climatol.*, **55**, 1139–1149, <https://doi.org/10.1175/JAMC-D-15-0146.1>.
- , and F. P. D. Delage, 2018: El Niño–Southern Oscillation and associated climatic conditions around the world during the latter half of the twenty-first century. *J. Climate*, **31**, 6189–6207, <https://doi.org/10.1175/JCLI-D-18-0138.1>.
- , F. Tseitkin, S. Torok, B. Lavery, and B. McAvaney, 1998: Australian temperature, Australian rainfall, and the Southern Oscillation, 1910–1996: Coherent variability and recent changes. *Aust. Meteor. Mag.*, **47**, 85–101.
- , C. Folland, A. Colman, and V. Mehta, 1999: Inter-decadal modulation of the impact of ENSO on Australia. *Climate Dyn.*, **15**, 319–324, <https://doi.org/10.1007/s003820050284>.
- , M. Haylock, R. Colman, and X. Wang, 2006: The predictability of interdecadal changes in ENSO and ENSO teleconnections. *J. Climate*, **19**, 4755–4771, <https://doi.org/10.1175/JCLI3868.1>.
- , F. Delage, C. Chung, G. Kociuba, and K. Keay, 2013: Robust twenty-first-century projections of El Niño and related precipitation variability. *Nature*, **502**, 541–545, <https://doi.org/10.1038/nature12580>.
- , —, —, H. Ye, and B. F. Murphy, 2017a: Humans have already increased the risk of major



- disruptions to Pacific rainfall. *Nat. Commun.*, **8**, 14368, <https://doi.org/10.1038/ncomms14368>.
- , —, G. Wang, I. Smith, and G. Kociuba, 2017b: Apparent limitations in the ability of CMIP5 climate models to simulate recent multi-decadal change in surface temperature: Implications for global temperature projections. *Climate Dyn.*, **49**, 53–69, <https://doi.org/10.1007/s00382-016-3326-x>.
- Pui, A., A. Sharma, A. Santoso, and S. Westra, 2012: Impact of the El Niño–Southern Oscillation, Indian Ocean dipole, and southern annular mode on daily to subdaily rainfall characteristics in east Australia. *Mon. Wea. Rev.*, **140**, 1665–1682, <https://doi.org/10.1175/MWR-D-11-00238.1>.
- Puy, M., J. Vialard, M. Lengaigne, and E. Guilyardi, 2016: Modulation of equatorial Pacific westerly/easterly wind events by the Madden–Julian oscillation and convectively-coupled Rossby waves. *Climate Dyn.*, **46**, 2155–2178, <https://doi.org/10.1007/s00382-015-2695-x>.
- Rashid, H. A., and A. C. Hirst, 2016: Investigating the mechanisms of seasonal ENSO phase locking bias in the ACCESS Coupled Model. *Climate Dyn.*, **46**, 1075–1090, <https://doi.org/10.1007/s00382-015-2633-y>.
- , —, and S. Marsland, 2016: An atmospheric mechanism for ENSO amplitude changes under an abrupt quadrupling of CO<sub>2</sub> concentration in CMIP5 models. *Geophys. Res. Lett.*, **43**, 1687–1694, <https://doi.org/10.1002/2015GL066768>.
- Risbey, J. S., M. J. Pook, P. C. McIntosh, M. C. Wheeler, and H. H. Hendon, 2009: On the remote drivers of rainfall variability in Australia. *Mon. Wea. Rev.*, **137**, 3233–3253, <https://doi.org/10.1175/2009MWR2861.1>.
- Ropelewski, C. F., and M. S. Halpert, 1987: Global and regional scale precipitation patterns associated with the El Niño/Southern Oscillation. *Mon. Wea. Rev.*, **115**, 1606–1626, [https://doi.org/10.1175/1520-0493\(1987\)115<1606:GARSPP>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<1606:GARSPP>2.0.CO;2).
- Saji, N. H., B. N. Goswami, P. N. Vinayachandran, and T. Yamagata, 1999: A dipole mode in the tropical Indian Ocean. *Nature*, **401**, 360–363, <https://doi.org/10.1038/43854>.
- Santoso, A., S. McGregor, F.-F. Jin, W. Cai, M. H. England, S.-I. An, M. McPhaden, and E. Guilyardi, 2013: Late-twentieth-century emergence of the El Niño propagation asymmetry and future projections. *Nature*, **504**, 126–130, <https://doi.org/10.1038/nature12683>.
- , and Coauthors, 2015: ENSO extremes and diversity: Dynamics, teleconnections, and impacts. *Bull. Amer. Meteor. Soc.*, **96**, 1969–1972, <https://doi.org/10.1175/BAMS-D-15-00141.1>.
- , M. J. McPhaden, and W. Cai, 2017: The defining characteristics of ENSO extremes and the strong 2015/2016 El Niño. *Rev. Geophys.*, **55**, 1079–1129, <https://doi.org/10.1002/2017RG000560>.
- Sasaki, W., K. J. Richards, and J.-J. Luo, 2013: Impact of vertical mixing induced by small vertical scale structures above and within the equatorial thermocline on the tropical Pacific in a CGCM. *Climate Dyn.*, **41**, 443–453, <https://doi.org/10.1007/s00382-012-1593-8>.
- Sullivan, A., J.-J. Luo, A. C. Hirst, D. Bi, W. Cai, and J. He, 2016: Robust contribution of decadal anomalies to the frequency of central-Pacific El Niño. *Sci. Rep.*, **6**, 38540, <https://doi.org/10.1038/srep38540>.
- Takahashi, K., and B. Dewitte, 2016: Strong and moderate nonlinear El Niño regimes. *Climate Dyn.*, **46**, 1627–1645, <https://doi.org/10.1007/s00382-015-2665-3>.
- , A. Montecinos, K. Goubanova, and B. Dewitte, 2011: ENSO regimes: Reinterpreting the canonical and Modoki El Niño. *Geophys. Res. Lett.*, **38**, L10704, <https://doi.org/10.1029/2011GL047364>.
- Taschetto, A. S., and M. H. England, 2009: El Niño Modoki impacts on Australian rainfall. *J. Climate*, **22**, 3167–3174, <https://doi.org/10.1175/2008JCLI2589.1>.
- , A. Sen Gupta, H. H. Hendon, C. C. Ummenhofer, and M. H. England, 2011: The contribution of Indian Ocean sea surface temperature anomalies on Australian summer rainfall during El Niño events. *J. Climate*, **24**, 3734–3747, <https://doi.org/10.1175/2011JCLI3885.1>.
- , —, N. C. Jourdain, A. Santoso, C. C. Ummenhofer, and M. H. England, 2014: Cold tongue and warm pool ENSO events in CMIP5: Mean state and future projections. *J. Climate*, **27**, 2861–2885, <https://doi.org/10.1175/JCLI-D-13-00437.1>.
- Thompson, D. W. J., and J. M. Wallace, 2000: Annular modes in the extratropical circulation. Part I: Month-to-month variability. *J. Climate*, **13**, 1000–1016, [https://doi.org/10.1175/1520-0442\(2000\)013<1000:AMITEC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<1000:AMITEC>2.0.CO;2).
- Timmermann, A., and Coauthors, 2018: El Niño–Southern Oscillation complexity. *Nature*, **559**, 535–545, <https://doi.org/10.1038/s41586-018-0252-6>.
- Trenberth, K. E., and D. P. Stepaniak, 2001: Indices of El Niño evolution. *J. Climate*, **14**, 1697–1701, [https://doi.org/10.1175/1520-0442\(2001\)014<1697:LIOENO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<1697:LIOENO>2.0.CO;2).
- Ummenhofer, C. C., A. Sen Gupta, M. H. England, A. S. Taschetto, P. R. Briggs, and M. R. Raupach, 2015: How did ocean warming affect Australian rainfall extremes during the 2010/2011 La Niña event? *Geophys. Res. Lett.*, **42**, 9942–9951, <https://doi.org/10.1002/2015GL065948>.
- van Rensch, P., A. J. E. Gallant, W. Cai, and N. Nicholls, 2015: Evidence of local sea surface temperatures

- overriding the southeast Australian rainfall response to the 1997–1998 El Niño. *Geophys. Res. Lett.*, **42**, 9449–9456, <https://doi.org/10.1002/2015GL066319>.
- Vecchi, G. A., and D. E. Harrison, 2000: Tropical Pacific sea surface temperature anomalies, El Niño, and equatorial westerly wind events. *J. Climate*, **13**, 1814–1830, [https://doi.org/10.1175/1520-0442\(2000\)013<1814:TPSSTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<1814:TPSSTA>2.0.CO;2).
- , B. J. Soden, A. T. Wittenberg, I. M. Held, A. Leetmaa, and M. J. Harrison, 2006: Weakening of tropical Pacific atmospheric circulation due to anthropogenic forcing. *Nature*, **441**, 73–76, <https://doi.org/10.1038/nature04744>.
- Vijayeta, A., and D. Dommenges, 2018: An evaluation of ENSO dynamics in CMIP simulations in the framework of the recharge oscillator model. *Climate Dyn.*, **51**, 1753–1771, <https://doi.org/10.1007/s00382-017-3981-6>.
- Voice, M. E., and F. J. Gauntlett, 1984: The 1983 Ash Wednesday fires in Australia. *Mon. Wea. Rev.*, **112**, 584–590, [https://doi.org/10.1175/1520-0493\(1984\)112<0584:TAWFIA>2.0.CO;2](https://doi.org/10.1175/1520-0493(1984)112<0584:TAWFIA>2.0.CO;2).
- Wang, G., and H. H. Hendon, 2007: Sensitivity of Australian rainfall to inter-El Niño variations. *J. Climate*, **20**, 4211–4226, <https://doi.org/10.1175/JCLI4228.1>.
- , and —, 2017: Why 2015 was a strong El Niño and 2014 was not. *Geophys. Res. Lett.*, **44**, 8567–8575, <https://doi.org/10.1002/2017GL074244>.
- , S. B. Power, and S. McGree, 2016: Unambiguous warming in the western tropical Pacific primarily caused by anthropogenic forcing. *Int. J. Climatol.*, **36**, 933–944, <https://doi.org/10.1002/joc.4395>.
- , W. Cai, B. Gan, L. Wu, A. Santoso, X. Lin, Z. Chen, and M. McPhaden, 2017: Continued increase of extreme El Niño frequency long after 1.5°C warming stabilization. *Nat. Climate Change*, **7**, 568–572, <https://doi.org/10.1038/nclimate3351>.
- Watkins, A., 2015: BOM: We're calling it, the 2015 El Niño is here. *Conversation*, 12 May, <http://theconversation.com/bom-were-calling-it-the-2015-el-nino-is-here-41598>.
- Webster, P. J., and S. Yang, 1992: Monsoon and ENSO: Selectively interactive systems. *Quart. J. Roy. Meteor. Soc.*, **118**, 877–926, <https://doi.org/10.1002/qj.49711850705>.
- Wengel, C. M., W. Latif, W. Park, J. Harlaß, and T. Bayr, 2018: Seasonal ENSO phase locking in the Kiel Climate Model: The importance of the equatorial cold sea surface temperature bias. *Climate Dyn.*, **50**, 901–919, <https://doi.org/10.1007/s00382-017-3648-3>.
- Werner, A., and N. J. Holbrook, 2011: A Bayesian forecast model of Australian region tropical cyclone formation. *J. Climate*, **24**, 6114–6131, <https://doi.org/10.1175/2011JCLI4231.1>.
- Williams, A. A. J., and D. J. Karoly, 1999: Extreme fire weather in Australia and the impact of the El Niño–Southern Oscillation. *Aust. Meteor. Mag.*, **48**, 15–22.
- Wittenberg, A., 2009: Are historical records sufficient to constrain ENSO simulations? *Geophys. Res. Lett.*, **36**, L12702, <https://doi.org/10.1029/2009GL038710>.
- , 2015: Low-frequency variations of ENSO. *US CLIVAR Variations*, Vol. 13, No. 1, US CLIVAR Program, Washington, DC, 26–31, [https://usclivar.org/sites/default/files/documents/2015/Variations2015Winter\\_0.pdf](https://usclivar.org/sites/default/files/documents/2015/Variations2015Winter_0.pdf).
- , and Coauthors, 2014: ENSO modulation: Is it decadal predictable? *J. Climate*, **27**, 2667–2681, <https://doi.org/10.1175/JCLI-D-13-00577.1>.
- Xie, P., and P. A. Arkin, 1997: Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs. *Bull. Amer. Meteor. Soc.*, **78**, 2539–2558, [https://doi.org/10.1175/1520-0477\(1997\)078<2539:GPAYMA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<2539:GPAYMA>2.0.CO;2).
- Xue, Y., and A. Kumar, 2017: Evolution of the 2015/16 El Niño and historical perspective since 1979. *Sci. China Earth Sci.*, **60**, 1572–1588, <https://doi.org/10.1007/s11430-016-0106-9>.
- Zhao, M., H. H. Hendon, O. Alves, G. Liu, and G. Wang, 2016: Weakened eastern Pacific El Niño predictability in the early twenty-first century. *J. Climate*, **29**, 6805–6822, <https://doi.org/10.1175/JCLI-D-15-0876.1>.
- Zhu, Y., and R.-H. Zhang, 2018: An Argo-derived background diffusivity parameterization for improved ocean simulations in the tropical Pacific. *Geophys. Res. Lett.*, **45**, 1509–1517, <https://doi.org/10.1002/2017GL076269>.