

1 **Estimating crop area using seasonal time series of Enhanced Vegetation Index from**
2 **MODIS satellite imagery**

3
4 **A B Potgieter¹, A Apan², P Dunn³ and G Hammer⁴**

5
6 ¹Emerging Technologies, Queensland Department of Primary Industries & Fisheries, Toowoomba QLD
7 4350, Australia

8 ²Australian Centre for Sustainable Catchments & Faculty of Engineering and Surveying University of
9 Southern, Toowoomba QLD 4350, Australia

10 ³ Australian Centre for Sustainable Catchments & Faculty of Sciences, University of Southern
11 Queensland, Toowoomba QLD 4350, Australia

12 ⁴School of Land and Food Sciences, The University of Queensland, Brisbane QLD 4072, Australia

13
14
15 **KEYWORDS:** multi-temporal MODIS, harmonic analysis, principal component analysis

16
17 **Abstract**

18 Cereal grain is one of the main export commodities of Australian agriculture. Over the past decade,
19 crop yield forecasts for wheat and sorghum have shown appreciable utility for industry planning at
20 shire, state and national scales. There is now an increasing drive from industry for more accurate and
21 cost effective crop production forecasts. In order to generate production estimates, accurate crop
22 area estimates are needed by the end of the cropping season. A range of multivariate methods for
23 analysing remotely sensed Enhanced Vegetation Index (EVI) from 16-day Moderate Resolution
24 Imaging Spectroradiometer (MODIS) satellite imagery within the cropping period (i.e. April to
25 November) were investigated to estimate crop area for wheat, barley, chickpea and total winter
26 cropped area for a case study region in NE Australia. Each pixel classification method was trained on
27 ground truth data collected from the study region. Three approaches to pixel classification were

1 examined: (i) cluster analysis of trajectories of EVI values from consecutive multi-date imagery
2 during the crop growth period, (ii) Harmonic Analysis of the Time Series (HANTS) of the EVI values,
3 and (iii) Principal Component Analysis (PCA) of the time series of EVI values. Images classified using
4 these three approaches were compared with each other, and with a classification based on the single
5 MODIS image taken at peak EVI. Imagery for the 2003 and 2004 seasons was used to assess the
6 ability of the methods to determine wheat, barley, chickpea and total cropped area estimates. The
7 accuracy at pixel scale was determined by the percent correct classification metric by contrasting all
8 pixel scale samples with independent pixel observations. At a shire level, aggregated total crop area
9 estimates were compared with surveyed estimates. All multi-temporal methods showed significant
10 overall capability to estimate total winter crop area. There was high accuracy at a pixel scale (>98%
11 correct classification) for identifying overall winter cropping at pixel scale. Discrimination among
12 crops was less accurate, however. Although the use of single-date EVI data produced high accuracy
13 for estimates of wheat area at shire-scale, the result contradicted the poor pixel scale accuracy
14 associated with this approach, due to fortuitous compensating errors. Further studies are needed to
15 extrapolate the multi-temporal approaches to other geographical areas and to improve the lead time
16 for deriving cropped area estimates before harvest.

17

18

1 **Introduction**

2 Cereal grain is one of the main agricultural export commodities of Australia. Grain production,
3 particularly wheat, has increased rapidly during the latter part of the 20th century (Knopke *et al.*
4 2000). This has increased the need of government bodies and industry for crop production forecasts
5 at various spatial and temporal scales. In Australia, variability in cereal production is chiefly affected
6 by climate variability (Nix 1975). This variability can generate significant macro-economic
7 consequences. For example, the severe drought of 2002 reduced the economic growth of the
8 Australian economy by about 0.75 percentage points (Penm 2002). During the last decade,
9 numerous objective information tools have been developed to assist agri-industry in managing this
10 variability at the paddock/farm level (Hammer *et al.* 2001; Nelson *et al.* 2002) and at the regional
11 level (Potgieter *et al.* 2002; Potgieter *et al.* 2005; Stephens *et al.* 2000). Having access to such
12 decision support tools has become increasingly necessary to better deal with production risk in such a
13 highly variable environment.

14 An example of such an objective information tool is the Regional Commodity Forecasting System
15 (RCFS), which is being used operationally by the Queensland Department of Primary Industries &
16 Fisheries (QDPI&F) to predict shire-scale wheat and sorghum yield on a monthly basis
17 (www.dpi.qld.gov.au/fieldcrops). This system, which has operated since 1999, generates a forecast
18 yield distribution for wheat and sorghum on a monthly basis through the cropping season. The
19 system involves the integration of an agro-climatic based simple crop stress index model (Potgieter *et*
20 *al.* 2005; Potgieter *et al.* 2006; Stephens 1998), weather data for the season up to the time of the
21 forecast, and an El Niño Southern Oscillation (ENSO) based seasonal climate forecast system (Stone
22 *et al.* 1996) for the remainder of the season. The RCFS is run each month throughout the crop-
23 growing seasons (winter and summer) for all main crop production shires in Australia. A shortcoming
24 of this system, however, is that it generates only a yield per unit area estimate. To estimate total
25 production, decision-makers must combine this with their subjective knowledge of total area sown (to
26 be harvested) at a spatial scale. Thus, in order to generate total production predictions, a real-time
27 or near real-time estimate of the cropping area is needed throughout the cropping season.

1 Production predictions can be used in updating supply chain information at the regional, state and
2 national levels.

3 Currently, no real-time objective estimates of end of season shire-scale cropped area estimates
4 exist. Although the Australian Bureau of Statistics (ABS) collates an annual shire-scale survey, these
5 data are usually not available until up to 2 years after the survey/census. The use of satellite
6 information, therefore, offers more objectivity, timeliness, repeatability and accuracy. Up to now,
7 however, remote sensing based regional crop production forecasting systems have not become
8 operational at a regional scale mainly because of the high resource costs (i.e. imagery, computer disk
9 space and speed) and the tediousness of applying fine resolution imagery to large areas. With the
10 advent of MODIS imagery, from the satellites launched in Dec 1999 and May 2000 (i.e. the Terra
11 platform, which captures morning images and the Aqua platform, which captures afternoon images,
12 respectively), there is a potential to address the issues of cost and useable pixel size for regional
13 applications.

14 In this study, we examine the use of MODIS imagery to derive specific crop area estimates for
15 agricultural forecasting systems aimed at estimating crop production at a regional scale. Various
16 studies have utilised MODIS in determining land use patterns (Muchoney *et al.* 2000; Price 2003;
17 Zhan *et al.* 2002), vegetation phenology (Zhang *et al.* 2003), and crop (rice) production in the
18 northern Hemisphere (Xiao *et al.* 2005). Near real-time MODIS imagery has also been used in the
19 crop explorer framework developed by the United States Department of Agriculture (USDA), which
20 uses accumulated Normalised Difference Vegetation Index (NDVI) to describe crop conditions relative
21 to a base year (see www.pecad.fas.usda.gov/cropexplorer). This system generates vegetation
22 canopy condition indices at an aggregated continental scale for high level decision makers. Such a
23 non-crop specific (i.e. generalised vegetation canopy condition) approach is likely to have limited
24 value to industry where commodity management decisions need to be made at a much finer spatial
25 resolution (e.g. shire-scale). Currently, no near real-time crop specific area estimates exist for crop
26 specific agricultural systems at a shire-scale in Australia.

27
28 The main objective of this study was to determine the utility of multi-temporal MODIS satellite
29 imagery in estimating area of specific and total winter crops at the end of any specific cropping

1 season. This was achieved by contrasting three multivariate approaches to analyse time series of
2 enhanced vegetation index (EVI) temporal profiles throughout the cropping period. Pixel and shire-
3 scale accuracies for each season studied were assessed based on in-season ground truthing, using
4 data for two selected shires in the Darling Downs region, Queensland, Australia. For each analysis
5 method, pixel classification was trained on ground truth data and accuracy tested on an independent
6 set of ground truth data and on survey data at the aggregate shire-scale.

7

8 **Methods**

9 *Study area*

10 The study area is located in the central Darling Downs region, approximately 150 km west of
11 Brisbane, Queensland, Australia (Figure 1). The Jondaryan and Pittsworth shires (ca 200,000 ha)
12 were selected for this study. The typical crop area planted in both shires equates to nearly half of
13 the total potential cropping area during either winter or summer cropping seasons. Crop
14 management practices are variable, and paddock sizes can range from small (~ 20 ha) to very large
15 (> 400 ha). Some larger paddocks might be divided into cropping strips. These strips can vary in
16 width from 50 m to 180 m in some areas and are usually used in crop rotation practices. The
17 practice of strip cropping was introduced as a preventative measure to counteract the potential loss
18 of topsoil via water runoff and erosion during wet seasons. Soils in this region are generally deep
19 and high in clay content and therefore have very high potential soil water holding capacities. In
20 addition, the high variability in in-crop (i.e. May to October period) rainfall¹, combined with the
21 advantage of deep soils and high soil moisture storing capacity, have shaped crop management
22 practices in the northern region to be more dependent on starting soil moisture at sowing than
23 regions further south in the more winter dominant rainfall areas (Nix 1975). The summer dominant
24 rainfall makes the region highly suited to summer cropping and the soil storage capacity also makes
25 it favourable for winter cropping (e.g. wheat, barley & chickpea) with sowing occurring between
26 middle of April to the end of June. Rotations traditionally incorporate both winter and summer crops.

¹ Coefficients of variation for in-crop (i.e. May to October period) shire rainfall was > 46% for the period 1977 – 2004 with rainfall station data weighted within a shire based on area represented.

1 In these shires, land use patterns over the last 10 years have been dominated by cropping (78% of
2 total shire area in both shires), with total winter crop area planted (which includes wheat and barley)
3 very similar to summer crop area planted (which includes sorghum and cotton)
4 (<http://www.nrm.qld.gov.au/>).

5

6 [Insert Figure 1 here]
7

8 Spatial crop yield variability within a specific season can be caused by either variability in rainfall
9 amount, soil type, crop management practices, timing of rainfall or any combination of these factors.
10 Although variability in rainfall amount might be small across the study area in some years (e.g.
11 2004), there is significant variability in the other factors, constituting a heterogeneous spatial
12 cropping landscape. This was evident in the differences in aggregated shire wheat and barley yields
13 of 2.96 t/ha and 2.69 t/ha for the 2003 season for the Jondaryan and Pittsworth shires, respectively.
14 Differences in aggregated shire wheat and barley yields were less during drier seasons like 2004 with
15 2.52 and 2.5 t/ha for the Jondaryan and Pittsworth shires, respectively (ABARE 2005).

16 *Vegetation Index*

17 The 16-day MODIS Enhanced Vegetation Index (EVI) imagery, which is derived from
18 transformations of the red (620-670 nanometers, 250m pixel size), near-infrared (841-876
19 nanometers, 250m pixel size), and blue (459-479 nanometers, 500m pixel size) spectral bands, was
20 used to form a continuous time series of data that represented the crop growth EVI temporal curve
21 for each pixel in the study area. The MODIS EVI was selected for its insensitivity to atmospheric and
22 canopy soil background noise. In addition, it optimises the vegetation signal with improved
23 sensitivity at higher biomass, which is a significant improvement on the traditional NDVI measure
24 (Huete *et al.* 2002).

25 The EVI is computed as,
26

$$27 \quad EVI = G \frac{\rho_{NIR} - \rho_R}{\rho_{NIR} + C_1 \rho_R - C_2 \rho_B + L} \quad [1]$$

28

1 where ρ is the atmospherically corrected or partially atmospherically corrected (Rayleigh and ozone
2 absorption) surface reflectances, L is the canopy background adjustment that addresses non-linear,
3 differential NIR and red radiant transfer through a canopy, and C1 and C2 are the coefficients of the
4 aerosol resistance term, which uses the blue band to correct for aerosol influences in the red band
5 (Huete *et al.* 2002). The coefficients adopted are L = 1, C1 = 6, C2 = 7.5 and G = 2.5, which
6 represents a gain factor (Huete et al., 1994; Huete et al., 1997). The EVI values thus have an
7 extended sensitivity, which makes it more likely to discriminate between canopy structure differences,
8 such as LAI differences (Justice et al., 1998). The EVI is MODIS specific and is composed based on a
9 high quality EVI values during the 16-day cycle. A filter to the data is applied, which is based on
10 quality, cloud cover and viewing angle in order to create the high quality EVI values (Huete *et al.*
11 2002). The MODIS EVI values range from -2000 to 10000, with a scale factor of 10000, and have a
12 fill value for missing data of -3000. On this scale water bodies have a negative EVI value or close to
13 zero while canopy cover has positive EVI values up to a maximum of 10000 (dense forest canopy).

14

15 *Satellite imagery and re-projection*

16 The "MOD13Q1" MODIS satellite product, which includes the 16-day 250-m VI data, was
17 downloaded from NASA's Earth Observing System (EOS)
18 (<http://edcimswww.cr.usgs.gov/pub/imswelcome/>) web site for the period 2003 to 2004. This
19 resulted in 46 images (i.e. 23 images x 2 years) each of which had a file size of 500 megabytes. The
20 23 images within each season were downloaded for the period January to December. The NDVI and
21 EVI MODIS products were geometrically, atmospherically and bidirectional reflectance distribution
22 fraction (BRDF) corrected, validated and quality assured through the EOS program (Huete *et al.*
23 2002; Justice *et al.* 2002). The MODIS reprojecting tool (<http://edcdaac.usgs.gov/datatools.asp>) was
24 used to sub-sample the "granule" to an area covering the study area. An image was created by
25 stacking the 23 images for each season with a GDA94 projection in ENVI software (RSI, 2005) thus
26 creating a single image with 23 layers. This resulted in a continuous sequence of EVI temporal
27 values for each pixel for each season.

1 Landsat TM 5 images (14 Sept 2004 and 16 Sept 2004), in combination with farm boundaries and
2 the 1999 land use map (Department of Natural Resources and Water 2006) of the study area, were
3 used to assure that selected ground truth points were “pure”, i.e. each selected pixel was near the
4 centre of a paddock and that the pixels were mainly from large paddocks.

6 *Multi-temporal analysis methods for EVI time series*

7 Major constraints in the use of medium to high resolution satellite imagery for estimating crop area or
8 yield are: aligning the image date with maximum crop canopy cover during the crop growth period
9 and the high costs involved in acquiring such imagery are. This is further confounded by variability in
10 climate, soil and crop practices within a specific region, making crop yield and area estimates less
11 accurate and more tedious to compute. To overcome this problem in this study, we focused on the
12 use of multiple consecutive images spanning the whole calendar year (i.e. January to December).
13 This allowed the capture of crop canopy information before, during, and after the crop growth period.

14 The efficacy of three analytical approaches to the multi-temporal data was examined: (i)
15 Clustering of multi-date MODIS EVI (MEVI) image values between day of year (DOY) 97 (early April)
16 and DOY 305 (end of October), (ii) Harmonic Analysis of the Time-series (HANTS) (Jakubauskas *et al.*
17 2001; 2002) of EVI data, and (iii) Principal Component Analysis (PCA) of the time series of EVI data.
18 The methods were assessed based on their ability to correctly classify image pixels based on field
19 observations over a period of 2 years (2003 and 2004) and the degree of association with surveyed
20 shire-scale crop area data (ABARE 2005).

21 The first approach involves classifying EVI values (see next section for details) from the
22 consecutive MODIS imagery during the main winter crop growth period, which spans from early April
23 to late October in this region. This constitutes the MEVI approach.

24 The second approach (HANTS) is based on decomposing the time series of EVI data from the
25 imagery into harmonic components or terms. In this study, for each pixel within the study area, the
26 time series encompassing 23 16-day MODIS EVI composites in each year was decomposed using a
27 discrete Fast Fourier Transform algorithm into a set of amplitude and phase terms at different

1 temporal frequencies. This technique was applied through the use of the Harmonic Analysis of Time
2 Series software (Verhoef *et al.* 1996).

3 Thirdly, the PCA approach uses traditional multivariate analysis to reduce the multidimensional
4 complexity in the temporal EVI profile. In this study, principal component analysis (Campbell 2002;
5 Davis 2002; Richards and Jia 1999) was used to reduce the EVI time series at each pixel from the 23-
6 image sequence into a smaller set of transformed variables or principal components (PC), which
7 explained 90% or more of the temporal variability in the series.

8 Finally, a benchmark (or control) classification approach was included. This was derived from a
9 single date EVI MODIS image acquired around the peak of the average EVI (PEVI) profile. In the
10 analysis, peak EVI was selected at day of year (DOY) 225.

11 *Crop/image feature classes and pixel classification*

12 For each analysis method, pixel classification was trained on ground truth data and its accuracy
13 tested on an independent set of ground truth data. Ground truth data were collated during field trips
14 undertaken in each year. In total, 1302 (wheat = 252; barley = 96; chickpea = 36; other = 918)
15 and 1365 (wheat = 243; barley = 45; chickpea = 9; other = 1068) sampling points were selected
16 from the ground truth data for the 2003 and 2004 season, respectively. Locations sampled within
17 the study area were classified according to crop/feature classes (i.e. wheat planted early, wheat
18 planted late, barley, etc.) given in Table 1. All features were identified from ground truth data
19 gathered during field trips except for the vegetation and forest classes, which were identified from
20 the 1999 land use map (Department of Natural Resources and Water 2006). The feature class
21 selections encompass classes of main interest, i.e. wheat, barley, and chickpea.

22 The ability to discriminate between crops is directly related to the amount of reflectance,
23 specifically in the NIR bandwidth, by the leaf and canopy structures (Campbell, 2002). For wheat
24 and barley, these features are very similar. The main factor contributing to differences in canopy
25 reflectance between wheat and barley relates to canopy architecture and density, which is a function
26 of the number of tillers and rate of growth. For barley, tillering and early growth are nearly double
27 that of wheat, causing more rapid crop canopy closure (Meinke *et al.* 1998). This is a significant
28 feature because discriminatory ability is likely to be associated with this attribute. The different crop

1 architecture and phenology of chickpea causes its leaf and canopy structure development to be
 2 almost in all cases quite different from wheat and barley, thus enabling discrimination between these
 3 crops.

4 The inclusion of crop feature classes or merging of specific classes was determined using
 5 separability metrics such as the Jeffries-Matusita (JM) measure. This metric constitutes the
 6 separability between two feature classes and is a function of the average distance between the
 7 spectral means of two classes. Output values range from 0 to 2.0 and indicate how well the selected
 8 feature class pairs are statistically separate. Values greater than 1.9 indicate that the feature class
 9 pairs have good separability (Richards and Jia 1999).

10

11 Table 1: Feature classes and data collating method used in the first level of
 12 classification for 2003 and 2004 seasons. *Double cropped* represents cropping in
 13 consecutive summer and winter seasons; *fed off* is traditionally hayed or grazed;
 14 *late plantings* are usually plantings occurring at the end or after the close of the
 15 traditional wheat planting window; and *na* represents no available data.

Feature Class	2003	2004
Barley	Field trip	Field trip
Barley double cropped	Field trip	na
Barley fed off	Field trip	na
Chickpeas	Field trip	Field trip
Grazing & natural vegetation	Field trip & Land use map	Field trip & Land use map
Natural forest	Land use map	Land use map
Production forest	Land use map	Land use map
Stubble & soil	Field trip	Field trip
Wheat	Field trip	Field trip
Wheat late plantings	Field trip	na

16

17

18

19

20

21

22

23

24

Supervised classification was performed via the maximum likelihood classification (MLC)
 algorithm (Richards and Jia 1999), which was available as part of the ENVI software. When only one
 layer or band was used, as in the case of PEVI approach, the minimum distance classifier (MDC)
 method was used. The classifiers (i.e. MLC and MDC) were trained using “pure” pixels within the
 ground truth data sample set (i.e. those pixels that fall completely within a large and homogeneous
 paddock for a specific feature type).

1 *Independent validation and accuracy assessment*

2 The accuracy of classification was assessed by contrasting the classified image (as described in the
3 previous section) with independent randomly selected sub-samples from the ground truthing
4 (collated through field trips). This was done to reduce artificial accuracy, i.e. minimise classification
5 bias. In total, 316 and 344 independent random ground truth pixels were selected and used to
6 calculate accuracy for the 2003 and 2004 seasons, respectively. This represented approximately
7 25% of the total ground truth samples in each year. The proportion of pixels correctly classified was
8 expressed empirically in a contingency table known as the confusion or error matrix. The statistic,
9 percent correctly classified (PCC) was used to determine the overall and between-crop accuracies for
10 each classification approach (Richards and Jia 1999). The results allowed inferences about the
11 comparative discriminatory ability of the multi-temporal decomposition approaches used in this study.

12 Accuracy at the aggregate shire-scale was determined by comparing derived estimates of total
13 and specific winter crop area with results of extended farm surveys conducted in the study region for
14 the 2003 and 2004 seasons (ABARE 2005). The degree of correspondence within a specific season at
15 a shire-scale was measured by calculating the percent error (PE). PE is computed as the ratio of the
16 difference of the remotely sensed area estimate and the surveyed area estimate to that of the
17 surveyed area estimate for each method for each year within a shire. The average of the absolute
18 PE was calculated to determine the accuracy across seasons and shires (MAPE).

19

20 **Results and discussion**

21 *Feature class selection*

22 The temporal separability between class means of wheat and wheat late plantings was moderate (JM
23 = 1.6) when the distance measures were compared. Hence, all wheat samples were merged into
24 one feature class with 252 and 243 sampling points in 2003 and 2004, respectively. Although good
25 separability was evident between barley/barley double cropped (JM = 1.99) and barley/barley fed off
26 (JM = 1.99), both barley double cropped and barley fed off were excluded from the final
27 classification. This was mainly because double cropping and fed off and haying of crops are less
28 common practice and resulted in fewer training and independent sampling points for ground truthing.

1 This resulted in 96 and 45 sampling points in 2003 and 2004, respectively. Very few chickpea sites
2 were observed and selected in either season, mainly because very little area was sown to chickpea,
3 especially in the 2004 season. Although there were few sampling points for chickpea (36 in 2003 and
4 9 in 2004) it was retained as a separate class to assess the discriminatory ability of the proposed
5 methods between the two main winter crops (i.e. wheat and barley), and the less important winter
6 crop (i.e. chickpea). The separability between barley and chickpea was larger than that between
7 wheat and chickpea. For simplicity, all other features (e.g. vegetation, natural forest, bare fallow
8 etc.) were combined to form one feature class with 918 and 1068 sampling points for both seasons.
9 In total, four main feature classes (i.e. wheat, barley, chickpea and non-cropping) were formed for
10 further analysis and classification.

11

12 *Temporal crop EVI profiles*

13 The average temporal EVI profiles throughout each growing season showed distinct differences for
14 wheat, barley and chickpea (Figure 2). The profiles represent the temporal plant canopy responses
15 to soil, plant and water regime combinations within the study area for each season. The differences
16 among crops in slope of the temporal profiles from emergence (i.e. EVI >2000 after DOY 129) to
17 anthesis (i.e. flowering around peak EVI at DOY 225) were more evident during 2003 than in 2004.
18 The period from crop emergence to anthesis is known as the green-up period while the period after
19 anthesis to crop harvest is known as the senescence period. The temporal profiles for barley and
20 wheat suggested a very similar planting date as crop emergence was around the same time for both
21 seasons for both shires (Figure 2). The average crop emergence date of chickpea was at least 2
22 months after that of wheat and barley, which suggested a later average planting date in both
23 seasons within the study area.

24

25

[Insert Figure 2 here]

26

27 The average EVI temporal profile for barley was higher than that of wheat in both seasons. In
28 addition, the green-up rate for barley was quicker than that of wheat in both seasons, which was

1 mainly an effect of the higher (i.e. nearly double) tillering and early leaf area growth rate of barley
2 (Meinke *et al.* 1998) . There were, however, some instances where the green-up rate of wheat was
3 similar to that of barley. This could be possibly ascribed to differences in soil temperatures,
4 increased nitrogen levels or no water limitations (e.g. irrigated) (Meinke *et al.* 1997). Conversely,
5 chickpea had much lower average EVI values than that of wheat and barley in both seasons. The
6 differences in average peak EVI values were not as large for the 2004 season. Although there was
7 some overlap in the temporal profile distributions between crops, the differences in the shape of the
8 profiles between wheat, barley and chickpea were apparent for both seasons. The much lower EVI
9 peaks for wheat and barley during the 2004 season were mainly caused by the significantly below
10 average rainfall recorded during 2004 that resulted in a reduction in biomass and crop growth and
11 thus ensuing lower EVI values. During periods of severe moisture stress such as in 2004, the
12 reflectance of crops in the visible (blue, green and red) bands increases (due to less absorption by
13 chlorophyll), while reflectance in the near-infrared band decreases, resulting in smaller band ratio
14 values and ensuing EVI values. Some overlaps in EVI temporal profile distributions for wheat, barley
15 and chickpea indicate that there will be some confusion in separating these crops, and consequently
16 some pixels will likely be wrongly classified.

17 *Image classification*

18 Once each method was trained on ground truth data, classification of all pixels on the image was
19 done by applying the standard maximum likelihood classifier for the multi date EVI imagery (from
20 DOY 97 to 305) and the derived PCA and HANTS imagery. The minimum distance classifier was used
21 to classify the peak EVI approach at DOY 225 (PEVI). For the PCA approach, 11 principal components
22 were retained, which explained more than 90% of the total temporal variability in the time series
23 derived from the 23 images. For the HANTS approach three harmonic terms (each term consists of a
24 phase and amplitude value) and the zero amplitude were used in the final classification. This
25 included the EVI average (0th harmonic), first, second and third harmonics (amplitude and phase for
26 each harmonic). The three harmonics plus the average explained more than 90% of the temporal
27 variability, similar to the finding for the PCA approach.

1 Figure 3 shows the classified images using the PEVI (a, b) and HANTS (c, d) approaches for the
2 2003 and the 2004 seasons, respectively. In general the two seasons differ significantly in the
3 amount of total winter crops planted. Independent of the classification approach, more winter crop
4 was evident in 2003 than in 2004. This related mainly to the poor rainfall recorded during 2004 and
5 the lack of sowing opportunities during the winter crop planting window (i.e. May to June). In
6 addition, the PEVI approach overestimated chickpea occurrence in both seasons with much of the
7 non-cropping pixels classified as chickpea in both 2003 and 2004 (a, b). The HANTS approach
8 showed substantially better discriminatory ability between wheat, barley, chickpea and non-cropping
9 than the PEVI approach in both seasons. This was due to the better discriminatory ability of the
10 HANTS approach compared to that of the single-date approach at pixel scale (Table 2). Similar
11 results to that for the HANTS approach were found for the MEVI and PCA data reduction methods
12 (data not shown).

13

14

[Insert Figure 3 here]

15

16 *Independent validation and accuracy assessment*

17 The percent of pixels correctly classified (PCC) for each of the four methods is given in Table 2.
18 The overall accuracy among these methods ranged from 56% to 98%. The single date approach
19 (PEVI) had most pixels incorrectly classified with an overall accuracy of 56% and 61% for 2003 and
20 2004 seasons, respectively. Most of this error came from misclassifying wheat and non-cropping
21 classes during both seasons. The overall PCC values for the multi-temporal approaches were all very
22 high with the highest accuracies produced in 2004. All multi-temporal approaches classified the non-
23 cropping pixels correctly (100%). This is significant because it means that such approaches can be
24 effectively used to discriminate crops from non-cropping land use areas in future studies. All multi-
25 temporal approaches achieved much higher overall accuracy compared to the single date method for
26 both the 2003 and 2004 seasons, respectively. This is mainly a result of the better ability in
27 discriminating between wheat, barley, chickpea and non-cropping in both seasons by utilising the
28 temporal canopy signatures derived from the entire crop growth period.

29

Table 2: Accuracy (%) across all classes for (i.e. wheat, barley, chickpea and non-cropping) for each method for the 2003 and 2004 seasons

Percent Correctly Classified (%)					
2003	Overall	Wheat	Barley	Chickpea	Non-cropping
Single date	56	57	90	80	51
Multi date	94	76	76	93	100
PCA	93	60	86	93	100
HANTS	93	56	95	86	100
CF1	87	58	90	100	92
CF2	85	58	86	100	89
2004					
Single date	61	74	85	100	56
Multi date	98	89	100	25	100
PCA	98	92	93	0	100
HANTS	95	85	71	0	100
CF1	95	92	71	50	97
CF2	92	89	100	25	93

Comparing the total winter crop area estimates (i.e. wheat, barley and chickpea) to the surveyed shire-scale data as collated by ABARE (Table 3), the HANTS method produced the smallest error (i.e. highest accuracy) within the Jondaryan shire for both seasons. It has an average mean absolute percent error (MAPE) of 26% (PE of 18% and -35% for each season, respectively). The MAPE across both shires was 27% (Table 4). All other methods showed MAPE greater than 63% for the Jondaryan shire (Table 3) and 97% across both shires for both seasons (Table 4). The single-date method had the smallest PE for total wheat area estimated of 5% and 9% for the Jondaryan shire for 2003 and 2004, respectively. This result, however, is fortuitous because of the very poor overall and within class pixel accuracies (Table 1). This artificial accuracy of the single-date approach is further confirmed by the very poor total winter crop shire-scale accuracy within 2003 (182%), 2004 (268%) and overall (225%) (Table 4). The high accuracy for the single-date wheat classification at an aggregated shire-scale is therefore spurious because of compensating errors when aggregating. Furthermore, the single-date approach is compounded by the question of the best date to use, which cannot be readily determined until after the season. Therefore the single date approach cannot be recommended as an acceptable method in determining winter crop area at a regional scale.

1 Table 3: Total shire-scale area estimates and ABARE surveyed (actual) data
 2 across all features (i.e. wheat, barley, chickpea and other) for each method for
 3 the 2003 and 2004 seasons within the Jondaryan shire. The accuracy is given in
 4 the PE (%) column, which is the difference between the estimated and actual
 5 values expressed as a percentage of the actual as collated by the ABARE survey
 6 (ABARE 2005).

	2003 Season			2004 Season		
	Estimate	Actual	PE (%)	Estimate	Actual	PE (%)
<i>Single date</i>						
Wheat	28597	27358	5	5922	5443	9
Barley	4566	10796	-58	1853	2714	-32
Chickpea	87110	7760	1023	18033	1650	993
Winter crop	120272	45914	162	25808	9807	163
<i>Multi date</i>						
Wheat	74502	27358	172	12417	5443	128
Barley	2865	10796	-73	7259	2714	167
Chickpea	14327	7760	85	0	1650	-100
Winter crop	91694	45914	100	19676	9807	101
<i>PCA</i>						
Wheat	72591	27358	165	8978	5443	65
Barley	2865	10796	-73	5521	2714	103
Chickpea	6877	7760	-11	0	1650	-100
Winter crop	82334	45914	79	14499	9807	48
<i>HANTS</i>						
Wheat	37824	27358	38	4909	5443	-10
Barley	2674	10796	-75	1509	2714	-44
Chickpea	13850	7760	78	0	1650	-100
Winter crop	54348	45914	18	6419	9807	-35

7
 8
 9
 10 Table 4: Aggregated temporal (2003, 2004 and All columns) scale accuracies
 11 (MAPE, %) for each of the remote sensing analysis approaches for the study area.

	2003	2004	All
Single Date			
Wheat	4	37	20
Barley	63	21	42
Chickpea	2645	1971	2308
Winter Crop	182	268	225
Multi Date			
Wheat	175	201	188
Barley	76	240	158
Chickpea	509	100	304
Winter Crop	128	172	150
PCA			
Wheat	165	116	140
Barley	68	145	106
Chickpea	171	100	135
Winter Crop	99	95	97
HANTS		HANTS	
Wheat	43	15	29
Barley	81	33	57
Chickpea	366	100	233
Winter Crop	33	21	27

1

2

3 The HANTS approach, showed moderate to high within season accuracy for total winter crop area
4 estimates, with MAPE values of 33% and 21% for the 2003 and 2004 seasons, respectively (Table 4).
5 All multi-temporal approaches showed significantly higher accuracy at the aggregated shire-scale
6 level within and across seasons compared to the accuracy of the single-date approach. The HANTS
7 method had the highest overall accuracy (27%) when determining total winter crop area estimates
8 across seasons within the study area.

9 Although the HANTS approach showed overall pixel accuracy similar to that of the other multi-
10 temporal approaches, it had the smallest total winter crop area error across both seasons and is thus
11 likely to be more reliable than any of the other analysis approaches. The shire-scale accuracy of
12 HANTS could be further increased by including ground truth data on areas that have been double
13 cropped with barley (i.e. cropping barley immediately after a summer crop). The degree of
14 discrimination between wheat and barley relates to how similar/dissimilar the temporal profile
15 trajectories are within the cropping window (Figure 2). The discriminatory ability of the HANTS
16 approach seems to be weaker during wetter seasons and stronger during the drier seasons as was
17 the case during 2003 and 2004, respectively. This weaker discriminatory ability in wet years is likely
18 to be related to spatial variability in rainfall and soil types, as well as the different crop management
19 practices, such as increased plant density rates, fertilizer application rates or a combination of these.

1 During 2004, which was classified as an El Niño year, there was less classification error between
2 wheat and barley crops, resulting in more accurate area estimates at the shire-scale. In addition,
3 almost all of the area that could be planted was planted to wheat and barley, which resulted in very
4 few ground truth fields been collated during the 2004 season. This resulted in chickpea been
5 excluded as a feature class, which further contributed to the poor discrimination of chickpea from
6 wheat, barley and other crops for the 2004 season. Thus, future studies would need a large number
7 of ground truth sampling points to enable rigorous discriminatory ability of chickpea from other
8 winter crops.

9 The temporal profile trajectory represents the crop life cycle (e.g. emergence, anthesis, maturity,
10 etc.) at a specific location and incorporates canopy reflectance responses to immediate environmental
11 conditions (i.e. temperature, soil, moisture, light, etc.). Thus, applying these multi-temporal
12 approaches to other geographical regions with soils and climate regimes not captured within the
13 study area needs further investigation.

14

15 *Implications for industry*

16 Accurate and objective crop area estimates are required along with yield estimates for accurate
17 crop production estimates. Managing storage, transport and marketing of bulk grain commodities
18 requires estimates of likely quantities throughout the production regions and with sufficient advance
19 warning for appropriate responses. The proposed remote sensing based multi-temporal analysis
20 approaches showed appreciable accuracy and are thus likely to be able to be adapted to assist
21 industry decision-making processes and enhance handling and marketing efficiencies. This will assist
22 the role of agri-business at national and international scales and should also be reflected in enhanced
23 returns to growers.

24 Annual winter crop production estimates can be created by incorporating the application of
25 remote sensing approaches, such as proposed in this study, with the end of season crop yield
26 forecast issued by QDPI&F. Although end-of-year winter crop production estimates are a significant
27 improvement on the current ABS and ABARE survey estimates (and will be of value to industry), the
28 need exists to generate crop area and production estimates that are available with the monthly crop

1 yield forecast available before harvest. This will avoid situations where an average crop yield is
2 forecast (t/ha) but little to no crop could be planted because of insufficient timely rainfall events.
3 Further research and development is necessary to address this issue of improving the lead time and
4 frequency of accurate remote sensing based crop area estimates.

5

6 **Acknowledgements**

7 We thank W Verhoef and A van der Kamp from the National Aerospace Laboratory (NLR) in the
8 Netherlands for supplying the HANTS software and providing guidance and advice on its use. We
9 also thank Land and Water Australia, through their Managing Climate Variability Program, for partly
10 funding this project.

11

1 **References**

2
3 ABARE. 2005. Australian Bureau of Agricultural Resource and Economics farm survey for
4 the 2003 and 2004. Canberra, Australia.

5
6 Department of Natural Resources and Water 2006, Queensland Land Use Mapping Program,
7 viewed May 2005, <http://www.nrm.qld.gov.au/science/lump>.

8
9 Campbell JB (2002) 'Introduction to Remote Sensing.' (Guilford Press: New York).

10
11 Davis JC (2002) 'Statistics and Data Analysis in Geology.' (John Wiley & Sons Inc.: New
12 York).

13
14 Hammer GL, Hansen JW, Phillips JG, Mjelde JW, Hill H, Love A, Pogieter AB (2001)
15 Advances in application of climate prediction in agriculture. *Agricultural Systems* **70**, 515-
16 553.

17
18 Huete A, Justice C, Liu H (1994) Development of Vegetation and Soil Indices for MODIS-
19 EOS. *Remote Sensing of Environment* **49**, 224-234.

20
21 Huete AR, Liu HQ, Batchily K, van Leeuwen W (1997) A Comparison of Vegetation Indices
22 over a Global set of TM Images for EOS-MODIS. *Remote Sensing of Environment* **59**, 440-
23 451.

24
25 Huete A, Didan K, Miura T, Rodriguez EP, Gao X, Ferreira LG (2002) Overview of the
26 radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing
27 of Environment* **83**, 195-213.

28
29 Jakubauskas ME, Legates DR, Kastens JH (2001) Harmonic Analysis of Time Series
30 AVHRR NDVI Data. *Photogrametric Engineering & Remote Sensing* **67**, 461-470.

31
32 Jakubauskas ME, Legates DR, Kastens JH (2002) Crop identification using harmonic
33 analysis of time -series AVHRR NDVI data. *Computers and Electronics in Agriculture* **37**,
34 127-139.

35
36 Justice CO, Townsend JRG, Vermote EF, Masuoka E, Wolfe RE, Saleous N, Roy DP,
37 Morisette JT (2002) An overview of MODIS Land data processing and product status.
38 *Remote Sensing of Environment* **83**, 3-15.

39
40 Knopke P, O'Donnell V, Shepherd A (2000) Productivity Growth in the Australian Grains
41 Industry. ABARE0 642 76413 1, Canberra.

42
43 Meinke H, Hammer GL, van Keulen H, Rabbinge R (1998) Improving wheat simulation
44 capabilities in Australia from a cropping systems perspective. III. The integrated wheat
45 model (I_WHEAT). *Europ. J. Agron* **8**, 101-116.

46
47 Meinke H, Hammer GL, van Keulen H, Rabbinge R, Keating BA (1997) Improving wheat
48 simulation capabilities in Australia from a cropping systems perspective. Water and nitrogen
49 effects on spring wheat in a semi-arid environment. . *Europ. J. Agron.* **7**, 75-88.

50

1 Muchoney D, Borak J, Chi M, Friedl M, Gopal S, Hodges J, Morrow N, Strahler A (2000)
2 Application of MODIS global supervised classification model to vegetation and land cover
3 mapping of Central America. *Int. J. Remote Sensing* **21**, 1115-1138.
4
5 Nelson RA, Holzworth DP, Hammer GL, Hayman PT (2002) Infusing the use of seasonal
6 climate forecasting into crop management practice in North East Australia using discussion
7 support software. *Agricultural Systems* **74**, 393-414.
8
9 Nix HA (1975) The Australian Climate and its Effects on Grain Yield and Quality. In
10 'Australian Field Crops'. (Eds A Lazenby, EM Matheson) pp. 183-226. (Angus and
11 Robertson: Sydney, Australia).
12
13 Penm J (2002) Economic Overview. ABARE Volume 9, Number 4, Canberra.
14
15 Potgieter AB, Hammer GL, Butler D (2002) Spatial and temporal patterns in Australian
16 wheat yield and their relationship with ENSO. *Australian Journal of Agricultural Research*
17 **53**, 77-89.
18
19 Potgieter AB, Hammer GL, deVoil P (2005) A simple regional-scale model for forecasting
20 sorghum yield across North-Eastern Australia. *Agriculture and Forest Meteorology* **132**, 143-
21 153.
22
23 Potgieter AB, Hammer GL, Doherty A (2006) Oz-Wheat: a regional-scale crop yield
24 simulation model for Australian wheat. Queensland Department of Primary Industries &
25 Fisheries. Information Series No, QI06033, Brisbane, Australia. (ISSN 0727-6273).
26
27 Price JC (2003) Comparing MODIS and ETM+ data for regional and global land
28 classification. *Remote Sensing of Environment* **86**, 491-499.
29
30 Richards JA, Jia X (1999) 'An Introduction Remote Sensing Digital Image Analysis.'
31 (Springer-Verlag: Berlin Heidelberg).
32
33 RSI (2005) Research Systems Inc., www.rsinc.com
34
35 Stephens DJ (1998) Objective criteria for estimating the severity of drought in the wheat
36 areas of Australia. *Agricultural Systems* **57**, 333-350.
37
38 Stephens DJ, Butler D, Hammer GL (2000) Using seasonal climate forecasts in forecasting
39 the Australian wheat crop. In 'Applications of Seasonal Climate Forecasting in Agriculture and
40 Natural Ecosystems: The Australian Experience'. (Eds GL Hammer, N Nicholls, C Mitchell)
41 pp. 351-366. (Kluwer Academic Publishers: Dordrecht, Netherlands).
42
43 Stone RC, Hammer GL, Marcussen T (1996) Prediction of global rainfall probabilities using
44 phases of the Southern Oscillation Index. *Nature* **384**, 252-255.
45
46 Verhoef W, Menenti M, Azzali S (1996) A colour composite of NOAA-AVHRR-NDVI
47 based on time series analysis (1981-1992). *Int. J. Remote Sensing* **17**, 231-235.
48

- 1 Xiao X, Boles S, Liu J, Zhuang D, Froking S, Li C, Salas W, Moore III B (2005) Mapping
2 paddy rice agriculture in southern China using multi-temporal MODIS images. *Remote*
3 *Sensing of Environment* **95**, 480-492.
4
- 5 Zhan X, Sohlberg RA, Townsend JRG, DiMiceli C, Carroll ML, Eastman JC, Hansen MC,
6 DeFries RS (2002) Detection of land cover changes using MODIS 250 m data. *Remote*
7 *Sensing of Environment* **83**, 336-350.
8
- 9 Zhang X, Friedl MA, Schaaf CB, Strahler AH, Hodges JCF, Gao F, Reed C, Huete A (2003)
10 Monitoring vegetation phenology using MODIS. *Remote Sensing of Environment* **84**, 471-
11 475.

1 Figure Captions:
2

3 Figure 1: Location of the Jondaryan and Pittsworth shires (hatched in black) within the north
4 eastern region of Australia. Shire boundaries are given by black solid lines.


5
6 Figure 2: Average temporal EVI profile throughout the growing season for wheat (green,
7 square), barley (brown, triangle) and chickpea (yellow, diamond) for (a) 2003 winter crop
8 season and (b) 2004 winter crop season.

9
10 Figure 3: Classified images using the PEVI classification for the 2003 and 2004 seasons (a, b)
11 and classified images using the HANTS approach for 2003 and 2004 seasons respectively (c,
12 d). Wheat is coloured in green, barley in yellow, chickpea in cyan and non crop (e.g. natural
13 and production forest, vegetation, stubble, bare soil etc.) in brown.

14
15

1 Figures:

2




3

4 Figure 1

5

6



7


8 Figure 2

9


10

11

1



2



3

4 Figure 3