

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

| | |
|-----------|--|
| Title | Metabolomic profiling of rhodosporidium toruloides grown on glycerol for carotenoid production during different growth phases(Main article) |
| Author(s) | Lee, Jaslyn Jie Lin; Chen, Liwei; Shi, Jiahua; Trzcinski, Antoine; Chen, Wei Ning |
| Citation | Lee, J. J. L., Chen, L., Shi, J., Trzcinski, A., & Chen, W.-N. (2014). Metabolomic profiling of rhodosporidium toruloides grown on glycerol for carotenoid production during different growth phases. Journal of agricultural and food chemistry, 62(41), 10203-10209. |
| Date | 2014 |
| URL | http://hdl.handle.net/10220/25205 |
| Rights | © 2014 American Chemical Society. This is the author created version of a work that has been peer reviewed and accepted for publication by Journal of Agricultural and Food Chemistry, American Chemical Society. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [http://dx.doi.org/10.1021/jf502987q]. |

Metabolomic Profiling of *Rhodospiridium toruloides* Grown on Glycerol for Carotenoid Production During Different Growth Phases


Jaslyn Jie Lin Lee[†], Liwei Chen[‡], Jiahua Shi[‡], Antoine Trzcinski[†] and Wei-Ning Chen^{*†,‡}

[†] Advanced Environmental Biotechnology Centre, Nanyang Environment and Water Research Institute, Nanyang Technological University, 1 Cleantech Loop, Singapore 637141

[‡] School of Chemical and Biomedical Engineering, Nanyang Technological University, 62 Nanyang Drive, Singapore 637459

*corresponding author: Tel. (W.-N.C.) Phone: +65 63162870. Fax: +65 67947553. E-mail: WNChen@ntu.edu.sg

Carotenoid production from three strains of *Rhodospiridium toruloides* grown on glycerol was studied. A time-dependent metabolomics approach was used to understand its metabolism on glycerol and mechanism for carotenoid production in three strains, during different growth phases (1, 4, 7, and 12 days). The strain CBS 5490 was the highest carotenoid producer (28.5 mg/L) and had a unique metabolic profile. In this strain, metabolites belonging to the TCA cycle and amino acids were produced in lower amounts, as compared to the other strains. On the other hand, it produced the highest amounts of carotenoid and fatty acid metabolites. This indicated that the lower production of the TCA cycle and amino acid metabolites was to promote energy and metabolic flux toward the carotenoid and fatty acid synthesis metabolic pathways. This study shows that metabolomic profiling is a useful tool to gain insight into the metabolic pathways in the cell, and to shed light on the different molecular mechanisms between strains.



Keywords: Metabolomics; glycerol; *R. toruloides*; carotenoids; GC-MS

SI File: jf502987q_si_001.pdf

Introduction

Glycerol is a widely available product due to the increased demand for biodiesel production worldwide. For every gallon of biodiesel produced, 1.05 pounds of glycerol is produced as a side product. It is predicted that the biodiesel industry would produce 37 billion gallons of biodiesel by 2016. As a result, the price of glycerol has decreased to \$0.30 per pound.¹ Thus, this has generated a wide-spread interest in using glycerol as a low-cost substrate for microbial fermentation to produce value-added compounds.²


Rhodospiridium toruloides is a red yeast that is gaining interest for its robust ability to grow on a variety of waste substrates.³⁻⁶ It is used to produce valuable compounds such as cephalosporin esterase, phenylalanine ammonia-lyase, and epoxide hydrolase.⁷⁻⁹ In recent years, there is a growing body of literature on this yeast for its ability to produce the valuable metabolic products carotenoids and fatty acids.^{6,10,11}

Carotenoids are a group of colored pigments. There are over 700 different types of carotenoids. They occur naturally in plants, yeasts, and algae. Demand for microbial sources of carotenoids has risen tremendously, and they are preferred over plant sources, for their fast growing rates. The most well studied carotenoid is β -carotene, a pro-vitamin A molecule, which possesses antioxidant properties that are beneficial for health, and is used as a nutritional

supplement and food colorant.¹² The β -carotene market potential is estimated to be worth over USD U.S. \$280 million in 2015. The current industrial producer of β -carotene is from the microalga microalga *Dunaliella salina* *Dunaliella salina*. However carotenoids, carotenoid production from algae is generally expensive and requires large areas for cultivation;³ hence, there is increasing demand for studies of other microorganisms and methods of cultivation.¹³

R. toruloides produces the carotenoids: torularhodin, **1**;³ torulene, **2**;³ and β -carotene, **3** (Figure 1).¹⁴ All three carotenoids have applications as food colorants and poultry feed.¹⁵⁻¹⁵ Apart from β -carotene, other lesser known carotenoids were also studied for their antioxidant activities. Torulene, **2**, was found to have more antioxidant potential than β -carotene, due to its higher number of double bonds.^{16,17} Torularhodin, **1**, may be a more potent singlet oxygen quencher than β -carotene, due to its longer polyene chain.

Figure 1. The structure Structures and absorption maxima (λ_{\max}) of the carotenoids: torularhodin, **1**;³ torulene, **2**;³ and β -carotene, **3**.



The emerging Foodomicsfoodomics trend uses a mass spectrophotometry (MS) based approach to advance food science and nutrition, for its fast and high -throughput nature.¹⁸ Under the foodomics field, untargeted metabolomics is an analytical strategy to identify all metabolites in an organism under a given set of conditions.¹⁹ This use of metabolomics -rangeranges from analyzing wine profiles -to developfor the development of a predictive model to aid wine development by nontargeted GC-MS,²⁰ to studying microbial metabolism,²¹ and is also used to identify the main compound in strawberry responsible for its anti-inflammatory benefits.²²

The ability of *R. toruloides* to use glycerol as a substrate for fermentation has been well explored.^{6,23,24} Its proteomic profile has also been studied in lipid -accumulating conditions using LC-MS-based metabolomic technology.¹¹ However, studies on the metabolism of this unique yeast on glycerol, and its mechanism to convert glycerol into carotenoids -isare lacking. In this study we analyze the potential of carotenoids carotenoid production from three strains grown on glycerol using HPLC, and analyzed their metabolic changes at four different growth phases via GC-MS. The differential metabolite profile would also be useful to understand the differences in -carotenoids carotenoid production between the strains.

Materials and Methods

Fermentation of *R. toruloides*

Over the past years, many studies have shown that that the yeast *R. toruloides* can use glycerol as the sole carbon source for fermentation. From over a decade of work on this species, we have chosen three strains: *R. toruloides* ATCC 10788 (AC) is from the American Type Culture Collection as itane was one of the first -strainstrains to be investigated for its lipid -producing ability,²⁵ and AS 2.1389 (AS) is from the China General Microbiological Culture Collection Center -sinceand its genome was recently sequenced, which enables future genetic work.¹¹ To both of these strains we added CBS 5490 (CBS), which was purchased from the largest fungal culture collection worldwide, (Central Bureau voor Schimmelcultures, Utrecht, The Netherlands) -as; it is a novel strain -whichthat has not yet been studied. All strains were maintained on YPG agar plates containing: (per liter) 20 g of glycerol, 20 g of peptone, 10 g of yeast extract, and 20 g of agar. Overnight culture was used to inoculate each 250 mL Erlenmeyer flask at an optical density (OD) of

0.2 containing 50 mL of fermentation media. Fermentation media contained (per liter) 20 g of peptone, 10 g of yeast extract, and 60 g of glycerol, purity = 99% (Sigma, St. Louis, MO, USA). Fermentation was carried out in an incubator for 12 days in 250 mL Erlenmeyer flasks at 30 °C, with shaking at 250 rpm. Three biological replicates were carried out for each strain.

Growth and Carotenoids Extraction

To measure the growth of the yeasts, the OD of the culture media was measured at a wavelength of 600 nm, using a Nano Drop 2000 UV spectrophotometer (Thermo Scientific, Waltham, MA, USA). Dry cell weight was quantitated by drying 1 mL of washed culture supernatant in an oven set at 70 °C until constant dry cell weight (DCW). Biomass was quantitated by using a standard curve correlating OD at 600 nm to dry cell weight. To analyze the intracellular carotenoid content of the cells, 1 mL of culture media was taken. The cells were centrifuged at 10,000 × g, washed with Milli-Q water, and broken with glass beads using a Fast Prep bead grinder (MP Biomedicals, Solon, OH, USA). Carotenoids were extracted with acetone until the cell pellet was colorless. Carotenoids were measured and quantitated using standards from CaroteNature (Ostermundigen, Switzerland). The spectrophotometric measurement of carotenoids was according to their absorption maxima peak (λ_{max}), which is the middle maximum of their UV-vis spectra in acetone. They were measured as follows: torularhodin, **1**, at 495 nm; torulene, **2**, at 485 nm; and β -carotene, **3**, at 455 nm.

Carotenoids Characterization and Quantitation

To confirm the quantitation and characterization of carotenoids extracted in acetone in each strain, high performance liquid chromatography (HPLC) equipped with a photodiode array detector was used as it is a validated method.²⁶ The carotenoids torularhodin, (**1**), torulene, (**2**), and β -carotene, (**3**), were identified based on the basis of their retention times using standards from CaroteNature (Ostermundigen, Switzerland). An Agilent 1100 Series instrument was employed, and the column used was a 250 mm × 4.6 mm i.d., 5 μ m, Li-Chrospher 100 RP-18, with a 4 mm × 4 mm i.d. guard column of the same material (Merck, Rahway, NJ, USA). A gradient from 70 to 100% acetone in Milli-Q water at a flow rate of 0.5 mL/min was used as the mobile phase. Detection was performed at 450 nm, and the UV-vis absorption spectra were recorded online with the photodiode array detection system.

Extracellular Glycerol Analysis

To analyze extracellular glycerol levels in the culture media, 1 mL of culture medium was taken every 24 h and filtered. Glycerol was quantitated using HPLC equipped with a refractive index detector using a standard of purity of 99% (Sigma, St. Louis, USA). The column used was an Aminex HPX-87H column (Bio-Rad, Hercules, CA, USA). Milli-Q water was used as the mobile phase at a flow rate of 0.6 mL/min, with temperature controlled at 65 °C, and detection was performed at 210 nm.

Metabolic Profile of each Strain at Different Growth Phases and GC-MS Analysis

The intracellular metabolites were extracted from all three strains at four different time points: early log growth phase on day 1, middle of log growth phase on day 4, as cells enter stationary phase on day 7, and during death phase at the end of fermentation on day 12. Cells suspensions with an OD of 10 were lysed and quenched in 800 μ L of ice-cold methanol using glass beads and a bead grinder. Ten microliters of 2 mg/mL ribitol dissolved in Milli-Q water was added to each sample as an internal standard to correct for any loss of metabolite during the extraction process. The supernatant was extracted and then air-dried using a heat block set at 30 °C overnight. Next, 50 μ L of 20 mg/mL solution of methoxyamine hydrochloride in pyridine was added to protect the carbonyl groups and incubated at 37 °C for 1 h. Silylation was then carried out by adding 99 μ L of *N*-methyl-*N*-(trimethylsilyl)-trifluoroacetamide (MSTFA) with 1 μ L of trimethylchlorosilane (TMCS) and incubated for 30 min at 70 °C.


Samples were then shaken vigorously for 2 h at room temperature before transferring them to sample vials, and covered with Parafilm to prevent evaporation. Metabolite analysis was carried out using gas chromatography coupled with a mass spectrophotometer (GC-MS). All samples for GC-MS were run within 24 h. One microliter of sample was injected using a Shimadzu QP 2010 Plus GC-MS system. The oven temperature was kept at 75 °C for 4 min, and was raised at 4 °C/min, to a final temperature of 280 °C, and held for 2 min. Mass spectra were recorded from *m/z* 35 to 600 with a scan time of 0.2 s.²⁷ The total elution time for mass spectra collected for each sample was 58 min. Data processing and metabolite identification was carried out accordingly.²⁸

Results and Discussion

Growth of Three *R. toruloides* Strains on Glycerol

We have found 60 g/L to be the optimized glycerol concentration to ensure growth and carotenoid production in the three strains (data not shown). Glycerol is able to enter the cell through diffusion and/or active transport.²⁹ Extracellular glycerol in the culture media was depleted from the culture media by day 3 for strains AC and AS, and by day 4 for strain CBS. This showed that the three strains could assimilate glycerol and grew comparatively well (Figure 2). From the biomass curve of the three strains in Figure 2, we define that all three strains were in the logarithmic (log) phase of growth from days 1 and 6, entered stationary phase on day 7, and were in the death phase by day 12. Cells underwent division and replication by obtaining energy from glycerol through glycolysis. The best strain to use glycerol as a substrate for cell growth was CBS. The cells produced the highest concentration of biomass (42.3 g/L) during late stationary phase, as compared to 37.2 g/L by AC and 38.4 g/L by AS (Figure 2).

Figure 2. Biomass concentration, extracellular glycerol levels, and the spectrophotometric quantitation of carotenoids: torularhodin, (1), torulene, (2), and β -carotene, (3) of all the three strains. The error bar indicates the SD.



Carotenoids quantitation Carotenoid Quantitation by UV-spectrophotometer Spectrophotometer

Rhodospiridium yeasts are known to produce the carotenoids: torularhodin, (1); torulene, (2); and β -carotene, (3).¹⁴ In the three strains, carotenoid concentration reached a plateau during stationary growth phase (Figure 2). The strain CBS was the highest carotenoid producer of all strains, for the three types of carotenoids: torularhodin, (1); torulene, (2); and β -carotene, (3).

Characterization and quantitation Quantitation of carotenoids Carotenoids by HPLC

To our knowledge, there has been no report of the characterization of carotenoids in these strains. It was found that the carotenoid values obtained by spectrophotometry and HPLC greatly differed, so the values quantitated by HPLC were taken as it is a more accurate method.³⁰ The milli-absorbance units (mAU) obtained from the HPLC profile were quantitated using carotene standards. Our analysis showed that all strains produced torularhodin, (1), torulene, (2), and β -carotene, (3) in different percentages (Figure 3), although torularhodin, (1) was the most abundant carotenoid in all three strains. All of the carotenoids eluted within 24 to 28 min as shown in Table 1. The HPLC results confirmed the UV spectrophotometer findings that CBS was the highest carotenoid producer. As quantitated from HPLC, at the end of fermentation on day 12, it produced 28.5 mg/L, as compared to 18 mg/L by AC and 19.3 mg/L by AS. It also produced the highest amount of the industrially relevant compound β -carotene, (3) (Figure 3).

Figure 3. Representative HPLC profile of carotenoids in the three strains, AC, AS, and CBS. In each HPLC profile the following carotenoids were identified: torularhodin, (1); torulene, (2); and β -carotene, (3).




Table 1. Production of Carotenoids by the Three Strains Quantitated by HPLC

| Carotenoid | Ret-time (min) | Strain | | | | | |
|----------------------|----------------|--------|--------------------------|------|--------------------------|------|--------------------------|
| | | AC | | AS | | CBS | |
| carotenoid | ret time (min) | mg/L | (% of total carotenoids) | mg/L | (% of total carotenoids) | mg/L | (% of total carotenoids) |
| Torularhodin, 1 | 24.5 | 7.4 | 41 | 12.5 | 65 | 19.7 | 69 |
| Torulenone, 2 | 27.3 | 7 | 39 | 2.5 | 13 | 2.0 | 7 |
| β -carotene, 3 | 28.0 | 3.6 | 20 | 4.3 | 22 | 6.8 | 24 |
| Total | | 18 | | 19.3 | | 28.5 | |

Comparison of the metabolites profiles during different growth phases. Metabolomics was used to investigate and shed light into the differences in carotenoid accumulation in the three strains on a metabolic level. The changes in intermediates would help to explain the cellular metabolism

accounting for the different carotenoids production in the three strains. In total, 13 important differential metabolites were identified, and the changes in their levels are shown in Figures 5 and 6. The metabolites could be classified into carbohydrates, TCA cycle intermediates, fatty acids, and amino acids. To apply the knowledge of metabolomic changes seen in the cell, we have mapped the changes in relation to biochemical pathways, as depicted in Figure 4. This gives a view of the metabolite changes in comparison to the other strains AC and AS, which could help us to understand the reason for higher carotenoid accumulation in strain CBS.

Figure 4. Metabolic changes in the strain CBS mapped onto metabolic pathways during the four different growth phases on days 1, 4, 7, and 12. Coloured symbols represent an increase, decrease, or no change in the level of the metabolite at each time point, as compared to the other strains AC and AS. Orange arrows represent the general flow of metabolic flux in strain CBS.




Figure 5. Quantitated metabolite profiling in the three strains AC, AS, and CBS during different growth phases by GC-MS: changes in the carbohydrate metabolite glycerol and amino acid levels of serine, glycine, threonine, alanine, glutamine, and proline. The error bar indicates the SD of three biological replicates.





Figure 6. Quantitated metabolite profiling in the three strains AC, AS, and CBS during different growth phases by GC-MS. Changes: changes in the TCA cycle intermediate metabolites malic acid and succinic acid, and levels of the total intracellular fatty acids metabolites (oleic acid, stearic acid, linoleic acid, and palmitic acid). The error bar indicates the SD of three biological replicates.



Carbohydrates metabolites Carbohydrate Metabolites

Glycerol was used as the sole carbon source in this study. In the cell, glycolysis is the pathway which supplies the cells with energy, by breaking down sugars such as glycerol, to form the final pyruvate molecule. During glycolysis, glycerol is converted into glyceraldehyde-3-phosphate, transformed into phosphoenolpyruvate (PEP), and finally into pyruvate (Figure 4). Pyruvate may enter the mitochondria and is converted into acetyl-CoA where it, which enters the citric acid cycle (TCA). Acetyl-CoA is an important molecule as it is the precursor of the fatty acid and carotenoid synthesis pathway.³¹ In Figure 5, the intracellular metabolite glycerol was found to decrease in all three strains in comparatively the same manner, as glycolysis took place.

Amino acids metabolites Acid Metabolites

Amino acids can be synthesized from compounds in the glycolysis pathway and TCA cycle. The glycolysis pathway provides the substrate glyceraldehyde-3-phosphate, for the synthesis of threonine, serine, and glycine, which are used to form branched chain amino acids (Figure 4). The amino acid alanine is synthesized from the pyruvate molecule provided by the glycolysis pathway. Glutamate and proline are synthesized from the intermediate of the TCA cycle, α -ketoglutarate (α -KG).

Interestingly, we observe that the strain CBS demonstrated a unique amino acid metabolic profile. In comparison to AC and AS strains, all of the amino acids were generally synthesized in lower amounts (Figure 5). This might suggest that the cellular mechanism of the CBS strain preferably synthesized less amino acids, which resulted in more available acetyl-CoA to enter the carotenoid and fatty acid synthesis pathways. This could contribute to the higher levels of carotenoids and total intracellular fatty acids observed in strain CBS.

TCA cycle intermediate metabolites Cycle Intermediate Metabolites

The TCA cycle is a key metabolic pathway in all cells. After glycolysis, pyruvate enters the mitochondria, and is converted into acetyl-CoA, which undergoes oxidative phosphorylation to provide energy for the cell in the form of NADH and NADPH. The TCA cycle contains intermediates such as succinic acid and malic acid.

In strain CBS we also observed a general down-regulation in production of the TCA cycle intermediates, malic acid and succinic acid, as compared to strains AC and AS (Figure 6). Therefore, we propose that the TCA cycle was down-regulated, to enable more acetyl-CoA from glycolysis to enter the carotenoid and fatty acid synthesis pathway. This contributes to higher carotenoid production in CBS strain CBS. This is also seen in a previous study wherein which the decrease of the TCA cycle intermediate malic acid, enhanced β -carotene production in the yeast *Blakeslea trispora*.³²

Fatty acid metabolites Acid Metabolites

R. toruloides is an oleaginous yeast which has been studied for its ability to accumulate high amounts of lipids. The total intracellular fatty acids synthesized by all three strains was palmitic acid, oleic acid, linoleic acid, and stearic acid, which is in accordance with previous results.²³ The highest carotenoid-producing strain, CBS, was also the highest fatty acid producer, especially the unsaturated oleic acid and linoleic acid (Figure 6).

The cell membrane is made up of a phospholipid bilayer where many biochemical processes take place. Both fatty acids and carotenoids are localized in the lipid bilayer membrane. Carotenoid-lipid interactions play a role in the regulation of membrane fluidity and integrity. β -carotene molecules are randomly integrated in the hydrophobic interior of the lipid bilayer. Its ability to penetrate deeper into the membrane is dependent on the fluidity of the membrane. Higher amounts of unsaturated fatty acids in the bilayer result in increased membrane fluidity, thereby allowing more β -carotene to accumulate deeper into the interior of the bilayer.³³ Therefore, the high content of the unsaturated fatty acids, oleic acid and linoleic acid, in strain CBS, may have served not to increase the membrane stability, but improved to improve the cell membrane fluidity to enhance the accumulation of all three carotenoids. In support of this, a study by Li et al.³⁴ has also shown that oleic acid could help increase membrane fluidity to overcome ethanol stress in *Saccharomyces cerevisiae*.

In addition, intracellular fatty acids form TAG and sterol esters, which are stored inside lipid bodies.³⁵ The increased synthesis of total intracellular fatty acid metabolites observed in strain CBS means more lipid bodies can be formed. Oleic acid has been suggested to be the main constituent in the formation of lipid bodies in *Dunaliella*.³⁶ It has been proposed that formation of lipid bodies was the driving force of β -carotene synthesis. This is because lipid bodies have the ability to sequester β -carotene molecules, stimulating more β -carotene production.³⁷ In light of our results it is possible that the higher production of fatty acids, especially the oleic acid levels in strain CBS, resulted in more lipid bodies formed, which can sequester all three carotenoids including torularhodin, 1-; torulene, 2-; and β -carotene, 3. The sequestering by lipid bodies thus promoted an increase for overall total carotenoid accumulation which was not reported previously. Therefore, our study found a strong correlation between fatty acid

production to carotenoids and carotenoid biosynthesis, especially in the unsaturated fatty acids oleic acid, which could help increase membrane fluidity and more lipid bodies body formation.

Carotenoids synthesis pathway Carotenoid Synthesis Pathway

Isoprenoids such as carotenoids are secondary metabolites produced at highest levels during late log phase to early stationary phase,³⁸ which is consistent with our observations. As seen in Figure 4, in the carotenoid synthesis pathway, γ -carotene is the metabolic branch point for β -carotene and the torulene/torularhodin pathway. From γ -carotene, cyclization forms the dicyclic carotenoid β -carotene, while whereas oxidation takes place to form the monocyclic carotenoid torulene, 2, and further oxidation into torularhodin, 1.³⁹ In all strains the highly oxidized end product torularhodin, 1, was the major carotenoid formed (Table 1). The regulatory mechanisms controlling the carotenoid synthesis pathways are not well understood, and further studies are required.

In summary, the use of a metabolomics strategy to study the metabolic profile at different time points has allowed us to gain insight into the metabolism and biochemical pathways of *R. toruloides*. Also, it helped us understand the underlying differences in carotenoid-producing ability between strains. The results suggested that fatty acid synthesis was beneficial to increase carotenoid production. However, more studies on the relationship between fatty acids and carotenoid synthesis is needed. Knowledge from this study could aid future work to increase carotenoid yields by metabolic engineering in microorganisms. For example, to down-regulate the TCA cycle and amino acid synthesis, and increase the unsaturated fatty acid production such as oleic acid.

The authors declare no competing financial interest.

Abbreviations Used

Strain ATCC, 10788 strain AC; Strain AS, 2.1389 strain AS; Strain CBS, 5490 strain CBS; BCAA, branched-chain amino acid; α -KG, α -ketoglutarate; OAA, oxaloacetate; PEP, phosphoenolpyruvate

Acknowledgments. Strain AS 2.1389 was a kind gift from Professor Zongbao (Kent) Zhao from Dalian Institute of Chemical Physics in China. We thank NEWRI for support.

Supporting Information

Supporting information

Supplementary Table S1 lists the retention times of all differential metabolites identified using the NIST library. Figure S1 shows the total ion chromatogram from GC-MS analysis of all three strains on d 12. Figure S2 shows the neutral lipid content in each strain at the end of fermentation by Nile red staining. Figure S3 shows the analysis of total nitrogen in the culture media during fermentation. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Yang, F.; Hanna, M.; Sun, R. Value-added uses for crude glycerol—a byproduct of biodiesel production. *Biotechnol. Biofuels* **2012**, *5*, 1310.1186/1754-6834-5-13.
- (2) Dobson, R.; Gray, V.; Rumbold, K. Microbial utilization of crude glycerol for the production of value-added products. *J. Ind. Microbiol. Biotechnol.* **2012**, *39*, 217–226.
- (3) Zhao, X.; Peng, F.; Du, W.; Liu, C.; Liu, D. Effects of some inhibitors on the growth and lipid accumulation of oleaginous yeast *Rhodospiridium toruloides* and preparation of biodiesel by enzymatic transesterification of the lipid. *Bioprocess Biosyst. Eng.* **2012**, *35*, 993–1004.
- (4) Wang, Q.; Guo, F. J.; Rong, Y. J.; Chi, Z. M. Lipid production from hydrolysate of cassava starch by *Rhodospiridium toruloides* 21167 for biodiesel making. *Renewable Energy* **2012**, *46*, 164–168.
- (5) Zhao, X.; Wu, S.; Hu, C.; Wang, Q.; Hua, Y.; Zhao, Z. K. Lipid production from Jerusalem artichoke by *Rhodospiridium toruloides* Y4. *J. Ind. Microbiol. Biotechnol.* **2010**, *37*, 581–585.
- (6) Saenge, C.; Cheirsilp, B.; Suksaroge, T. T.; Bourtoom, T. Potential use of oleaginous red yeast *Rhodotorula glutinis* for the bioconversion of crude glycerol from biodiesel plant to lipids and carotenoids. *Process Biochem.* **2011**, *46*, 210–218.
- (7) Politino, M.; Tonzi, S. M.; Burnett, W. V.; Romancik, G.; Usher, J. J. Purification and characterization of a cephalosporin esterase from *Rhodospiridium toruloides*. *Appl. Environ. Microbiol.* **1997**, *63*, 4807–4811.
- (8) Zhu, L.; Cui, W.; Fang, Y.; Liu, Y.; Gao, X.; Zhou, Z. Cloning, expression and characterization of phenylalanine ammonia-lyase from *Rhodotorula glutinis*. *Biotechnol. Lett.* **2013**, *35*, 751–756.

- (9) Botes, A. L. Affinity purification and characterization of a yeast epoxide hydrolase. *Biotechnol. Lett.* **1999**, *21*, 511–517.
- (10) Li, Y.; Zhao, Z.; Bai, F. High-density cultivation of oleaginous yeast *Rhodospiridium toruloides* Y4 in fed-batch culture. *Enzyme Microb. Technol.* **2007**, *41*, 312–317.
- (11) Zhu, Z.; Zhang, S.; Liu, H.; Shen, H.; Lin, X.; Yang, F.; Zhou, Y. J.; Jin, G.; Ye, M.; Zou, H.; Zhao, Z. K.; A multi-omic map of the lipid-producing yeast *Rhodospiridium toruloides*. *Nat. Comm. Commun.* **2012**, *3*, DOI: 10.1038/ncomms2112.
- (12) Ribeiro, B. D.; Barreto, D. W.; Coelho, M. A. Z. Technological aspects of β -carotene production. *Food Bioprocess Technol.* **2011**, *4*, 693–701.
- (13) Guedes, A. C.; Amaro, H. M.; Malcata, F. X. Microalgae as sources of carotenoids. *Mar. Drugs* **2011**, *9*, 625–44644.
- (14) Buzzini, P.; Innocenti, M.; Turchetti, B.; Libkind, D.; van Broock, M.; Mulinacci, N. Carotenoid profiles of yeasts belonging to the genera *Rhodotorula*, *Rhodospiridium*, *Sporobolomyces*, and *Sporidiobolus*. *Can. J. Microbiol.* **2007**, *53*, 1024–311031.
- (15) Buzzini, P. Production of yeast carotenoids by using agro-industrial by-products. *Agro Food Ind. Hi-Tech* **2001**, *12*, 7–10.
- (16) Sakaki, H.; Nakanishi, T.; Tada, A.; Miki, W.; Komemushi, S. Activation of torularhodin production by *Rhodotorula glutinis* using weak white light irradiation. *J. Biosci. Bioeng.* **2001**, *92*, 294–297.
- (17) Sakaki, H.; Nochide, H.; Komemushi, S.; Miki, W. Effect of active oxygen species on the productivity of torularhodin by *Rhodotorula glutinis* no. 21. *J. Biosci. Bioeng.* **2002**, *93*, 338–340.
- (18) Capozzi, F.; Bordoni, A. Foodomics: A new comprehensive approach to food and nutrition. *Genes Nutri Nutr.* **2013**, *8*, 1–4.
- (19) Herrero, M.; Simão, C.; García-Cañas, V.; Ibáñez, E.; Cifuentes, A. Foodomics: MS-based strategies in modern food science and nutrition. *Mass Speet Spectrom. Rev.* **2012**, *31*, 49–69.
- (20) Schmidtke, L. M.; Blackman, J. W.; Clark, A. C.; Grant-Preece, P. Wine metabolomics: objective measures of sensory properties of semillon Semillon from GC-MS profiles. *J. Agri Agric. Food Chem.* **2013**, *61*, 11957–11967.
- (21) Cevallos-Cevallos, J. M.; Reyes-De-Corcuera, J. I.; Etxeberria, E.; Danyluk, M. D.; Rodrick, G. E. Metabolomic analysis in food science: a review. *Trends Food Sci. Technol.* **2009**, *20*, 557–566.
- (22) Lee, J.; Kim, S.; Namgung, H.; Jo, Y.-H.; Bao, C.; Choi, H.-K.; Auh, J.-H.; Lee, H. J. Ellagic acid identified through metabolomic analysis is an active metabolite in strawberry ('seolhyang') regulating lipopolysaccharide-induced inflammation. *J. Agric. Food Chem.* **2013**, *62*, 3954–3962.
- (23) Xu, J.; Zhao, X.; Wang, W.; Du, W.; Liu, D. Microbial conversion of biodiesel byproduct glycerol to triacylglycerols by oleaginous yeast *Rhodospiridium toruloides* and the individual effect of some impurities on lipid production. *Biochem. Eng. J.* **2012**, *65*, 30–36.
- (24) Uckun Kiran, E.; Trzcinski, A.; Webb, C. Microbial oil produced from biodiesel by-products could enhance overall production. *Bioresour. Technol.* **2013**, *129*, 650–4654.
- (25) Turcotte, G.; Kosaric, N. The effect of C/N ratio on lipid production by *Rhodospiridium toruloides* ATCC 10788. *Biotechnol. Lett.* **1989**, *11*, 637–642.
- (26) Kimura, M.; Kabori, C. N.; Rodriguez-Amaya, D. B.; Nestel, P. Screening and HPLC methods for carotenoids in sweetpotato, cassava and maize for plant breeding trials. *Food Chem.* **2007**, *100*, 1734–1746.
- (27) Wang, M.; Bai, J.; Chen, W. N.; Ching, C. B. Metabolomic profiling of cellular responses to carvedilol enantiomers in vascular smooth muscle cells. *PLoS ONE One* **2010**, 510.1371/journal.pone.0015441.
- (28) Seiber, J. N.; Molyneux, R. J.; Schieberle, P. Targeted metabolomics: a new section in the journal of agricultural and food chemistry. *J. Agric. Food Chem.* **2013**, *62*, 22–23.
- (29) Chatzifragkou, A.; Dietz, D.; Komaitis, M.; Zeng, A. P.; Papanikolaou, S. Effect of biodiesel-derived waste glycerol impurities on biomass and 1,3-propanediol production of *Clostridium butyricum* VPI 1718. *Biotechnol. Bioeng.* **2010**, *107*, 76–84.
- (30) Rodriguez-Amaya, D. B.; Kimura, M.; Institute, I. F. P. R.; Tropical, C. I. d. A.; HarvestPlus Handbook for Carotenoid Analysis; International Food Policy Research Institute (IFPRI): Washington, DC, USA, 2004.
- (31) Fakas, S.; Makri, A.; Bellou, S.; Aggelis, G. Pathways to aerobic glycerol catabolism and their regulation. *Microbial Conversions of Raw Glycerol* **2009**, 9–18. Fakas, S.; Makri, A.; Bellou, S.; Aggelis, G. Pathways

- to aerobic glycerol catabolism and their regulation. In *Microbial Conversions of Raw Glycerol*; Nova Science Publishers: New York, 2009; pp 9–18.
- (32) Hu, X.; Li, H.; Tang, P.; Sun, J.; Yuan, Q.; Li, C. GC-MS-based metabolomics study of the responses to arachidonic acid in *Blakeslea trispora*. *Fungal Gen. Biol.* **2013**, *57*, 33–41.
- (33) Popova, A. V.; Andreeva, A. S. Carotenoid-Lipid Interactions lipid interactions. *Adv. Planar Lipid Bilayers and Liposomes* **2013**, *17*, 215–236.
- (34) Li, Z.; Ma, X.; Li, A.; Zhang, C. A novel potential source of β -carotene: *Eustigmatos cf. polyphem* (Eustigmatophyceae) and pilot β -carotene production in bubble column and flat panel photobioreactors. *Bioresour. Technol.* **2012**, *117*, 257–263.
- (35) Fujimoto, T.; Ohsaki, Y.; Cheng, J.; Suzuki, M.; Shinohara, Y. Lipid droplets: a classic organelle with new outfits. *Histochem. Cell Biol.* **2008**, *130*, 263–279.
- (36) Lamers, P. P.; Janssen, M.; De Vos, R. C. H.; Bino, R. J.; Wijffels, R. H. Carotenoid and fatty acid metabolism in nitrogen-starved *Dunaliella salina*, a unicellular green microalga. *J. Biotechnol.* **2012**, *162*, 21–27.
- (37) Rabbani, S.; Beyer, P.; Lintig, J. V.; Huguency, P.; Kleinig, H. Induced β -carotene synthesis driven by triacylglycerol deposition in the unicellular alga *Dunaliella bardawil*. *Plant Physiol.* **1998**, *116*, 1239–1248.
- (38) Siddiqui, M. S.; Thodey, K.; Trenchard, I.; Smolke, C. D. Advancing secondary metabolite biosynthesis in yeast with synthetic biology tools. *FEMS Yeast Res.* **2012**, *12*, 144–170.
- (39) Sperstad, S.; Lutnaes, B. F.; Stormo, S. K.; Liaaen-Jensen, S.; Landfald, B. Torularhodin and torulene are the major contributors to the carotenoid pool of marine *Rhodospiridium babjevae* (Golubev). *J. Ind. Microbiol. BiotechBiotechnol.* **2006**, *33*, 269–273.